

# Tina Kyndt

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4917499/publications.pdf>

Version: 2024-02-01

96  
papers

4,444  
citations

87723

38  
h-index

118652

62  
g-index

100  
all docs

100  
docs citations

100  
times ranked

3513  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Spatiotemporal expression profile of novel and known small RNAs throughout rice plant development focussing on seed tissues. <i>BMC Genomics</i> , 2022, 23, 44.  | 1.2 | 4         |
| 2  | Rice diterpenoid phytoalexins are involved in defence against parasitic nematodes and shape rhizosphere nematode communities. <i>New Phytologist</i> , 2022, 235, 1231-1245.  | 3.5 | 12        |
| 3  | Dehydroascorbate induces plant resistance in rice against root-knot nematode <i>Meloidogyne graminicola</i> . <i>Molecular Plant Pathology</i> , 2022, 23, 1303-1319.   | 2.0 | 13        |
| 4  | Genetic disruption of <i>Arabidopsis</i> secondary metabolite synthesis leads to microbiome-mediated modulation of nematode invasion. <i>ISME Journal</i> , 2022, 16, 2230-2241.  | 4.4 | 9         |
| 5  | Plant defense priming in the field: a review. , 2021, , 87-124.   |     | 9         |
| 6  | Genome-wide shifts in histone modifications at early stage of rice infection with <i>Meloidogyne graminicola</i> . <i>Molecular Plant Pathology</i> , 2021, 22, 440-455.  | 2.0 | 14        |
| 7  | Chitin in Strawberry Cultivation: Foliar Growth and Defense Response Promotion, but Reduced Fruit Yield and Disease Resistance by Nutrient Imbalances. <i>Molecular Plant-Microbe Interactions</i> , 2021, 34, 227-239. | 1.4 | 19        |
| 8  | Benzoxazinoids selectively affect maize root-associated nematode taxa. <i>Journal of Experimental Botany</i> , 2021, 72, 3835-3845.   | 2.4 | 15        |
| 9  | The phenylpropanoid pathway inhibitor piperonylic acid induces broad-spectrum pest and disease resistance in plants. <i>Plant, Cell and Environment</i> , 2021, 44, 3122-3139.  | 2.8 | 31        |
| 10 | The Induced Resistance Lexicon: Do <sup>TM</sup> s and Don <sup>TM</sup> ts. <i>Trends in Plant Science</i> , 2021, 26, 685-691.  | 4.3 | 84        |
| 11 | Phytohormones selectively affect plant parasitic nematodes associated with <i>Arabidopsis</i> roots. <i>New Phytologist</i> , 2021, 232, 1272-1285.   | 3.5 | 11        |
| 12 | Non-coding RNAs in the interaction between rice and <i>Meloidogyne graminicola</i> . <i>BMC Genomics</i> , 2021, 22, 560.   | 1.2 | 12        |
| 13 | Plant parasitic cyst nematodes redirect host indole metabolism via NADPH oxidase-mediated ROS to promote infection. <i>New Phytologist</i> , 2021, 232, 318-331.  | 3.5 | 9         |
| 14 | Biochar-Enhanced Resistance to <i>Botrytis cinerea</i> in Strawberry Fruits (But Not Leaves) Is Associated With Changes in the Rhizosphere Microbiome. <i>Frontiers in Plant Science</i> , 2021, 12, 700479.            | 1.7 | 11        |
| 15 | Induced Resistance by Ascorbate Oxidation Involves Potentiating of the Phenylpropanoid Pathway and Improved Rice Tolerance to Parasitic Nematodes. <i>Frontiers in Plant Science</i> , 2021, 12, 713870.                | 1.7 | 8         |
| 16 | Short-term effects of cadmium on leaf growth and nutrient transport in rice plants. <i>Plant Science</i> , 2021, 313, 111054.   | 1.7 | 15        |
| 17 | Beneficial worm allies warn plants of parasite attack belowground and reduce aboveground herbivore preference and performance. <i>Molecular Ecology</i> , 2021, , .   | 2.0 | 5         |
| 18 | Cucurbitaceae COLD Peeling Extracts (CCOPEs) Protect Plants From Root-Knot Nematode Infections Through Induced Resistance and Nematicidal Effects. <i>Frontiers in Plant Science</i> , 2021, 12, 785699.                | 1.7 | 4         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Vitamin C in Plants: Novel Concepts, New Perspectives, and Outstanding Issues. <i>Antioxidants and Redox Signaling</i> , 2020, 32, 463-485.   | 2.5 | 84        |
| 20 | The Use of PTI-Marker Genes to Identify Novel Compounds that Establish Induced Resistance in Rice. <i>International Journal of Molecular Sciences</i> , 2020, 21, 317.  | 1.8 | 16        |
| 21 | Ascorbate Oxidase Induces Systemic Resistance in Sugar Beet Against Cyst Nematode <i>Heterodera schachtii</i> . <i>Frontiers in Plant Science</i> , 2020, 11, 591715.   | 1.7 | 22        |
| 22 | A Phytochemical Perspective on Plant Defense Against Nematodes. <i>Frontiers in Plant Science</i> , 2020, 11, 602079.   | 1.7 | 43        |
| 23 | Chorismate mutase and isochorismatase, two potential effectors of the migratory nematode <i>Hirschmanniella oryzae</i> , increase host susceptibility by manipulating secondary metabolite content of rice. <i>Molecular Plant Pathology</i> , 2020, 21, 1634-1646.                               | 2.0 | 12        |
| 24 | Molecular insights into the compatible and incompatible interactions between sugar beet and the beet cyst nematode. <i>BMC Plant Biology</i> , 2020, 20, 483.   | 1.6 | 21        |
| 25 | Evaluation of Metabarcoding Primers for Analysis of Soil Nematode Communities. <i>Diversity</i> , 2020, 12, 388.  | 0.7 | 20        |
| 26 | Genome-wide DNA hypomethylation shapes nematode pattern-triggered immunity in plants. <i>New Phytologist</i> , 2020, 227, 545-558.  | 3.5 | 44        |
| 27 | Root-knot nematodes induce gall formation by recruiting developmental pathways of post-embryonic organogenesis and regeneration to promote transient pluripotency. <i>New Phytologist</i> , 2020, 227, 200-215.   | 3.5 | 41        |
| 28 | Ascorbate oxidation activates systemic defence against root-knot nematode <i>Meloidogyne graminicola</i> in rice. <i>Journal of Experimental Botany</i> , 2020, 71, 4271-4284.  | 2.4 | 26        |
| 29 | Systemic defense activation by COS-OGA in rice against root-knot nematodes depends on stimulation of the phenylpropanoid pathway. <i>Plant Physiology and Biochemistry</i> , 2019, 142, 202-210.  | 2.8 | 45        |
| 30 | Selection of miRNA reference genes for plant defence studies in rice ( <i>Oryza sativa</i> ). <i>Planta</i> , 2019, 250, 2101-2110.   | 1.6 | 9         |
| 31 | Strigolactones enhance root-knot nematode ( <i>Meloidogyne graminicola</i> ) infection in rice by antagonizing the jasmonate pathway. <i>New Phytologist</i> , 2019, 224, 454-465.  | 3.5 | 47        |
| 32 | Jasmonate-Induced Defense Mechanisms in the Belowground Antagonistic Interaction Between <i>Pythium arrhenomanes</i> and <i>Meloidogyne graminicola</i> in Rice. <i>Frontiers in Plant Science</i> , 2019, 10, 1515.  | 1.7 | 15        |
| 33 | Gibberellin antagonizes jasmonate-induced defense against <i>Meloidogyne graminicola</i> in rice. <i>New Phytologist</i> , 2018, 218, 646-660.  | 3.5 | 71        |
| 34 | Mechanisms of resistance in the rice cultivar Manikpukha to the rice stem nematode <i>Ditylenchus angustus</i> . <i>Molecular Plant Pathology</i> , 2018, 19, 1391-1402.  | 2.0 | 22        |
| 35 | Trace analysis of multi-class phytohormones in <i>Oryza sativa</i> using different scan modes in high-resolution Orbitrap mass spectrometry: method validation, concentration levels, and screening in multiple accessions. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 4527-4539. | 1.9 | 28        |
| 36 | Trichoderma-Inoculated Miscanthus Straw Can Replace Peat in Strawberry Cultivation, with Beneficial Effects on Disease Control. <i>Frontiers in Plant Science</i> , 2018, 9, 213.   | 1.7 | 28        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | <i>Meloidogyne graminicola</i> : a major threat to rice agriculture. <i>Molecular Plant Pathology</i> , 2017, 18, 3-15.   | 2.0 | 134       |
| 38 | Below-Ground Attack by the Root Knot Nematode <i>Meloidogyne graminicola</i> Predisposes Rice to Blast Disease. <i>Molecular Plant-Microbe Interactions</i> , 2017, 30, 255-266.  | 1.4 | 28        |
| 39 | Transcriptomic and histological responses of African rice ( <i>Oryza glaberrima</i> ) to <i>Meloidogyne graminicola</i> provide new insights into root-knot nematode resistance in monocots. <i>Annals of Botany</i> , 2017, 119, 885-899.                      | 1.4 | 54        |
| 40 | Interplay between Carotenoids, Abscisic Acid and Jasmonate Guides the Compatible Rice- <i>Meloidogyne graminicola</i> Interaction. <i>Frontiers in Plant Science</i> , 2017, 8, 951.  | 1.7 | 58        |
| 41 | Interactions between the oomycete <i>Pythium arrhenomanes</i> and the rice root-knot nematode <i>Meloidogyne graminicola</i> in aerobic Asian rice varieties. <i>Rice</i> , 2016, 9, 36.  | 1.7 | 9         |
| 42 | Redirection of auxin flow in <i>Arabidopsis thaliana</i> roots after infection by root-knot nematodes. <i>Journal of Experimental Botany</i> , 2016, 67, 4559-4570.   | 2.4 | 69        |
| 43 | Thiamine-induced priming against root-knot nematode infection in rice involves lignification and hydrogen peroxide generation. <i>Molecular Plant Pathology</i> , 2016, 17, 614-624.  | 2.0 | 54        |
| 44 | Identification of Bangladeshi rice varieties resistant to ufra disease caused by the nematode <i>Ditylenchus angustus</i> . <i>Crop Protection</i> , 2016, 79, 162-169.   | 1.0 | 6         |
| 45 | Analysis of fungal endophytes associated with rice roots from irrigated and upland ecosystems in Kenya. <i>Plant and Soil</i> , 2016, 405, 371-380.   | 1.8 | 23        |
| 46 | Biochar-amended potting medium reduces the susceptibility of rice to root-knot nematode infections. <i>BMC Plant Biology</i> , 2015, 15, 267.   | 1.6 | 92        |
| 47 | A high-resolution melt (HRM) assay to characterize CYP51 haplotypes of the wheat pathogen <i>Mycosphaerella graminicola</i> . <i>Crop Protection</i> , 2015, 71, 12-18.   | 1.0 | 7         |
| 48 | The role of thionins in rice defence against root pathogens. <i>Molecular Plant Pathology</i> , 2015, 16, 870-881.  | 2.0 | 33        |
| 49 | Î <sup>2</sup> -Aminobutyric Acid-induced Resistance Against Root-Knot Nematodes in Rice Is Based on Increased Basal Defense. <i>Molecular Plant-Microbe Interactions</i> , 2015, 28, 519-533.  | 1.4 | 75        |
| 50 | Recent Advances in Understanding Plant-Nematode Interactions in Monocots. <i>Advances in Botanical Research</i> , 2015, 73, 189-219.  | 0.5 | 8         |
| 51 | The genome of cultivated sweet potato contains <i>Agrobacterium</i> T-DNAs with expressed genes: An example of a naturally transgenic food crop. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 5844-5849. | 3.3 | 236       |
| 52 | Systemic Suppression of the Shoot Metabolism upon Rice Root Nematode Infection. <i>PLoS ONE</i> , 2014, 9, e106858.   | 1.1 | 13        |
| 53 | Vasconcellea for Papaya Improvement. , 2014, , 47-79.   |     | 15        |
| 54 | Analysis of the transcriptome of <i>Hirschmanniella oryzae</i> to explore potential survival strategies and host-nematode interactions. <i>Molecular Plant Pathology</i> , 2014, 15, 352-363.   | 2.0 | 23        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Plant-Parasitic Nematode Infections in Rice: Molecular and Cellular Insights. Annual Review of Phytopathology, 2014, 52, 135-153.  | 3.5 | 123       |
| 56 | Sensitivity towards DMI fungicides and haplotypic diversity of their CYP51 target in the <i>Mycosphaerella graminicola</i> population of Flanders. Journal of Plant Diseases and Protection, 2014, 121, 156-163. | 1.6 | 6         |
| 57 | Nematode feeding sites: unique organs in plant roots. Planta, 2013, 238, 807-818.  | 1.6 | 158       |
| 58 | Identification of candidate effector genes in the transcriptome of the rice root knot nematode <i>Meloidogyne graminicola</i> . Molecular Plant Pathology, 2013, 14, 379-390.                                    | 2.0 | 69        |
| 59 | Brassinosteroids Suppress Rice Defense Against Root-Knot Nematodes Through Antagonism With the Jasmonate Pathway. Molecular Plant-Microbe Interactions, 2013, 26, 106-115.                                       | 1.4 | 118       |
| 60 | How far a protected area contributes to conserve habitat species composition and population structure of endangered African tree species (Benin, West Africa). Ecological Complexity, 2013, 13, 60-68.           | 1.4 | 40        |
| 61 | Transcriptional analysis through RNA sequencing of giant cells induced by <i>Meloidogyne graminicola</i> in rice roots. Journal of Experimental Botany, 2013, 64, 3885-3898.                                     | 2.4 | 128       |
| 62 | Transcriptional silencing of RNAi constructs against nematode genes in Arabidopsis. Nematology, 2013, 15, 519-528.   | 0.2 | 12        |
| 63 | An insight into critical endocycle genes for plant-parasitic nematode feeding sites establishment. Plant Signaling and Behavior, 2013, 8, e24223.  | 1.2 | 12        |
| 64 | Transcriptome analysis of rice mature root tissue and root tips in early development by massive parallel sequencing. Journal of Experimental Botany, 2012, 63, 2141-2157.  | 2.4 | 41        |
| 65 | <i>CCS52</i> and <i>DEL1</i> genes are key components of the endocycle in nematode-induced feeding sites. Plant Journal, 2012, 72, 185-198.  | 2.8 | 75        |
| 66 | Abscisic acid interacts antagonistically with classical defense pathways in rice-migratory nematode interaction. New Phytologist, 2012, 196, 901-913.  | 3.5 | 120       |
| 67 | Transcriptional reprogramming by root knot and migratory nematode infection in rice. New Phytologist, 2012, 196, 887-900.  | 3.5 | 157       |
| 68 | Genetic Evidence of the Contribution of Ethnic Migrations to the Propagation and Persistence of the Rare and Declining Scrambling Shrub <i>Caesalpinia bonduc</i> L. Human Ecology, 2012, 40, 117-128.           | 0.7 | 7         |
| 69 | Comparing systemic defence-related gene expression changes upon migratory and sedentary nematode attack in rice. Plant Biology, 2012, 14, 73-82.   | 1.8 | 76        |
| 70 | Vasconcellea, 2011, 213-249.   |     | 22        |
| 71 | Quantitative morphological descriptors confirm traditionally classified morphotypes of <i>Tamarindus indica</i> L. fruits. Genetic Resources and Crop Evolution, 2011, 58, 299-309.                              | 0.8 | 36        |
| 72 | Natural variation in fruit characteristics, seed germination and seedling growth of <i>Adansonia digitata</i> L. in Benin. New Forests, 2011, 41, 113-125.   | 0.7 | 44        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | The Jasmonate Pathway Is a Key Player in Systemically Induced Defense against Root Knot Nematodes in Rice. <i>Plant Physiology</i> , 2011, 157, 305-316.  | 2.3 | 318       |
| 74 | Nematode Resistant GM Crops in Industrialised and Developing Countries. , 2011, , 517-541.  |     | 35        |
| 75 | The Role of Pseudo-Endoglucanases in the Evolution of Nematode Cell Wall-Modifying Proteins. <i>Journal of Molecular Evolution</i> , 2010, 70, 441-452.   | 0.8 | 20        |
| 76 | Women's Traditional Knowledge, Use Value, and the Contribution of Tamarind ( <i>Tamarindus indica</i> L.) to Rural Households' Cash Income in Benin. <i>Economic Botany</i> , 2010, 64, 248-259.  | 0.8 | 65        |
| 77 | Analysis of ITS of the rDNA to infer phylogenetic relationships among Vietnamese Citrus accessions. <i>Genetic Resources and Crop Evolution</i> , 2010, 57, 183-192.  | 0.8 | 16        |
| 78 | Genetic diversity, population structure and taxonomy of <i>Calopteryx splendens</i> (Odonata): Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542 Td  | 1.2 | 8         |
| 79 | Spatial genetic structuring of baobab ( <i>Adansonia digitata</i> ), Malvaceae) in the traditional agroforestry systems of West Africa. <i>American Journal of Botany</i> , 2009, 96, 950-957.  | 0.8 | 45        |
| 80 | Expressed sequence tags of the peanut pod nematode <i>Ditylenchus africanus</i> : The first transcriptome analysis of an Anguinid nematode. <i>Molecular and Biochemical Parasitology</i> , 2009, 167, 32-40.                           | 0.5 | 50        |
| 81 | Genetic fingerprinting using AFLP cannot distinguish traditionally classified baobab morphotypes. <i>Agroforestry Systems</i> , 2009, 75, 157-165.  | 0.9 | 44        |
| 82 | Occurrence of DNA methylation in <i>Daphnia magna</i> and influence of multigeneration Cd exposure. <i>Environment International</i> , 2009, 35, 700-706.   | 4.8 | 87        |
| 83 | Folk Classification, Perception, and Preferences of Baobab Products in West Africa: Consequences for Species Conservation and Improvement. <i>Economic Botany</i> , 2008, 62, 74-84.  | 0.8 | 130       |
| 84 | <i>AtCDKA1</i> silencing in <i>Arabidopsis thaliana</i> reduces reproduction of sedentary plant-parasitic nematodes. <i>Plant Biotechnology Journal</i> , 2008, 6, 749-757.   | 4.1 | 22        |
| 85 | A family of GHF5 endo- $\beta$ -glucanases in the migratory plant-parasitic nematode <i>Radopholus similis</i> . <i>Plant Pathology</i> , 2008, 57, 581-590.  | 1.2 | 36        |
| 86 | Evolution of GHF5 endoglucanase gene structure in plant-parasitic nematodes: no evidence for an early domain shuffling event. <i>BMC Evolutionary Biology</i> , 2008, 8, 305.   | 3.2 | 50        |
| 87 | Purification and characterization of the cysteine proteinases in the latex of <i>Vasconcellea</i> spp.. <i>FEBS Journal</i> , 2007, 274, 451-462.   | 2.2 | 23        |
| 88 | Cross-species microsatellite amplification in <i>Vasconcellea</i> and related genera and their use in germplasm classification. <i>Genome</i> , 2006, 49, 786-798.  | 0.9 | 18        |
| 89 | Evidence of Natural Hybridization and Introgression between <i>Vasconcellea</i> Species (Caricaceae) from Southern Ecuador Revealed by Chloroplast, Mitochondrial and Nuclear DNA Markers. <i>Annals of Botany</i> , 2006, 97, 793-805. | 1.4 | 38        |
| 90 | Patterns of Genetic and Morphometric Diversity in Baobab ( <i>Adansonia digitata</i> ) Populations Across Different Climatic Zones of Benin (West Africa). <i>Annals of Botany</i> , 2006, 97, 819-830.                                 | 1.4 | 110       |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 91 | Molecular phylogeny and evolution of Caricaceae based on rDNA internal transcribed spacers and chloroplast sequence data. <i>Molecular Phylogenetics and Evolution</i> , 2005, 37, 442-459.              | 1.2 | 39        |
| 92 | Isolation and characterization of microsatellite loci in the highland papaya <i>Vasconcellea</i> × <i>heilbornii</i> V. Badillo (Caricaceae). <i>Molecular Ecology Notes</i> , 2005, 5, 590-592.         | 1.7 | 4         |
| 93 | Maternal inheritance of cytoplasmic organelles in intergeneric hybrids of <i>Carica papaya</i> L. and <i>Vasconcellea</i> spp. (Caricaceae Dumort., Brassicales). <i>Euphytica</i> , 2005, 143, 161-168. | 0.6 | 14        |
| 94 | Species relationships in the genus <i>Vasconcellea</i> (Caricaceae) based on molecular and morphological evidence. <i>American Journal of Botany</i> , 2005, 92, 1033-1044.                              | 0.8 | 44        |
| 95 | Phylogenetic analysis of the highland papayas ( <i>Vasconcellea</i> ) and allied genera (Caricaceae) using PCR-RFLP. <i>Theoretical and Applied Genetics</i> , 2004, 108, 1473-1486.                     | 1.8 | 76        |
| 96 | AFLP analysis of genetic relationships among papaya and its wild relatives (Caricaceae) from Ecuador. <i>Theoretical and Applied Genetics</i> , 2002, 105, 289-297.                                      | 1.8 | 88        |