

# Nathan C Winn

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/4921695/nathan-c-winn-publications-by-year.pdf>

**Version:** 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

33  
papers

357  
citations

11  
h-index

18  
g-index

39  
ext. papers

526  
ext. citations

4.2  
avg, IF

3.77  
L-index

#	Paper	IF	Citations
33	Exercise and Adipose Tissue Immunity: Outrunning Inflammation. <i>Obesity</i> , <b>2021</b> , 29, 790-801	8	2
32	The Tailgate Study: Differing metabolic effects of a bout of excessive eating and drinking. <i>Alcohol</i> , <b>2021</b> , 90, 45-55	2.7	2
31	Voluntary Wheel Running Partially Compensates for the Effects of Global Estrogen Receptor- $\alpha$ Knockout on Cortical Bone in Young Male Mice. <i>International Journal of Molecular Sciences</i> , <b>2021</b> , 22,	6.3	2
30	Myeloid-specific deletion of ferroportin impairs macrophage bioenergetics but is disconnected from systemic insulin action in adult mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , <b>2021</b> , 321, E376-E391	6	1
29	Recruitment and remodeling of peridroplet mitochondria in human adipose tissue. <i>Redox Biology</i> , <b>2021</b> , 46, 102087	11.3	1
28	Global estrogen receptor- $\alpha$ knockout has differential effects on cortical and cancellous bone in aged male mice. <i>Facets</i> , <b>2020</b> , 5, 328-348	2.3	3
27	Regulation of tissue iron homeostasis: the macrophage "ferrostat". <i>JCI Insight</i> , <b>2020</b> , 5,	9.9	44
26	Post Meal Exercise May Lead to Transient Hypoglycemia Irrespective of Glycemic Status in Humans. <i>Frontiers in Endocrinology</i> , <b>2020</b> , 11, 578	5.7	
25	Impaired insulin signaling in the B10.D2- $\alpha$ /oSnJ mouse model of complement factor 5 deficiency. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , <b>2019</b> , 317, E200-E211	6	
24	A Thermogenic-Like Brown Adipose Tissue Phenotype Is Dispensable for Enhanced Glucose Tolerance in Female Mice. <i>Diabetes</i> , <b>2019</b> , 68, 1717-1729	0.9	8
23	Glycemic effects following the consumption of mixed soy protein isolate and alginate beverages in healthy adults. <i>Food and Function</i> , <b>2019</b> , 10, 1718-1725	6.1	1
22	Overproduction of endothelin-1 impairs glucose tolerance but does not promote visceral adipose tissue inflammation or limit metabolic adaptations to exercise. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , <b>2019</b> , 317, E548-E558	6	4
21	A thermogenic-like brown adipose tissue phenotype is dispensable for enhanced glucose tolerance in female mice. <i>FASEB Journal</i> , <b>2019</b> , 33, lb564	0.9	
20	Post Meal Hypoglycemia With and Without Exercise in Non-Obese and Obese Individuals.. <i>Medicine and Science in Sports and Exercise</i> , <b>2019</b> , 51, 139-140	1.2	
19	Metabolic Implications of Diet and Energy Intake during Physical Inactivity. <i>Medicine and Science in Sports and Exercise</i> , <b>2019</b> , 51, 995-1005	1.2	7
18	Estrogen receptor- $\beta$ signaling maintains immunometabolic function in males and is obligatory for exercise-induced amelioration of nonalcoholic fatty liver. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , <b>2019</b> , 316, E156-E167	6	17
17	Energy-matched moderate and high intensity exercise training improves nonalcoholic fatty liver disease risk independent of changes in body mass or abdominal adiposity - A randomized trial. <i>Metabolism: Clinical and Experimental</i> , <b>2018</b> , 78, 128-140	12.7	58

16	Removal of interscapular brown adipose tissue increases aortic stiffness despite normal systemic glucose metabolism in mice. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , <b>2018</b> , 314, R584-R597	3.2	16
15	Increased susceptibility to OVX-associated metabolic dysfunction in UCP1-null mice. <i>Journal of Endocrinology</i> , <b>2018</b> ,	4.7	5
14	Regular exercise reduces adipose tissue inflammation and improves glycemic control in Western diet-fed mice despite hyperendothelinemia. <i>FASEB Journal</i> , <b>2018</b> , 32, lb570	0.9	
13	Weight maintenance diets prevent short-term physical inactivity-induced glycemic dysregulation in young healthy subjects. <i>FASEB Journal</i> , <b>2018</b> , 32, 724.10	0.9	
12	Endothelial dysfunction occurs independently of adipose tissue inflammation and insulin resistance in ovariectomized Yucatan miniature-swine. <i>Adipocyte</i> , <b>2018</b> , 7, 35-44	3.2	1
11	When gain is greater than loss: effects of physical activity on insulin sensitivity after short-term inactivity in older subjects. <i>Journal of Physiology</i> , <b>2018</b> , 596, 5071-5072	3.9	
10	Loss of UCP1 exacerbates Western diet-induced glycemic dysregulation independent of changes in body weight in female mice. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , <b>2017</b> , 312, R74-R84	3.2	32
9	Plasma Irisin Modestly Increases during Moderate and High-Intensity Afternoon Exercise in Obese Females. <i>PLoS ONE</i> , <b>2017</b> , 12, e0170690	3.7	29
8	Exercise Normalizes Dysfunctional Adipose Tissue Phenotype in FGF21-Null Mice. <i>Medicine and Science in Sports and Exercise</i> , <b>2017</b> , 49, 1028	1.2	
7	Anti-inflammatory effects of exercise training in adipose tissue do not require FGF21. <i>Journal of Endocrinology</i> , <b>2017</b> , 235, 97-109	4.7	15
6	Deletion of UCP1 enhances ex vivo aortic vasomotor function in female but not male mice despite similar susceptibility to metabolic dysfunction. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , <b>2017</b> , 313, E402-E412	6	12
5	Prior exercise does not alter the incretin response to a subsequent meal in obese women. <i>Peptides</i> , <b>2015</b> , 71, 94-9	3.8	11
4	Walking Reduces Postprandial Insulin Secretion in Obese Adolescents Consuming a High-Fructose or High-Glucose Diet. <i>Journal of Physical Activity and Health</i> , <b>2015</b> , 12, 1153-61	2.5	6
3	Postdinner resistance exercise improves postprandial risk factors more effectively than predinner resistance exercise in patients with type 2 diabetes. <i>Journal of Applied Physiology</i> , <b>2015</b> , 118, 624-34	3.7	48
2	Moderate amounts of fructose- or glucose-sweetened beverages do not differentially alter metabolic health in male and female adolescents. <i>American Journal of Clinical Nutrition</i> , <b>2014</b> , 100, 796-805	7	27
1	Multomics reveals persistence of obesity-associated immune cell phenotypes in adipose tissue during weight loss and subsequent weight regain		2