## Fu Zhao

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4926721/publications.pdf Version: 2024-02-01



**ΕΠ 2ΗΛΟ** 

#	Article	IF	CITATIONS
1	Treatment and outcome of pediatric intracranial ependymoma after first relapse. Journal of Neuro-Oncology, 2022, , 1.	1.4	1
2	Novel patientâ€derived xenograft and cell line models for therapeutic screening in <scp>NF2</scp> â€associated schwannoma. Journal of Pathology, 2022, 257, 620-634.	2.1	3
3	Clinical risk factor analysis of bilateral vestibular schwannoma's growth pattern inconsistency in individual NF2 patients. Clinical Neurology and Neurosurgery, 2021, 200, 106365.	0.6	0
4	Survival and Prognostic Factors of Adult Intracranial Ependymoma. American Journal of Surgical Pathology, 2021, 45, 979-987.	2.1	11
5	Loss of 5-Hydroxymethylcytosine as an Epigenetic Signature That Correlates With Poor Outcomes in Patients With Medulloblastoma. Frontiers in Oncology, 2021, 11, 603686.	1.3	4
6	Establishment of nomograms for the prediction of useful hearing loss in patients with neurofibromatosis type 2. Journal of Neuro-Oncology, 2021, 155, 373-381.	1.4	1
7	Posterior fossa ependymoma with preoperative cerebrospinal metastases: a case report with literature review. British Journal of Neurosurgery, 2020, , 1-4.	0.4	1
8	The clinical treatment and outcome of cerebellopontine angle medulloblastoma: a retrospective study of 15 cases. Scientific Reports, 2020, 10, 9769.	1.6	7
9	Clinical features of newly developed NF2 intracranial meningiomas through comparative analysis of pediatric and adult patients. Clinical Neurology and Neurosurgery, 2020, 194, 105799.	0.6	8
10	Characterization of global 5-hydroxymethylcytosine in pediatric posterior fossa ependymoma. Clinical Epigenetics, 2020, 12, 19.	1.8	4
11	Selective impairment of the executive attentional network in adult patients with neurofibromatosis type 1. NeuroReport, 2019, 30, 921-926.	0.6	5
12	Comparison of Clinical, Histopathological, and Genomic Features Between Malignant Peripheral Nerve Sheath Tumors and Cellular Schwannomas of the Eighth Cranial Nerve: A Case Series. World Neurosurgery, 2019, 122, e487-e497.	0.7	4
13	Targeting the cMET pathway augments radiation response without adverse effect on hearing in NF2 schwannoma models. Proceedings of the National Academy of Sciences of the United States of America, 2018, 115, E2077-E2084.	3.3	32
14	Prognostic value of Ki-67 index in adult medulloblastoma after accounting for molecular subgroup: a retrospective clinical and molecular analysis. Journal of Neuro-Oncology, 2018, 139, 333-340.	1.4	9
15	Awake craniotomy for assisting placement of auditory brainstem implant in NF2 patients. Acta Oto-Laryngologica, 2018, 138, 548-553.	0.3	5
16	Surgical treatment of large vestibular schwannomas in patients with neurofibromatosis type 2: outcomes on facial nerve function and hearing preservation. Journal of Neuro-Oncology, 2018, 138, 417-424.	1.4	14
17	Deregulation of the Hippo Pathway Promotes Tumor Cell Proliferation Through YAP Activity in Human Sporadic Vestibular Schwannoma. World Neurosurgery, 2018, 117, e269-e279.	0.7	14
18	Distinctive localization and MRI features correlate of molecular subgroups in adult medulloblastoma. Journal of Neuro-Oncology, 2017, 135, 353-360.	1.4	23

Fu Zhao

#	Article	IF	CITATIONS
19	A Retrospective Analysis of Vision-Impairing Tumors Among 467 Patients with Neurofibromatosis Type 2. World Neurosurgery, 2017, 97, 557-564.	0.7	5
20	Clinical Features and Prognostic Factors of Primary Intracranial Malignant Fibrous Histiocytoma: A Report of 8 Cases and a Literature Review. Translational Neuroscience and Clinics, 2016, 2, 155-164.	0.1	0
21	Granular cell tumors in the central nervous system: a report on eight cases and a literature review. British Journal of Neurosurgery, 2016, 30, 611-618.	0.4	13
22	Molecular subgroups of adult medulloblastoma: a long-term single-institution study. Neuro-Oncology, 2016, 18, 982-990.	0.6	75
23	Clinical features of clear cell meningioma: a retrospective study of 36 cases among 10,529 patients in a single institution. Acta Neurochirurgica, 2016, 158, 67-76.	0.9	32
24	Clinical features of spinal schwannomas in 65 patients with schwannomatosis compared with 831 with solitary schwannomas and 102 with neurofibromatosis Type 2: a retrospective study at a single institution. Journal of Neurosurgery: Spine, 2016, 24, 145-154.	0.9	58
25	Post-Traumatic Cerebrospinal Fluid Rhinorrhea Associated with Craniofacial Fibrous Dysplasia: Case Report and Literature Review. Translational Neuroscience and Clinics, 2016, 2, 188-194.	0.1	1
26	Multiple Large Dilated Virchow-Robin Spaces in a 12-Year-Old With Neurofibromatosis Type 2. Pediatric Neurology, 2014, 51, 856-857.	1.0	4
27	Altered Regional and Circuit Resting-State Activity Associated with Unilateral Hearing Loss. PLoS ONE, 2014, 9, e96126.	1.1	54