
## Malin Barman

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4941381/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Maternal characteristics and pregnancy outcomes in the NICE birth cohort: an assessment of self-selection bias. Journal of Maternal-Fetal and Neonatal Medicine, 2024, 35, 9014-9022.	0.7	10
2	Metabolomic profiles of mid-trimester amniotic fluid are not associated with subsequent spontaneous preterm delivery or gestational duration at delivery. Journal of Maternal-Fetal and Neonatal Medicine, 2022, 35, 2054-2062.	0.7	4
3	Low Concentration of Fecal Valeric Acid at 1 Year of Age Is Linked with Eczema and Food Allergy at 13 Years of Age: Findings from a Swedish Birth Cohort. International Archives of Allergy and Immunology, 2022, 183, 398-408.	0.9	16
4	Differences between Arterial and Venous Umbilical Cord Plasma Metabolome and Association with Parity. Metabolites, 2022, 12, 175.	1.3	6
5	Assessment of Joint Impact of Iodine, Selenium, and Zinc Status on Women's Third-Trimester Plasma Thyroid Hormone Concentrations. Journal of Nutrition, 2022, 152, 1737-1746.	1.3	4
6	Calprotectin levels in amniotic fluid in relation to intra-amniotic inflammation and infection in women with preterm labor with intact membranes: A retrospective cohort study. European Journal of Obstetrics, Gynecology and Reproductive Biology, 2022, 272, 24-29.	0.5	1
7	Umbilical cord blood metabolome differs in relation to delivery mode, birth order and sex, maternal diet and possibly future allergy development in rural children. PLoS ONE, 2021, 16, e0242978.	1.1	10
8	High-Moisture Meat Analogues Produced from Yellow Pea and Faba Bean Protein Isolates/Concentrate: Effect of Raw Material Composition and Extrusion Parameters on Texture Properties. Foods, 2021, 10, 843.	1.9	78
9	Maternal Dietary Selenium Intake during Pregnancy and Neonatal Outcomes in the Norwegian Mother, Father, and Child Cohort Study. Nutrients, 2021, 13, 1239.	1.7	7
10	Associations of maternal and infant metabolomes with immune maturation and allergy development at 12Âmonths in the Swedish NICE-cohort. Scientific Reports, 2021, 11, 12706.	1.6	5
11	Maternal selenium intake and selenium status during pregnancy in relation to preeclampsia and pregnancy-induced hypertension in a large Norwegian Pregnancy Cohort Study. Science of the Total Environment, 2021, 798, 149271.	3.9	17
12	Thyroid hormones in relation to toxic metal exposure in pregnancy, and potential interactions with iodine and selenium. Environment International, 2021, 157, 106869.	4.8	15
13	Maternal Dietary Selenium Intake during Pregnancy Is Associated with Higher Birth Weight and Lower Risk of Small for Gestational Age Births in the Norwegian Mother, Father and Child Cohort Study. Nutrients, 2021, 13, 23.	1.7	12
14	Proportions of Polyunsaturated Fatty Acids in Umbilical Cord Blood at Birth Are Related to Atopic Eczema Development in the First Year of Life. Nutrients, 2021, 13, 3779.	1.7	9
15	Infant lodine and Selenium Status in Relation to Maternal Status and Diet During Pregnancy and Lactation. Frontiers in Nutrition, 2021, 8, 733602.	1.6	15
16	Maternal probiotic milk intake during pregnancy and breastfeeding complications in the Norwegian Mother and Child Cohort Study. European Journal of Nutrition, 2020, 59, 2219-2228.	1.8	4
17	Maternal dietary selenium intake is associated with increased gestational length and decreased risk of preterm delivery. British Journal of Nutrition, 2020, 123, 209-219.	1.2	19
18	Cord Blood Levels of EPA, a Marker of Fish Intake, Correlate with Infants' T- and B-Lymphocyte Phenotypes and Risk for Allergic Disease. Nutrients, 2020, 12, 3000.	1.7	9

MALIN BARMAN

0.7

12

#	Article	IF	CITATIONS
19	Maternal Intake of Cow's Milk during Lactation Is Associated with Lower Prevalence of Food Allergy in Offspring. Nutrients, 2020, 12, 3680.	1.7	22
20	Comprehensive proteomic investigation of infectious and inflammatory changes in late preterm prelabour rupture of membranes. Scientific Reports, 2020, 10, 17696.	1.6	6
21	Mid-trimester amniotic fluid proteome's association with spontaneous preterm delivery and gestational duration. PLoS ONE, 2020, 15, e0232553.	1.1	2
22	Low-level maternal exposure to cadmium, lead, and mercury and birth outcomes in a Swedish prospective birth-cohort. Environmental Pollution, 2020, 265, 114986.	3.7	34
23	Comparison of Bacterial DNA Profiles in Mid-Trimester Amniotic Fluid Samples From Preterm and Term Deliveries. Frontiers in Microbiology, 2020, 11, 415.	1.5	31
24	Fecal short chain fatty acids in children living on farms and a link between valeric acid and protection from eczema. Scientific Reports, 2020, 10, 22449.	1.6	39
25	Food and Nutrient Intake during Pregnancy in Relation to Maternal Characteristics: Results from the NICE Birth Cohort in Northern Sweden. Nutrients, 2019, 11, 1680.	1.7	40
26	Exposure to a Farm Environment During Pregnancy Increases the Proportion of Arachidonic Acid in the Cord Sera of Offspring. Nutrients, 2019, 11, 238.	1.7	3
27	Fish Oil Supplementation in Pregnancy Increases Gestational Age, Size for Gestational Age, and Birth Weight in Infants: A Randomized Controlled Trial. Journal of Nutrition, 2019, 149, 628-634.	1.3	26
28	Nutritional impact on Immunological maturation during Childhood in relation to the Environment (NICE): a prospective birth cohort in northern Sweden. BMJ Open, 2018, 8, e022013.	0.8	24
29	Late introduction of fish and eggs is associated with increased risk of allergy development – results from the FARMFLORA birth cohort. Food and Nutrition Research, 2017, 61, 1393306.	1.2	9
30	Serum fatty acids in infants, reflecting family fish consumption, wereÂinversely associated with allergy development but not related to farmÂresidence. Acta Paediatrica, International Journal of Paediatrics, 2016, 105, 1462-1471.	0.7	14
31	Effect of maternal supplementation with fish oil during pregnancy and lactation on allergy development in childhood. Acta Paediatrica, International Journal of Paediatrics, 2016, 105, 1348-1348.	0.7	2
32	Diet in 1-year-old farm and control children and allergy development: results from the FARMFLORA birth cohort. Food and Nutrition Research, 2016, 60, 32721.	1.2	11
33	Fat intake and breast milk fatty acid composition in farming and nonfarming women and allergy development in the offspring. Pediatric Research, 2016, 79, 114-123.	1.1	27
34	Single Nucleotide Polymorphisms in the FADS Gene Cluster but not the ELOVL2 Gene are Associated with Serum Polyunsaturated Fatty Acid Composition and Development of Allergy (in a Swedish Birth) Tj ETQqO	0 0 <b>1</b> gBT /C	)veøløck 10 Tf
35	No association between allergy and current 25â€hydroxy vitamin D in serum or vitamin D intake. Acta Paediatrica, International Journal of Paediatrics, 2015, 104, 405-413.	0.7	12

Serum fatty acid profile does not reflect seafood intake in adolescents with atopic eczema. Acta Paediatrica, International Journal of Paediatrics, 2014, 103, 968-976.

#	Article	IF	CITATIONS
37	High Levels of Both n-3 and n-6 Long-Chain Polyunsaturated Fatty Acids in Cord Serum Phospholipids Predict Allergy Development. PLoS ONE, 2013, 8, e67920.	1.1	27