Suzanne N Avery

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4942901/publications.pdf

Version: 2024-02-01

36 1,147 18 33 papers citations h-index g-index

37 37 37 37 1706

times ranked

citing authors

docs citations

all docs

#	Article	IF	CITATIONS
1	Anterior hippocampal dysfunction in early psychosis: a 2-year follow-up study. Psychological Medicine, 2023, 53, 160-169.	2.7	3
2	<scp>ENIGMAâ€anxiety</scp> working group: Rationale for and organization of <scp>largeâ€scale</scp> neuroimaging studies of anxiety disorders. Human Brain Mapping, 2022, 43, 83-112.	1.9	31
3	Development of Thalamocortical Structural Connectivity in Typically Developing and Psychosis Spectrum Youths. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging, 2022, 7, 782-792.	1.1	8
4	Increased amplitude of hippocampal low frequency fluctuations in early psychosis: A two-year follow-up study. Schizophrenia Research, 2022, 241, 260-266.	1.1	3
5	Structural Brain Correlates of Childhood Inhibited Temperament: An ENIGMA-Anxiety Mega-analysis. Journal of the American Academy of Child and Adolescent Psychiatry, 2022, 61, 1182-1188.	0.3	2
6	Relational Memory in the Early Stage of Psychosis: A 2-Year Follow-up Study. Schizophrenia Bulletin, 2021, 47, 75-86.	2.3	12
7	Stable habituation deficits in the early stage of psychosis: a 2-year follow-up study. Translational Psychiatry, 2021, 11, 20.	2.4	6
8	Habituation during encoding: A new approach to the evaluation of memory deficits in schizophrenia. Schizophrenia Research, 2020, 223, 179-185.	1.1	6
9	Relational memory in the early stage of psychotic bipolar disorder. Psychiatry Research, 2020, 294, 113508.	1.7	1
10	Visual exploration differences during relational memory encoding in early psychosis. Psychiatry Research, 2020, 287, 112910.	1.7	5
11	BNST-insula structural connectivity in humans. Neurolmage, 2020, 210, 116555.	2.1	26
12	Disrupted Habituation in the Early Stage of Psychosis. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging, 2019, 4, 1004-1012.	1.1	21
13	Impaired relational memory in the early stage of psychosis. Schizophrenia Research, 2019, 212, 113-120.	1.1	21
14	F77. HABITUATION DEFICITS ARE ASSOCIATED WITH RELATIONAL MEMORY IMPAIRMENT IN THE EARLY STAGE OF PSYCHOSIS. Schizophrenia Bulletin, 2019, 45, S283-S284.	2.3	0
15	19.4 RELATIONAL MEMORY AND HIPPOCAMPAL FUNCTION IN EARLY AND CHRONIC SCHIZOPHRENIA. Schizophrenia Bulletin, 2019, 45, S120-S121.	2.3	0
16	Social anxiety is associated with BNST response to unpredictability. Depression and Anxiety, 2019, 36, 666-675.	2.0	68
17	Hippocampal Network Modularity Is Associated With Relational Memory Dysfunction in Schizophrenia. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging, 2018, 3, 423-432.	1.1	23
18	Impaired associative inference in the early stage of psychosis. Schizophrenia Research, 2018, 202, 86-90.	1.1	17

#	Article	IF	CITATIONS
19	Slow to warm up: the role of habituation in social fear. Social Cognitive and Affective Neuroscience, 2016, 11, 1832-1840.	1.5	35
20	Impaired face recognition is associated with social inhibition. Psychiatry Research, 2016, 236, 53-57.	1.7	32
21	NEUROCIRCUITRY UNDERLYING RISK AND RESILIENCE TO SOCIAL ANXIETY DISORDER. Depression and Anxiety, 2014, 31, 822-833.	2.0	36
22	Structural and functional bases of inhibited temperament. Social Cognitive and Affective Neuroscience, 2014, 9, 2049-2058.	1.5	49
23	BNST neurocircuitry in humans. Neurolmage, 2014, 91, 311-323.	2.1	145
24	Relational memory and hippocampal function in psychotic bipolar disorder. European Archives of Psychiatry and Clinical Neuroscience, 2014, 264, 199-211.	1.8	11
25	Differences in age-related effects on brain volume in Down syndrome as compared to Williams syndrome and typical development. Journal of Neurodevelopmental Disorders, 2014, 6, 8.	1.5	29
26	Amygdala–cingulate intrinsic connectivity is associated with degree of social inhibition. Biological Psychology, 2014, 99, 15-25.	1.1	63
27	Amygdala and hippocampus fail to habituate to faces in individuals with an inhibited temperament. Social Cognitive and Affective Neuroscience, 2013, 8, 143-150.	1.5	91
28	The effect of intellectual ability on functional activation in a neurodevelopmental disorder: preliminary evidence from multiple fMRI studies in Williams syndrome. Journal of Neurodevelopmental Disorders, 2012, 4, 24.	1.5	4
29	Intact Relational Memory and Normal Hippocampal Structure in the Early Stage of Psychosis. Biological Psychiatry, 2012, 71, 105-113.	0.7	19
30	White matter integrity deficits in prefrontal–amygdala pathways in Williams syndrome. NeuroImage, 2012, 59, 887-894.	2.1	23
31	Using novel control groups to dissect the amygdala's role in Williams syndrome. Developmental Cognitive Neuroscience, 2011, 1, 295-304.	1.9	21
32	Sustained amygdala response to both novel and newly familiar faces characterizes inhibited temperament. Social Cognitive and Affective Neuroscience, 2011, 6, 621-629.	1.5	62
33	Internal representation of hierarchical sequences involves the default network. BMC Neuroscience, 2010, 11, 54.	0.8	8
34	Eye-Movement Behavior Reveals Relational Memory Impairment in Schizophrenia. Biological Psychiatry, 2010, 68, 617-624.	0.7	46
35	A unique role for the human amygdala in novelty detection. NeuroImage, 2010, 50, 1188-1193.	2.1	158
36	Amygdala temporal dynamics: temperamental differences in the timing of amygdala response to familiar and novel faces. BMC Neuroscience, 2009, 10, 145.	0.8	61

3