

Dimitris P Makris

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/494358/publications.pdf>

Version: 2024-02-01

149
papers

6,475
citations

61984

43
h-index

74163

75
g-index

149
all docs

149
docs citations

149
times ranked

6040
citing authors

#	ARTICLE	IF	CITATIONS
1	Citric acid-based deep eutectic solvent for the anthocyanin recovery from <i>Hibiscus sabdariffa</i> through microwave-assisted extraction. <i>Biomass Conversion and Biorefinery</i> , 2022, 12, 351-360.	4.6	37
2	Extraction of Polyphenols from Olive Leaves Employing Deep Eutectic Solvents: The Application of Chemometrics to a Quantitative Study on Antioxidant Compounds. <i>Applied Sciences (Switzerland)</i> , 2022, 12, 831.	2.5	14
3	Organosolv Treatment/Polyphenol Extraction from Olive Leaves (<i>Olea europaea</i> L.) Using Glycerol and Glycerol-Based Deep Eutectic Solvents: Effect on Metabolite Stability. <i>Biomass</i> , 2022, 2, 46-61.	2.8	8
4	Mulching Effect on Quantitative and Qualitative Characteristics of Yield in Sweet Potatoes. <i>Horticulturae</i> , 2022, 8, 271.	2.8	3
5	Pulsed Electric Field-Based Extraction of Total Polyphenols from <i>Sideritis raiseri</i> Using Hydroethanolic Mixtures. <i>Oxygen</i> , 2022, 2, 91-98.	5.0	16
6	Process Optimization and Stability of Waste Orange Peel Polyphenols in Extracts Obtained with Organosolv Thermal Treatment Using Glycerol-Based Solvents. <i>ChemEngineering</i> , 2022, 6, 35.	2.4	11
7	Extraction of Polyphenolic Antioxidants from Red Grape Pomace and Olive Leaves: Process Optimization Using a Tailor-Made Tertiary Deep Eutectic Solvent. <i>Sustainability</i> , 2022, 14, 6864.	3.2	11
8	Enhanced-performance extraction of olive (<i>Olea europaea</i>) leaf polyphenols using L-lactic acid/ammonium acetate deep eutectic solvent combined with β -cyclodextrin: screening, optimisation, temperature effects and stability. <i>Biomass Conversion and Biorefinery</i> , 2021, 11, 1125-1136.	4.6	25
9	Extractor dimensions affect optimization of laboratory-scale batch solid-liquid extraction of polyphenols from plant material: potato peels as a case study. <i>Chemical Engineering Communications</i> , 2021, 208, 1618-1629.	2.6	2
10	Recovery and applications of enzymes from food wastes. , 2021, , 313-325.		1
11	Fortification of chocolate using <i>Moringa oleifera</i> extract encapsulated in microemulsions. <i>OCL - Oilseeds and Fats, Crops and Lipids</i> , 2021, 28, 38.	1.4	6
12	Empirical Kinetic Modelling and Mechanisms of Quercetin Thermal Degradation in Aqueous Model Systems: Effect of pH and Addition of Antioxidants. <i>Applied Sciences (Switzerland)</i> , 2021, 11, 2579.	2.5	5
13	Evaluation of Pulsed Electric Field Polyphenol Extraction from <i>Vitis vinifera</i> , <i>Sideritis scardica</i> and <i>Crocus sativus</i> . <i>ChemEngineering</i> , 2021, 5, 25.	2.4	24
14	Encapsulation of <i>Moringa oleifera</i> Extract in Ca-Alginate Chocolate Beads: Physical and Antioxidant Properties. <i>Journal of Food Quality</i> , 2021, 2021, 1-9.	2.6	9
15	Physical Properties of Chitosan Films Containing Pomegranate Peel Extracts Obtained by Deep Eutectic Solvents. <i>Foods</i> , 2021, 10, 1262.	4.3	19
16	Use of Pulsed Electric Field as a Low-Temperature and High-Performance "Green" Extraction Technique for the Recovery of High Added Value Compounds from Olive Leaves. <i>Beverages</i> , 2021, 7, 45.	2.8	22
17	Pulsed Electric Field and <i>Salvia officinalis</i> L. Leaves: A Successful Combination for the Extraction of High Value Added Compounds. <i>Foods</i> , 2021, 10, 2014.	4.3	10
18	Cyclodextrins as high-performance green co-solvents in the aqueous extraction of polyphenols and anthocyanin pigments from solid onion waste. <i>European Food Research and Technology</i> , 2021, 247, 2831-2845.	3.3	10

#	ARTICLE	IF	CITATIONS
19	Integrated Green Process for the Extraction of Red Grape Pomace Antioxidant Polyphenols Using Ultrasound-Assisted Pretreatment and β -Cyclodextrin. <i>Beverages</i> , 2021, 7, 59.	2.8	13
20	Enhancement of Polyphenols Recovery from <i>Rosa canina</i> , <i>Calendula officinalis</i> and <i>Castanea sativa</i> Using Pulsed Electric Field. <i>Beverages</i> , 2021, 7, 63.	2.8	10
21	Optimization of Pulsed Electric Field as Standalone "Green" Extraction Procedure for the Recovery of High Value-Added Compounds from Fresh Olive Leaves. <i>Antioxidants</i> , 2021, 10, 1554.	5.1	27
22	Development of a Green Methodology for Simultaneous Extraction of Polyphenols and Pigments from Red Winemaking Solid Wastes (Pomace) Using a Novel Glycerol-Sodium Benzoate Deep Eutectic Solvent and Ultrasonication Pretreatment. <i>Environments - MDPI</i> , 2021, 8, 90.	3.3	12
23	Appraisal of the combined effect of time and temperature on the total polyphenol yield in batch stirred-tank extraction of medicinal and aromatic plants: The extraction efficiency factor. <i>Journal of Applied Research on Medicinal and Aromatic Plants</i> , 2021, 25, 100340.	1.5	6
24	Pressurized Liquid Extraction of Polyphenols and Anthocyanins from Saffron Processing Waste with Aqueous Organic Acid Solutions: Comparison with Stirred-Tank and Ultrasound-Assisted Techniques. <i>Sustainability</i> , 2021, 13, 12578.	3.2	10
25	Kinetics and modeling of L-cysteine effect on the Cu(II)-induced oxidation of quercetin. <i>Chemical Engineering Communications</i> , 2020, 207, 139-152.	2.6	5
26	High-efficiency Extraction of Phenolics from Wheat Waste Biomass (Bran) by Combining Deep Eutectic Solvent, Ultrasound-assisted Pretreatment and Thermal Treatment. <i>Environmental Processes</i> , 2020, 7, 845-859.	3.5	20
27	Batch Stirred-Tank Green Extraction of <i>Salvia fruticosa</i> Mill. Polyphenols Using Newly Designed Citrate-Based Deep Eutectic Solvents and Ultrasonication Pretreatment. <i>Applied Sciences (Switzerland)</i> , 2020, 10, 4774.	2.5	8
28	Editorial "High-Performance Green Extraction of Natural Products". <i>Applied Sciences (Switzerland)</i> , 2020, 10, 7672.	2.5	0
29	Hydroxypropyl- β -Cyclodextrin as a Green Co-Solvent in the Aqueous Extraction of Polyphenols from Waste Orange Peels. <i>Beverages</i> , 2020, 6, 50.	2.8	19
30	Glycerol and Glycerol-Based Deep Eutectic Mixtures as Emerging Green Solvents for Polyphenol Extraction: The Evidence So Far. <i>Molecules</i> , 2020, 25, 5842.	3.8	38
31	High-Performance Green Extraction of Polyphenolic Antioxidants from <i>Salvia fruticosa</i> Using Cyclodextrins: Optimization, Kinetics, and Composition. <i>Applied Sciences (Switzerland)</i> , 2020, 10, 3447.	2.5	13
32	Hydroglycerolic Solvent and Ultrasonication Pretreatment: A Green Blend for High-Efficiency Extraction of <i>Salvia fruticosa</i> Polyphenols. <i>Sustainability</i> , 2020, 12, 4840.	3.2	7
33	Development of a Low-Temperature and High-Performance Green Extraction Process for the Recovery of Polyphenolic Phytochemicals from Waste Potato Peels Using Hydroxypropyl β -Cyclodextrin. <i>Applied Sciences (Switzerland)</i> , 2020, 10, 3611.	2.5	12
34	The Effect of Ultrasonication Pretreatment on the Production of Polyphenol-Enriched Extracts from <i>Moringa oleifera</i> L. (Drumstick Tree) Using a Novel Bio-Based Deep Eutectic Solvent. <i>Applied Sciences (Switzerland)</i> , 2020, 10, 220.	2.5	29
35	A Green Extraction Process for Polyphenols from Elderberry (<i>Sambucus nigra</i>) Flowers Using Deep Eutectic Solvent and Ultrasound-Assisted Pretreatment. <i>Molecules</i> , 2020, 25, 921.	3.8	46
36	Green Valorization of Olive Leaves to Produce Polyphenol-Enriched Extracts Using an Environmentally Benign Deep Eutectic Solvent. <i>AgriEngineering</i> , 2020, 2, 226-239.	3.2	19

#	ARTICLE	IF	CITATIONS
37	Polyphenol Extraction from <i>Humulus lupulus</i> (Hop) Using a Neoteric Glycerol/L-Alanine Deep Eutectic Solvent: Optimisation, Kinetics and the Effect of Ultrasound-Assisted Pretreatment. <i>AgriEngineering</i> , 2019, 1, 403-417.	3.2	27
38	Optimization of Bioactive Substances in the Wastes of Some Selective Mediterranean Crops. <i>Beverages</i> , 2019, 5, 42.	2.8	9
39	Development of sodium propionate-based deep eutectic solvents for polyphenol extraction from onion solid wastes. <i>Clean Technologies and Environmental Policy</i> , 2019, 21, 1563-1574.	4.1	29
40	Saffron Processing Wastes as a Bioresource of High-Value Added Compounds: Development of a Green Extraction Process for Polyphenol Recovery Using a Natural Deep Eutectic Solvent. <i>Antioxidants</i> , 2019, 8, 586.	5.1	70
41	Polyphenolic Antioxidants from Agri-Food Waste Biomass. <i>Antioxidants</i> , 2019, 8, 624.	5.1	20
42	Green extraction processes for the efficient recovery of bioactive polyphenols from wine industry solid wastes – Recent progress. <i>Current Opinion in Green and Sustainable Chemistry</i> , 2018, 13, 50-55.	5.9	27
43	Incorporation of 2-hydroxypropyl β -cyclodextrin in a biomolecule-based low-transition temperature mixture (LTTM) boosts efficiency of polyphenol extraction from <i>Moringa oleifera</i> Lam leaves. <i>Journal of Applied Research on Medicinal and Aromatic Plants</i> , 2018, 9, 62-69.	1.5	10
44	Highly Efficient Extraction of Antioxidant Polyphenols from <i>Olea europaea</i> Leaves Using an Eco-friendly Glycerol/Glycine Deep Eutectic Solvent. <i>Waste and Biomass Valorization</i> , 2018, 9, 1985-1992.	3.4	57
45	Methyl β -cyclodextrin as a booster for the extraction for <i>Olea europaea</i> leaf polyphenols with a bio-based deep eutectic solvent. <i>Biomass Conversion and Biorefinery</i> , 2018, 8, 345-355.	4.6	39
46	Empirical Kinetic Modelling of the Effect of L-Ascorbic Acid on the Cu(II)-Induced Oxidation of Quercetin. <i>ChemEngineering</i> , 2018, 2, 46.	2.4	2
47	Characterisation of Polyphenol-Containing Extracts from <i>Stachys mucronata</i> and Evaluation of Their Antiradical Activity. <i>Medicines (Basel, Switzerland)</i> , 2018, 5, 14.	1.4	7
48	Stability effects of methyl β -cyclodextrin on <i>Olea europaea</i> leaf extracts in a natural deep eutectic solvent. <i>European Food Research and Technology</i> , 2018, 244, 1783-1792.	3.3	14
49	A Green Extraction Process to Recover Polyphenols from Byproducts of Hemp Oil Processing. <i>Recycling</i> , 2018, 3, 15.	5.0	16
50	Polyphenol extraction from <i>Origanum dictamnus</i> using low-transition temperature mixtures composed of glycerol and organic salts: Effect of organic anion carbon chain length. <i>Chemical Engineering Communications</i> , 2018, 205, 1494-1506.	2.6	14
51	Natural food colorants derived from onion wastes: Application in a yoghurt product. <i>Electrophoresis</i> , 2018, 39, 1975-1983.	2.4	45
52	The effect of 2-hydroxypropyl β -cyclodextrin on the stability of polyphenolic compounds from <i>Moringa oleifera</i> Lam leaf extracts in a natural low-transition temperature mixture. <i>Nova Biotechnologica Et Chimica</i> , 2018, 17, 29-37.	0.1	3
53	Optimised extraction of antioxidant polyphenols from <i>Satureja thymbra</i> using newly designed glycerol-based natural low-transition temperature mixtures (LTTMs). <i>Journal of Applied Research on Medicinal and Aromatic Plants</i> , 2017, 6, 31-40.	1.5	34
54	Screening of Natural Sodium Acetate-Based Low-Transition Temperature Mixtures (LTTMs) for Enhanced Extraction of Antioxidants and Pigments from Red Vinification Solid Wastes. <i>Environmental Processes</i> , 2017, 4, 123-135.	3.5	30

#	ARTICLE	IF	CITATIONS
55	Enhanced extraction of antioxidant polyphenols from <i>Moringa oleifera</i> Lam. leaves using a biomolecule-based low-transition temperature mixture. <i>European Food Research and Technology</i> , 2017, 243, 1839-1848.	3.3	32
56	Nutritional Characterization of Leaves and Herbal Tea of <i>Moringa oleifera</i> Cultivated in Greece. <i>Journal of Herbs, Spices and Medicinal Plants</i> , 2017, 23, 320-333.	1.1	20
57	Implementation of kinetics and response surface methodology reveals contrasting effects of catechin and chlorogenic acid on the development of browning in wine model systems containing either ascorbic acid or sulphite. <i>European Food Research and Technology</i> , 2017, 243, 565-574.	3.3	5
58	Ultrasound-Assisted Extraction of Polyphenolic Antioxidants from Olive (<i>Olea europaea</i>) Leaves Using a Novel Glycerol/Sodium-Potassium Tartrate Low-Transition Temperature Mixture (LTTM). <i>Environments - MDPI</i> , 2017, 4, 31.	3.3	30
59	Combination of Lactic Acid-Based Deep Eutectic Solvents (DES) with β -Cyclodextrin: Performance Screening Using Ultrasound-Assisted Extraction of Polyphenols from Selected Native Greek Medicinal Plants. <i>Agronomy</i> , 2017, 7, 54.	3.0	56
60	Accelerated Aging of the Traditional Greek Distillate Tsipouro Using Wooden Chips. Part I: Effect of Static Maceration vs. Ultrasonication on the Polyphenol Extraction and Antioxidant Activity. <i>Beverages</i> , 2017, 3, 5.	2.8	11
61	Low-Transition Temperature Mixtures (LTTMs) Made of Bioorganic Molecules: Enhanced Extraction of Antioxidant Phenolics from Industrial Cereal Solid Wastes. <i>Recycling</i> , 2017, 2, 3.	5.0	20
62	Effect of Methyl β -cyclodextrin on Radical Scavenging Kinetics of Olive Leaf Extracts and Interactions with Ascorbic Acid. <i>ChemEngineering</i> , 2017, 1, 6.	2.4	2
63	Optimization of a Green Extraction/Inclusion Complex Formation Process to Recover Antioxidant Polyphenols from Oak Acorn Husks (<i>Quercus Robur</i>) Using Aqueous 2-Hydroxypropyl- β -Cyclodextrin/Glycerol Mixtures. <i>Environments - MDPI</i> , 2016, 3, 3.	3.3	17
64	Extraction of Antioxidant Phenolics from Agri-Food Waste Biomass Using a Newly Designed Glycerol-Based Natural Low-Transition Temperature Mixture: A Comparison with Conventional Eco-Friendly Solvents. <i>Recycling</i> , 2016, 1, 194-204.	5.0	57
65	Kinetics of Ultrasound-Assisted Flavonoid Extraction from Agri-Food Solid Wastes Using Water/Glycerol Mixtures. <i>Resources</i> , 2016, 5, 7.	3.5	17
66	Development of a Green Process for the Preparation of Antioxidant and Pigment-Enriched Extracts from Winery Solid Wastes Using Response Surface Methodology and Kinetics. <i>Chemical Engineering Communications</i> , 2016, 203, 1317-1325.	2.6	27
67	Novel Glycerol-Based Natural Eutectic Mixtures and Their Efficiency in the Ultrasound-Assisted Extraction of Antioxidant Polyphenols from Agri-Food Waste Biomass. <i>Waste and Biomass Valorization</i> , 2016, 7, 1377-1387.	3.4	120
68	Novel lactic acid-based natural deep eutectic solvents: Efficiency in the ultrasound-assisted extraction of antioxidant polyphenols from common native Greek medicinal plants. <i>Journal of Applied Research on Medicinal and Aromatic Plants</i> , 2016, 3, 120-127.	1.5	136
69	Optimization of a green extraction method for the recovery of polyphenols from olive leaf using cyclodextrins and glycerin as co-solvents. <i>Journal of Food Science and Technology</i> , 2016, 53, 3939-3947.	2.8	47
70	Kinetics of Ultrasound-Assisted Polyphenol Extraction from Spent Filter Coffee Using Aqueous Glycerol. <i>Chemical Engineering Communications</i> , 2016, 203, 407-413.	2.6	36
71	Optimization of polyphenol extraction from red grape pomace using aqueous glycerol/tartaric acid mixtures and response surface methodology. <i>Preparative Biochemistry and Biotechnology</i> , 2016, 46, 176-182.	1.9	27
72	Ultrasound-Assisted Green Extraction of Eggplant Peel (<i>Solanum melongena</i>) Polyphenols Using Aqueous Mixtures of Glycerol and Ethanol: Optimisation and Kinetics. <i>Environmental Processes</i> , 2016, 3, 369-386.	3.5	57

#	ARTICLE	IF	CITATIONS
91	Factorial design optimisation of hydrocaffeic acid removal from an aqueous matrix by the use of a crude potato polyphenol oxidase. <i>Biocatalysis and Agricultural Biotechnology</i> , 2013, 2, 305-310.	3.1	2
92	The effect of chlorogenic acid, catechin and SO ₂ on browning development in white wine model solutions. <i>Journal of the Institute of Brewing</i> , 2013, 119, 309-313.	2.3	4
93	Characterization of Polyphenolic Phytochemicals in Red Grape Pomace. <i>International Journal of Waste Resources</i> , 2013, 03, .	0.2	23
94	Removal of Olive Mill Wastewater Phenolics with the Use of a Polyphenol Oxidase Homogenate from Potato Peel Waste. <i>Journal of Waste Management</i> , 2013, 2013, 1-7.	0.5	3
95	Use of response surface methodology to evaluate the reducing power in binary solutions of ascorbic acid with natural polyphenolic antioxidants. <i>International Journal of Food Studies</i> , 2013, 2, .	0.8	8
96	Stability and transformation of major flavonols in onion (<i>Allium cepa</i>) solid wastes. <i>Journal of Food Science and Technology</i> , 2012, 49, 489-494.	2.8	28
97	Implementation of response surface methodology to assess the antiradical behaviour in mixtures of ascorbic acid and Î±-tocopherol with grape (<i>Vitis vinifera</i>) stem extracts. <i>Food Chemistry</i> , 2012, 132, 351-359.	8.2	21
98	Chlorogenic Acid Oxidation by a Crude Peroxidase Preparation: Biocatalytic Characteristics and Oxidation Products. <i>Food and Bioprocess Technology</i> , 2012, 5, 243-251.	4.7	12
99	Deployment of response surface methodology to optimize recovery of grape (<i>Vitis vinifera</i>) stem and seed polyphenols. <i>Procedia Food Science</i> , 2011, 1, 1686-1693.	0.6	9
100	Interactions of natural antioxidants with red grape pomace anthocyanins in a liquid model matrix: Stability and copigmentation effects. <i>Chemical Industry and Chemical Engineering Quarterly</i> , 2011, 17, 59-66.	0.7	10
101	BROWNING DEVELOPMENT IN WINE-LIKE LIQUID MODEL MATRICES: DEPENDENCE ON PHENOLIC, FE(III) AND SO ₂ CONCENTRATIONS. <i>Journal of Food Process Engineering</i> , 2010, 33, 934-945.	2.9	2
102	Removal of olive mill waste water phenolics using a crude peroxidase extract from onion by-products. <i>Environmental Chemistry Letters</i> , 2010, 8, 271-275.	16.2	15
103	Comparison of fisetin and quercetin oxidation with a cell-free extract of onion trimmings and peel, plant waste, containing peroxidase enzyme: a further insight into flavonol degradation mechanism. <i>International Journal of Food Science and Technology</i> , 2010, 45, 2265-2271.	2.7	12
104	Polyphenol characterization and encapsulation in Î²-cyclodextrin of a flavonoid-rich <i>Hypericum perforatum</i> (St John's wort) extract. <i>LWT - Food Science and Technology</i> , 2010, 43, 882-889.	5.2	103
105	Optimisation of Anthocyanin Recovery from Onion (<i>Allium cepa</i>) Solid Wastes Using Response Surface Methodology. <i>Journal of Food Technology</i> , 2010, 8, 183-186.	0.5	9
106	Biocatalytic properties of a peroxidase-active cell-free extract from onion solid wastes: caffeic acid oxidation. <i>Biodegradation</i> , 2009, 20, 143-153.	3.0	14
107	Polyphenolic Composition and Antioxidant Characteristics of Kumquat (<i>Fortunella margarita</i>) Peel Fractions. <i>Plant Foods for Human Nutrition</i> , 2009, 64, 297-302.	3.2	62
108	An Investigation on the Recovery of Antioxidant Phenolics from Onion Solid Wastes Employing Water/Ethanol-Based Solvent Systems. <i>Food and Bioprocess Technology</i> , 2009, 2, 337-343.	4.7	74

#	ARTICLE	IF	CITATIONS
109	Factorial design optimisation of grape (<i>Vitis vinifera</i>) seed polyphenol extraction. <i>European Food Research and Technology</i> , 2009, 229, 731-742.	3.3	35
110	Anthocyanin profiles of major red grape (<i>Vitis vinifera</i> L.) varieties cultivated in Greece and their relationship with <i>in vitro</i> antioxidant characteristics. <i>International Journal of Food Science and Technology</i> , 2009, 44, 2385-2393.	2.7	37
111	Deployment of response surface methodology to optimise recovery of grape (<i>Vitis vinifera</i>) stem polyphenols. <i>Talanta</i> , 2009, 79, 1311-1321.	5.5	65
112	Implementation of response surface methodology to optimise extraction of onion (<i>Allium cepa</i>) solid waste phenolics. <i>Innovative Food Science and Emerging Technologies</i> , 2009, 10, 246-252.	5.6	85
113	Investigation on biocatalytic properties of a peroxidase-active homogenate from onion solid wastes: An insight into quercetin oxidation mechanism. <i>Process Biochemistry</i> , 2008, 43, 861-867.	3.7	45
114	Optimisation of the extraction of olive (<i>Olea europaea</i>) leaf phenolics using water/ethanol-based solvent systems and response surface methodology. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 392, 977-985.	3.7	147
115	Characterisation of certain major polyphenolic antioxidants in grape (<i>Vitis vinifera</i> cv. Roditis) stems by liquid chromatography-mass spectrometry. <i>European Food Research and Technology</i> , 2008, 226, 1075-1079.	3.3	47
116	Hydrocaffeic acid oxidation by a peroxidase homogenate from onion solid wastes. <i>European Food Research and Technology</i> , 2008, 227, 1379-1386.	3.3	12
117	Evolution of benzoate derivatives and their hydroxycinnamate analogues during ageing of white wines in oak barrels. <i>Journal of Food Composition and Analysis</i> , 2008, 21, 667-671.	3.9	11
118	Extraction of phenolics in liquid model matrices containing oak chips: Kinetics, liquid chromatography-mass spectroscopy characterisation and association with <i>in vitro</i> antiradical activity. <i>Food Chemistry</i> , 2008, 110, 263-272.	8.2	28
119	Thermal Stability of Anthocyanin Extract of <i>Hibiscus sabdariffa</i> L. in the Presence of β -Cyclodextrin. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 10303-10310.	5.2	88
120	Peroxidase-active cell free extract from onion solid wastes: biocatalytic properties and putative pathway of ferulic acid oxidation. <i>Journal of Bioscience and Bioengineering</i> , 2008, 106, 279-285.	2.2	17
121	An Investigation on Factors Affecting Recovery of Antioxidant Phenolics and Anthocyanins from Red Grape (<i>Vitis vinifera</i> L.) Pomace Employing Water/Ethanol-Based Solutions. <i>American Journal of Food Technology</i> , 2008, 3, 164-173.	0.2	40
122	Differentiation of Young Red Wines Based on Chemometrics of Minor Polyphenolic Constituents. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 3233-3239.	5.2	32
123	Biomimetic oxidation of quercetin: Isolation of a naturally occurring quercetin heterodimer and evaluation of its <i>in vitro</i> antioxidant properties. <i>Food Research International</i> , 2007, 40, 7-14.	6.2	74
124	Interactions between quercetin and catechin in a model matrix: Effects on the <i>in vitro</i> antioxidant behaviour. <i>Food Research International</i> , 2007, 40, 819-826.	6.2	13
125	Recovery of antioxidant phenolics from white vinification solid by-products employing water/ethanol mixtures. <i>Bioresource Technology</i> , 2007, 98, 2963-2967.	9.6	134
126	Polyphenolic content and <i>in vitro</i> antioxidant characteristics of wine industry and other agri-food solid waste extracts. <i>Journal of Food Composition and Analysis</i> , 2007, 20, 125-132.	3.9	425

#	ARTICLE	IF	CITATIONS
127	Copper(II)-mediated biomimetic oxidation of quercetin: generation of a naturally occurring oxidation product and evaluation of its in vitro antioxidant properties. <i>European Food Research and Technology</i> , 2007, 225, 435-441.	3.3	22
128	Differentiation of young red wines based on cultivar and geographical origin with application of chemometrics of principal polyphenolic constituents. <i>Talanta</i> , 2006, 70, 1143-1152.	5.5	101
129	Kinetics of browning onset in white wines: influence of principal redox-active polyphenols and impact on the reducing capacity. <i>Food Chemistry</i> , 2006, 94, 98-104.	8.2	46
130	Storage of olives (<i>Olea europaea</i>) under CO ₂ atmosphere: Effect on anthocyanins, phenolics, sensory attributes and in vitro antioxidant properties. <i>Food Chemistry</i> , 2006, 99, 342-349.	8.2	31
131	Flavonols in grapes, grape products and wines: Burden, profile and influential parameters. <i>Journal of Food Composition and Analysis</i> , 2006, 19, 396-404.	3.9	146
132	Determination of major anthocyanin pigments in Hellenic native grape varieties (<i>Vitis vinifera</i> sp.): association with antiradical activity. <i>Journal of Food Composition and Analysis</i> , 2005, 18, 375-386.	3.9	146
133	Determination of low molecular weight polyphenolic constituents in grape (<i>Vitis vinifera</i> sp.) seed extracts: Correlation with antiradical activity. <i>Food Chemistry</i> , 2005, 89, 1-9.	8.2	168
134	Browning development in white wines: dependence on compositional parameters and impact on antioxidant characteristics. <i>European Food Research and Technology</i> , 2005, 220, 326-330.	3.3	29
135	An analytical survey of the polyphenols of seeds of varieties of grape (<i>Vitis vinifera</i>) cultivated in Greece: implications for exploitation as a source of value-added phytochemicals. <i>Phytochemical Analysis</i> , 2005, 16, 17-23.	2.4	42
136	Oxidation of caffeic acid in the presence of l-cysteine: isolation of 2-S-cysteinylcaffeic acid and evaluation of its antioxidant properties. <i>Food Research International</i> , 2005, 38, 395-402.	6.2	77
137	The effect of polyphenolic composition as related to antioxidant capacity in white wines. <i>Food Research International</i> , 2003, 36, 805-814.	6.2	83
138	Polyphenols in Hellenic wines: creating composition tables as a tool for epidemiological studies. <i>Journal of Wine Research</i> , 2003, 14, 103-114.	1.5	2
139	Anthocyanin Composition and Colour Characteristics of Selected Aged Wines Produced in Greece. <i>Journal of Wine Research</i> , 2002, 13, 23-34.	1.5	25
140	Evaluation of the antiradical and reducing properties of selected Greek white wines: correlation with polyphenolic composition. <i>Journal of the Science of Food and Agriculture</i> , 2002, 82, 1014-1020.	3.5	53
141	Effect of natural antioxidants on heat-induced, copper(II)-catalysed, oxidative degradation of quercetin and rutin (quercetin 3-O-rutinoside) in aqueous model systems. <i>Journal of the Science of Food and Agriculture</i> , 2002, 82, 1147-1153.	3.5	21
142	Hydroxyl Free Radical-Mediated Oxidative Degradation of Quercetin and Morin: A Preliminary Investigation. <i>Journal of Food Composition and Analysis</i> , 2002, 15, 103-113.	3.9	76
143	Correlation of Pigment and Flavanol Content with Antioxidant Properties in Selected Aged Regional Wines from Greece. <i>Journal of Food Composition and Analysis</i> , 2002, 15, 655-665.	3.9	316
144	An investigation on structural aspects influencing product formation in enzymic and chemical oxidation of quercetin and related flavonols. <i>Food Chemistry</i> , 2002, 77, 177-185.	8.2	63

#	ARTICLE	IF	CITATIONS
145	Domestic Processing of Onion Bulbs (<i>Allium cepa</i>) and Asparagus Spears (<i>Asparagus officinalis</i>): Effect on Flavonol Content and Antioxidant Status. <i>Journal of Agricultural and Food Chemistry</i> , 2001, 49, 3216-3222.	5.2	209
146	Comparison of Quercetin and a Non-Orthohydroxy Flavonol As Antioxidants by Competing In Vitro Oxidation Reactions. <i>Journal of Agricultural and Food Chemistry</i> , 2001, 49, 3370-3377.	5.2	63
147	Effect of Principal Polyphenolic Components in Relation to Antioxidant Characteristics of Aged Red Wines. <i>Journal of Agricultural and Food Chemistry</i> , 2001, 49, 5736-5742.	5.2	338
148	Heat-Induced, Metal-Catalyzed Oxidative Degradation of Quercetin and Rutin (Quercetin) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 627 Td</i> 2000, 48, 3830-3838.	5.2	179
149	High Performance Liquid Chromatography Studies on Free Radical Oxidation of Flavonols. , 2000, , 249-251.		4