

# Kim Ravnskjaer

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4943809/publications.pdf>

Version: 2024-02-01

17  
papers

1,803  
citations

687363

13  
h-index

888059

17  
g-index

17  
all docs

17  
docs citations

17  
times ranked

5295  
citing authors

#	ARTICLE	IF	CITATIONS
1	Circulating TREM2 as a noninvasive diagnostic biomarker for NASH in patients with elevated liver stiffness. <i>Hepatology</i> , 2023, 77, 558-572.	7.3	17
2	The Role of Diagnostic Biomarkers, Omics Strategies, and Single-Cell Sequencing for Nonalcoholic Fatty Liver Disease in Severely Obese Patients. <i>Journal of Clinical Medicine</i> , 2021, 10, 930.	2.4	6
3	Loss of CREB Coactivator CRTC1 in SF1 Cells Leads to Hyperphagia and Obesity by High-fat Diet But Not Normal Chow Diet. <i>Endocrinology</i> , 2021, 162, .	2.8	3
4	Liver-fibrosis-activated transcriptional networks govern hepatocyte reprogramming and intra-hepatic communication. <i>Cell Metabolism</i> , 2021, 33, 1685-1700.e9.	16.2	73
5	Osteoprotegerin Is more than a Possible Serum Marker in Liver Fibrosis: A Study into Its Function in Human and Murine Liver. <i>Pharmaceutics</i> , 2020, 12, 471.	4.5	15
6	Transcriptional Dynamics of Hepatic Sinusoid-Associated Cells After Liver Injury. <i>Hepatology</i> , 2020, 72, 2119-2133.	7.3	62
7	Transcriptional regulation of Hepatic Stellate Cell activation in NASH. <i>Scientific Reports</i> , 2019, 9, 2324.	3.3	65
8	Hepatic Insulin Resistance Following Chronic Activation of the CREB Coactivator CRTC2. <i>Journal of Biological Chemistry</i> , 2015, 290, 25997-26006.	3.4	26
9	Glucagon regulates gluconeogenesis through KAT2B- and WDR5-mediated epigenetic effects. <i>Journal of Clinical Investigation</i> , 2013, 123, 4318-4328.	8.2	73
10	Keystone Symposia on Epigenomics and Chromatin Dynamics: Keystone resort, CO, January 17-22, 2012. <i>Epigenetics</i> , 2012, 7, 522-523.	2.7	1
11	Class IIa Histone Deacetylases Are Hormone-Activated Regulators of FOXO and Mammalian Glucose Homeostasis. <i>Cell</i> , 2011, 145, 607-621.	28.9	486
12	PPAR $\gamma$ is a fatty acid sensor that enhances mitochondrial oxidation in insulin-secreting cells and protects against fatty acid-induced dysfunction. <i>Journal of Lipid Research</i> , 2010, 51, 1370-1379.	4.2	71
13	Targeted disruption of the CREB coactivator <i>Crtc2</i> increases insulin sensitivity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 3087-3092.	7.1	137
14	A fasting inducible switch modulates gluconeogenesis via activator/coactivator exchange. <i>Nature</i> , 2008, 456, 269-273.	27.8	481
15	Cooperative interactions between CBP and TORC2 confer selectivity to CREB target gene expression. <i>EMBO Journal</i> , 2007, 26, 2880-2889.	7.8	148
16	Glucose-induced repression of PPAR $\gamma$ gene expression in pancreatic $\beta$ -cells involves PP2A activation and AMPK inactivation. <i>Journal of Molecular Endocrinology</i> , 2006, 36, 289-299.	2.5	82
17	Pancreatic $\beta$ -Cell Lipotoxicity Induced by Overexpression of Hormone-Sensitive Lipase. <i>Diabetes</i> , 2003, 52, 2057-2065.	0.6	57