## **Cornelis P Tensen**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4945203/publications.pdf Version: 2024-02-01



| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Differential expression of CXCR3 targeting chemokines CXCL10, CXCL9, and CXCL11 in different types of skin inflammation. Journal of Pathology, 2001, 194, 398-405.  | 2.1 | 332       |
| 2  | The mutational landscape of cutaneous T cell lymphoma and Sézary syndrome. Nature Genetics, 2015, 47, 1465-1470.  | 9.4 | 322       |
| 3  | Distinct types of primary cutaneous large B-cell lymphoma identified by gene expression profiling.<br>Blood, 2005, 105, 3671-3678.  | 0.6 | 266       |
| 4  | Expression of MCP-1 by Reactive Astrocytes in Demyelinating Multiple Sclerosis Lesions. American<br>Journal of Pathology, 1999, 154, 45-51.   | 1.9 | 238       |
| 5  | Epigenetic Profiling of Cutaneous T-Cell Lymphoma: Promoter Hypermethylation of Multiple Tumor<br>Suppressor Genes Including BCL7a, PTPRG, and p73. Journal of Clinical Oncology, 2005, 23, 3886-3896.                        | 0.8 | 224       |
| 6  | Regulation of CD163 on human macrophages: cross-linking of CD163 induces signaling and activation.<br>Journal of Leukocyte Biology, 1999, 66, 858-866.  | 1.5 | 193       |
| 7  | Oncogenomic analysis of mycosis fungoides reveals major differences with Sézary syndrome. Blood, 2009, 113, 127-136.  | 0.6 | 188       |
| 8  | Novel and Highly Recurrent Chromosomal Alterations in Selzary Syndrome. Cancer Research, 2008, 68, 2689-2698.   | 0.4 | 176       |
| 9  | CXCR3-mediated chemotaxis of human T cells is regulated by a Gi- and phospholipase C–dependent<br>pathway and not via activation of MEK/p44/p42 MAPK nor Akt/PI-3 kinase. Blood, 2003, 102, 1959-1965.                        | 0.6 | 161       |
| 10 | Aberrant Expression of the Tyrosine Kinase Receptor EphA4 and the Transcription Factor Twist in<br>Sézary Syndrome Identified by Gene Expression Analysis. Cancer Research, 2004, 64, 5578-5586.                              | 0.4 | 155       |
| 11 | Discrepancy Between Molecular Structure and Ligand Selectivity of a Testicular Follicle-Stimulating<br>Hormone Receptor of the African Catfish (Clarias gariepinus)1. Biology of Reproduction, 2001, 64,<br>1633-1643.        | 1.2 | 153       |
| 12 | Distinct Efficacies for Two Endogenous Ligands on a Single Cognate Gonadoliberin Receptor. FEBS<br>Journal, 1997, 243, 134-140.   | 0.2 | 140       |
| 13 | Human IP-9: A Keratinocyte-Derived High Affinity CXC-Chemokine Ligand for the IP-10/Mig Receptor (CXCR3)1. Journal of Investigative Dermatology, 1999, 112, 716-722.  | 0.3 | 140       |
| 14 | The Human Cytomegalovirus–Encoded Chemokine Receptor US28 Promotes Angiogenesis and Tumor<br>Formation via Cyclooxygenase-2. Cancer Research, 2009, 69, 2861-2869.  | 0.4 | 139       |
| 15 | Evidence for a conformational polymorphism of invertebrate neurohormones. D-amino acid residue in crustacean hyperglycemic peptides. Journal of Biological Chemistry, 1994, 269, 18295-18298.                                 | 1.6 | 139       |
| 16 | Gene-expression profiling and array-based CGH classify CD4+CD56+ hematodermic neoplasm and cutaneous myelomonocytic leukemia as distinct disease entities. Blood, 2007, 109, 1720-1727.                                       | 0.6 | 137       |
| 17 | Array-Based Comparative Genomic Hybridization Analysis Reveals Recurrent Chromosomal Alterations<br>and Prognostic Parameters in Primary Cutaneous Large B-Cell Lymphoma. Journal of Clinical<br>Oncology, 2006, 24, 296-305. | 0.8 | 125       |
| 18 | The CXCR3 Activating Chemokines IP-10, Mig, and IP-9 are Expressed in Allergic but not in Irritant Patch<br>Test Reactions. Journal of Investigative Dermatology, 1999, 113, 574-578.   | 0.3 | 116       |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | MicroRNA-21 Expression in CD4+ T Cells Is Regulated by STAT3 and Is Pathologically Involved in Sézary<br>Syndrome. Journal of Investigative Dermatology, 2011, 131, 762-768.  | 0.3 | 116       |
| 20 | The CXCR3 Targeting Chemokine CXCL11 Has Potent Antitumor Activity In Vivo Involving Attraction of CD8+ T Lymphocytes But Not Inhibition of Angiogenesis. Journal of Immunotherapy, 2005, 28, 343-351.  | 1.2 | 114       |
| 21 | The antipsoriatic drug dimethylfumarate strongly suppresses chemokine production in human<br>keratinocytes and peripheral blood mononuclear cells. British Journal of Dermatology, 2001, 144,<br>1114-1120.   | 1.4 | 111       |
| 22 | Co-evolution of Ligand-Receptor Pairs in the Vasopressin/Oxytocin Superfamily of Bioactive Peptides.<br>Journal of Biological Chemistry, 1996, 271, 3619-3626.  | 1.6 | 104       |
| 23 | Cloning and sequence analysis of cDNA encoding two crustacean hyperglycemic hormones from the lobster Homarus americanus. FEBS Journal, 1991, 200, 103-106.   | 0.2 | 96        |
| 24 | Amino acid sequence of crustacean hyperglycemic hormone (CHH) from the crayfish, Orconectes limousus: Emergence of a novel neuropeptide family. Peptides, 1991, 12, 909-913.  | 1.2 | 91        |
| 25 | miRNA expression profiling of mycosis fungoides. Molecular Oncology, 2011, 5, 273-280.  | 2.1 | 91        |
| 26 | NPY in invertebrates: molecular answers to altered functions during evolution. Peptides, 2001, 22, 309-315.   | 1.2 | 86        |
| 27 | Expression profiling reveals that methylation of TIMP3 is involved in uveal melanoma development.<br>International Journal of Cancer, 2003, 106, 472-479.   | 2.3 | 86        |
| 28 | Fine-Mapping Chromosomal Loss at 9p21: Correlation with Prognosis in Primary Cutaneous Diffuse<br>Large B-Cell Lymphoma, Leg Type. Journal of Investigative Dermatology, 2009, 129, 1149-1155.  | 0.3 | 84        |
| 29 | Cloning, Characterization, and Expression of a G-Protein-Coupled Receptor from <i>Lymnaea<br/>stagnalis</i> and Identification of a Leucokinin-Like Peptide, PSFHSWSamide, as Its Endogenous Ligand.<br>Journal of Neuroscience, 1997, 17, 1197-1205. | 1.7 | 83        |
| 30 | Molecular profiling of cutaneous squamous cell carcinomas and actinic keratoses from organ transplant recipients. BMC Cancer, 2013, 13, 58.   | 1.1 | 83        |
| 31 | A novel G protein-coupled receptor mediating both vasopressin- and oxytocin-like functions of Lys-conopressin in Lymnaea stagnalis. Neuron, 1995, 15, 897-908.  | 3.8 | 82        |
| 32 | Evaluation of Immunophenotypic and Molecular Biomarkers for Sézary Syndrome Using Standard<br>Operating Procedures: A Multicenter Study of 59 Patients. Journal of Investigative Dermatology, 2016,<br>136, 1364-1372.                                | 0.3 | 78        |
| 33 | Increased CCL27–CCR10 expression in allergic contact dermatitis: implications for local skin memory.<br>Journal of Pathology, 2004, 204, 39-46.   | 2.1 | 77        |
| 34 | A Meta-Analysis of Gene Expression Data Identifies a Molecular Signature Characteristic for<br>Tumor-Stage Mycosis Fungoides. Journal of Investigative Dermatology, 2012, 132, 2050-2059.   | 0.3 | 75        |
| 35 | Site-Directed Mutagenesis of the Histamine H1-Receptor Reveals a Selective Interaction of Asparagine207 with Subclasses of H1-Receptor Agonists. Biochemical and Biophysical Research Communications, 1994, 201, 295-301.                             | 1.0 | 74        |
| 36 | Cucurbitacin I Inhibits Stat3 and Induces Apoptosis in Sézary Cells. Journal of Investigative Dermatology, 2008, 128, 1691-1695.  | 0.3 | 74        |

| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 37 | The <i>Lymnaea</i> Cardioexcitatory Peptide (LyCEP) Receptor: A G-Protein–Coupled Receptor for a Novel Member of the RFamide Neuropeptide Family. Journal of Neuroscience, 1998, 18, 9812-9821.  | 1.7  | 71        |
| 38 | Oligonucleotide Array-CCH Identifies Genomic Subgroups and Prognostic Markers for Tumor Stage<br>Mycosis Fungoides. Journal of Investigative Dermatology, 2010, 130, 1126-1135.  | 0.3  | 71        |
| 39 | Morphological changes during dendritic cell maturation correlate with cofilin activation and translocation to the cell membrane. European Journal of Immunology, 2004, 34, 156-164.  | 1.6  | 70        |
| 40 | Cutaneous T cell lymphoma. Nature Reviews Disease Primers, 2021, 7, 61.  | 18.1 | 70        |
| 41 | Aberrant DNA Methylation in Cutaneous Malignancies. Seminars in Oncology, 2005, 32, 479-487.   | 0.8  | 69        |
| 42 | Epidermal Interferon-Î <sup>3</sup> Inducible Protein-10 (IP-10) and Monokine Induced by Î <sup>3</sup> -Interferon (Mig) but not<br>IL-8 mRNA Expression is Associated with Epidermotropism in Cutaneous T Cell Lymphomas. Journal of<br>Investigative Dermatology, 1998, 111, 222-226. | 0.3  | 67        |
| 43 | Chromosomal Aberration Patterns Differ in Subtypes of Primary Cutaneous B Cell Lymphomas. Journal of Investigative Dermatology, 2004, 122, 1495-1502.  | 0.3  | 67        |
| 44 | Molecular cloning and characterization of an invertebrate homologue of a neuropeptide Y receptor.<br>European Journal of Neuroscience, 1998, 10, 3409-3416.  | 1.2  | 64        |
| 45 | Cutaneous Anaplastic Large Cell Lymphoma and Peripheral T-Cell Lymphoma NOS Show Distinct<br>Chromosomal Alterations and Differential Expression of Chemokine Receptors and Apoptosis<br>Regulators. Journal of Investigative Dermatology, 2010, 130, 563-575.                           | 0.3  | 62        |
| 46 | Cloning and characterization of dominant negative splice variants of the human histamine H4 receptor. Biochemical Journal, 2008, 414, 121-131.   | 1.7  | 61        |
| 47 | Haploinsufficiency for NR3C1, the gene encoding the glucocorticoid receptor, in blastic plasmacytoid dendritic cell neoplasms. Blood, 2016, 127, 3040-3053.  | 0.6  | 60        |
| 48 | A genomic and expression study of APâ€1 in primary cutaneous Tâ€cell lymphoma: evidence for<br>dysregulated expression of JUNB and JUND in MF and SS. Journal of Cutaneous Pathology, 2008, 35,<br>899-910.  | 0.7  | 57        |
| 49 | Noncompetitive Antagonism and Inverse Agonism as Mechanism of Action of Nonpeptidergic<br>Antagonists at Primate and Rodent CXCR3 Chemokine Receptors. Journal of Pharmacology and<br>Experimental Therapeutics, 2008, 325, 544-555.   | 1.3  | 57        |
| 50 | Differences in Structure–Function Relations between Nonmammalian and Mammalian<br>Gonadotropin-Releasing Hormone Receptors. Biochemical and Biophysical Research Communications,<br>1997, 238, 517-522.  | 1.0  | 56        |
| 51 | Chemokine IP-10 expression in cultured human keratinocytes. Archives of Dermatological Research, 1998, 290, 335-341.   | 1.1  | 56        |
| 52 | Genomic analysis reveals recurrent deletion of JAKâ€STAT signaling inhibitors <i>HNRNPK</i> and <i>SOCS1</i> in mycosis fungoides. Genes Chromosomes and Cancer, 2018, 57, 653-664.  | 1.5  | 56        |
| 53 | Nuclear Factor-κB Pathway–Activating Gene Aberrancies in Primary Cutaneous Large B-Cell Lymphoma,<br>Leg Type. Journal of Investigative Dermatology, 2014, 134, 290-292.   | 0.3  | 54        |
| 54 | Genomeâ€wide promoter methylation analysis identifies epigenetic silencing of<br><scp><i>MAPK</i></scp> <i>13</i> in primary cutaneous melanoma. Pigment Cell and Melanoma<br>Research, 2013, 26, 542-554.   | 1.5  | 52        |

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | A Lack of Birbeck Granules in Langerhans Cells Is Associated with a Naturally Occurring Point<br>Mutation in the Human Langerin Gene. Journal of Investigative Dermatology, 2005, 124, 714-717.  | 0.3 | 51        |
| 56 | Primary cutaneous follicle center lymphoma and primary cutaneous large B-cell lymphoma, leg type,<br>are both targeted by aberrant somatic hypermutation but demonstrate differential expression of AID.<br>Blood, 2006, 107, 4926-4929. | 0.6 | 51        |
| 57 | Multiple release of peptides by electrically active neurosecretory caudo-dorsal cells of Lymnaea stagnalis. Neuroscience Letters, 1983, 41, 151-155.   | 1.0 | 49        |
| 58 | Genomic organization, sequence and transcriptional regulation of the human CXCL 11 gene.<br>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms, 1999, 1446, 167-172.   | 2.4 | 48        |
| 59 | A novel splice variant of the Fas gene in patients with cutaneous T-cell lymphoma. Cancer Research, 2002, 62, 5389-92.   | 0.4 | 48        |
| 60 | Primary cutaneous anaplastic large cell lymphoma shows a distinct mi <scp>RNA</scp> expression<br>profile and reveals differences from tumorâ€stage mycosis fungoides. Experimental Dermatology, 2012,<br>21, 632-634.                   | 1.4 | 47        |
| 61 | An <i>in vitro</i> threeâ€dimensional model of primary human cutaneous squamous cell carcinoma.<br>Experimental Dermatology, 2009, 18, 849-856.  | 1.4 | 46        |
| 62 | Epigenomic Analysis of Sézary Syndrome Defines Patterns of Aberrant DNA Methylation and Identifies<br>DiagnosticÂMarkers. Journal of Investigative Dermatology, 2016, 136, 1876-1884.  | 0.3 | 46        |
| 63 | Localization of messenger RNAs encoding crustacean hyperglycemic hormone and gonad inhibiting<br>hormone in the X-organ sinus gland complex of the lobster Homarus americanus. Neuroscience, 1992,<br>51, 121-128.                       | 1.1 | 45        |
| 64 | Proteomic analysis of skin irritation reveals the induction of HSP27 by sodium lauryl sulphate in human skin. British Journal of Dermatology, 2002, 146, 777-785.  | 1.4 | 45        |
| 65 | Chemokine/chemokine receptor interactions in extramedullary leukaemia of the skin in childhood<br>AML: Differential roles for CCR2, CCR5, CXCR4 and CXCR7. Pediatric Blood and Cancer, 2010, 55, 344-348.                                | 0.8 | 45        |
| 66 | Deep-Sequencing Analysis Reveals that the miR-199a2/214 Cluster within DNM3os Represents the Vast<br>Majority of Aberrantly Expressed MicroRNAs in Sézary Syndrome. Journal of Investigative<br>Dermatology, 2012, 132, 1520-1522.       | 0.3 | 42        |
| 67 | Title is missing!. Fish Physiology and Biochemistry, 1997, 17, 99-108.   | 0.9 | 41        |
| 68 | Genetic rearrangements result in altered gene expression and novel fusion transcripts in Sézary<br>syndrome. Oncotarget, 2017, 8, 39627-39639.   | 0.8 | 41        |
| 69 | Differential Expression of Thymus and Activation Regulated Chemokine and Its Receptor CCR4 in Nodal<br>and Cutaneous Anaplastic Large-Cell Lymphomas and Hodgkin's Disease. Modern Pathology, 2002, 15,<br>838-844.                      | 2.9 | 38        |
| 70 | An Integrated Data Resource for Genomic AnalysisÂof Cutaneous T-Cell Lymphoma. Journal of<br>Investigative Dermatology, 2018, 138, 2681-2683.  | 0.3 | 38        |
| 71 | Early and late effects of the immunosuppressants rapamycin and mycophenolate mofetil on UV carcinogenesis. International Journal of Cancer, 2010, 127, 796-804.  | 2.3 | 37        |
| 72 | Autocrine IL-21 Stimulation Is Involved in the Maintenance of Constitutive STAT3 Activation in Sézary Syndrome. Journal of Investigative Dermatology, 2012, 132, 440-447.  | 0.3 | 37        |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | Proteomic Profiling Identifies an UV-Induced Activation of Cofilin-1 and Destrin in Human Epidermis.<br>Journal of Investigative Dermatology, 2005, 124, 818-824.   | 0.3 | 35        |
| 74 | Synthesis and structure–activity relationship of 3-phenyl-3H-quinazolin-4-one derivatives as CXCR3 chemokine receptor antagonists. Bioorganic and Medicinal Chemistry Letters, 2005, 15, 2910-2913.               | 1.0 | 32        |
| 75 | Proteomic Analysis of Uveal Melanoma Reveals Novel Potential Markers Involved in Tumor<br>Progression. , 2006, 47, 786.   |     | 32        |
| 76 | Parasites flicking the NPY gene on the host's switchboard: why NPY?. FASEB Journal, 1999, 13, 1972-1984.  | 0.2 | 32        |
| 77 | Furin Is a Chemokine-modifying Enzyme. Journal of Biological Chemistry, 2004, 279, 13402-13411.   | 1.6 | 30        |
| 78 | Functional characterization of cancerâ€associated fibroblasts of human cutaneous squamous cell<br>carcinoma. Experimental Dermatology, 2011, 20, 737-742.   | 1.4 | 29        |
| 79 | Isolation and amino acid sequence of crustacean hyperglycemic hormone precursor-related peptides.<br>Peptides, 1991, 12, 673-681.   | 1.2 | 28        |
| 80 | Improved Sézary cell detection and novel insights into immunophenotypic and molecular<br>heterogeneity in Sézary syndrome. Blood, 2021, 138, 2539-2554.   | 0.6 | 28        |
| 81 | Genetic and epigenetic insights into cutaneous T-cell lymphoma. Blood, 2022, 139, 15-33.  | 0.6 | 28        |
| 82 | Clinical and pathogenic aspects of the severe cutaneous adverse reaction epidermal necrolysis (EN).<br>Journal of the European Academy of Dermatology and Venereology, 2020, 34, 1957-1971.                       | 1.3 | 25        |
| 83 | Detection of mRNA encoding Crustacean Hyperglycemic Hormone (CHH) in the eyestalk of the crayfish<br>Orconectes limosus using non-radioactive in situ hybridization. Neuroscience Letters, 1991, 124,<br>178-182. | 1.0 | 23        |
| 84 | Fragment based lead discovery of small molecule inhibitors for the EPHA4 receptor tyrosine kinase.<br>European Journal of Medicinal Chemistry, 2012, 47, 493-500.   | 2.6 | 23        |
| 85 | Exploring the IL-21–STAT3 Axis as Therapeutic Target for Sézary Syndrome. Journal of Investigative<br>Dermatology, 2014, 134, 2639-2647.  | 0.3 | 23        |
| 86 | Molecular advances in cutaneous T-cell lymphoma. Seminars in Cutaneous Medicine and Surgery, 2018,<br>37, 81-86.  | 1.6 | 22        |
| 87 | Processing of natural and recombinant CXCR3-targeting chemokines and implications for biological activity. FEBS Journal, 2001, 268, 4992-4999.  | 0.2 | 21        |
| 88 | Reduced ILâ€1Ra/ILâ€1 ratio in ultraviolet Bâ€exposed skin of patients with polymorphic light eruption.<br>Experimental Dermatology, 2009, 18, 212-217.   | 1.4 | 21        |
| 89 | Crystal structure of the EphA4 protein tyrosine kinase domain in the apo- and dasatinib-bound state.<br>FEBS Letters, 2011, 585, 3593-3599.   | 1.3 | 21        |
| 90 | Comparative characterization of hyperglycemic neuropeptides from the lobster Homarus americanus.<br>Peptides, 1991, 12, 241-249.  | 1.2 | 20        |

| #   | Article   | IF    | CITATIONS |
|-----|---|-------|-----------|
| 91  | Profiling of apoptosis genes identifies distinct types of primary cutaneous large B cell lymphoma.<br>Journal of Pathology, 2008, 215, 340-346.   | 2.1   | 20        |
| 92  | MicroRNA Profiling of Primary Cutaneous Large B-Cell Lymphomas. PLoS ONE, 2013, 8, e82471.  | 1.1   | 20        |
| 93  | Deregulation of JAK2 signaling underlies primary cutaneous CD8 <sup>+</sup> aggressive<br>epidermotropic cytotoxic T-cell lymphoma. Haematologica, 2022, 107, 702-714.  | 1.7   | 20        |
| 94  | Title is missing!. Fish Physiology and Biochemistry, 1997, 17, 45-51.   | 0.9   | 19        |
| 95  | NOTCH1 Signaling as a Therapeutic Target in Sézary Syndrome. Journal of Investigative Dermatology, 2012, 132, 2810-2817.  | 0.3   | 18        |
| 96  | A novel mouse model for <scp>S</scp> ézary <scp>S</scp> yndrome using xenotransplantation of<br><scp>S</scp> ézary cells into immunodeficient <scp>RAG</scp> 2 <sup>â^'/â^'</sup> l³c <sup>â^'/â^'</sup> mice<br>Experimental Dermatology, 2012, 21, 706-709. | e.1.4 | 18        |
| 97  | Diagnostic and prognostic significance of <i>CDKN2A</i> / <i>CDKN2B</i> deletions in patients with transformed mycosis fungoides and primary cutaneous CD30-positive lymphoproliferative disease. British Journal of Dermatology, 2015, 172, 784-788.         | 1.4   | 18        |
| 98  | Transmitter identification in neurons involved in male copulation behavior inLymnaea stagnalis.<br>Journal of Comparative Neurology, 1998, 395, 440-449.  | 0.9   | 17        |
| 99  | The Mutational Landscape of CTCL and Sezary Syndrome. Blood, 2015, 126, 573-573.  | 0.6   | 17        |
| 100 | IP-10 mRNA EXPRESSION IN CULTURED KERATINOCYTES IS SUPPRESSED BY INHIBITION OF PROTEIN KINASE-C AND TYROSINE KINASE AND ELEVATION OF cAMP. Cytokine, 1999, 11, 469-475.   | 1.4   | 14        |
| 101 | Rapamycin impairs UV induction of mutantâ€p53 overexpressing cell clusters without affecting tumor onset. International Journal of Cancer, 2012, 131, 1267-1276.  | 2.3   | 14        |
| 102 | Pathogenesis of Skin Carcinomas and a Stem Cell as Focal Origin. Frontiers in Medicine, 2018, 5, 165.   | 1.2   | 14        |
| 103 | Wholeâ€genome analysis uncovers recurrent <i>IKZF1</i> inactivation and aberrant cell adhesion in blastic plasmacytoid dendritic cell neoplasm. Genes Chromosomes and Cancer, 2020, 59, 295-308.  | 1.5   | 14        |
| 104 | Performance of the N/TERT epidermal model for skin sensitizer identification via Nrf2-Keap1-ARE pathway activation. Toxicology in Vitro, 2014, 28, 982-989.   | 1.1   | 12        |
| 105 | No TP63 rearrangements in a selected group of primary cutaneous CD30+ lymphoproliferative disorders with aggressive clinical course. Blood, 2016, 128, 141-143.   | 0.6   | 12        |
| 106 | Acquired N-Linked Glycosylation Motifs in B-Cell Receptors of Primary Cutaneous B-Cell Lymphoma and the Normal B-Cell Repertoire. Journal of Investigative Dermatology, 2019, 139, 2195-2203.   | 0.3   | 12        |
| 107 | Cucurbitacin E and I target the JAK/STAT pathway and induce apoptosis in Sézary cells. Biochemistry and Biophysics Reports, 2020, 24, 100832.   | 0.7   | 12        |
| 108 | Cell-of-origin classification using the Hans and Lymph2Cx algorithms in primary cutaneous large<br>B-cell lymphomas. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische<br>Medizin, 2022, 480, 667-675.                             | 1.4   | 12        |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Arrayâ€based CGH of primary cutaneous CD8+ aggressive EPIDERMOâ€tropic cytotoxic Tâ€cell lymphoma.<br>Genes Chromosomes and Cancer, 2018, 57, 622-629.  | 1.5 | 11        |
| 110 | Dietary Immunosuppressants Do Not Enhance UV-Induced Skin Carcinogenesis, and Reveal Discordance between p53-Mutant Early Clones and Carcinomas. Cancer Prevention Research, 2013, 6, 129-138.            | 0.7 | 9         |
| 111 | PLCG1 Gene Mutations in Cutaneous T-Cell Lymphomas Revisited. Journal of Investigative Dermatology, 2015, 135, 2153-2154.   | 0.3 | 9         |
| 112 | Whole-genome profiling of primary cutaneous anaplastic large cell lymphoma. Haematologica, 2022,<br>107, 1619-1632.   | 1.7 | 9         |
| 113 | Lgr5+ stem cells and their progeny in mouse epidermis under regimens of exogenous skin carcinogenesis, and their absence in ensuing skin tumors. Oncotarget, 2016, 7, 52085-52094.                        | 0.8 | 8         |
| 114 | Lgr6+ stem cells and their progeny in mouse epidermis under regimens of exogenous skin carcinogenesis, and their absence in ensuing skin tumors. Oncotarget, 2016, 7, 86740-86754.                        | 0.8 | 7         |
| 115 | Azathioprine-Induced Microsatellite Instability Is Not Observed in Skin Carcinomas of Organ<br>Transplant Recipients. Journal of Investigative Dermatology, 2009, 129, 1307-1309.                         | 0.3 | 6         |
| 116 | Micro RNA â€155 potentiates tumour development in mycosis fungoides. British Journal of Dermatology,<br>2017, 177, 618-620.   | 1.4 | 6         |
| 117 | EPHA4 is overexpressed but not functionally active in Sézary syndrome. Oncotarget, 2015, 6, 31868-31876.  | 0.8 | 6         |
| 118 | No Acceleration of UV-Induced Skin Carcinogenesis from Evenly Spread Dietary Intake of Cyclosporine in Contrast to Oral Bolus Dosages. Transplantation, 2013, 96, 871-876.                                | 0.5 | 4         |
| 119 | Primary Cutaneous Follicle Center Lymphomas (PCFCL) Express Heavily Mutated B-Cell Receptors with<br>Acquired N-Glycosylation Motifs and Lack Ongoing Somatic Hypermutation. Blood, 2018, 132, 1573-1573. | 0.6 | 3         |
| 120 | Serum and cutaneous transcriptional expression levels of IL31 are minimal in cutaneous T cell lymphoma variants. Biochemistry and Biophysics Reports, 2021, 26, 101007.                                   | 0.7 | 2         |
| 121 | Molecular Cloning of a Gonadotropin-Releasing Hormone Receptor cDNA from the Red Sea Bream,<br>Pagrus major. Annals of the New York Academy of Sciences, 1998, 839, 518-519.                              | 1.8 | 1         |
| 122 | Chemical Sensitization. , 2014, , 67-87.  |     | 1         |
| 123 | <scp>RNA</scp> â€seq analysis of Lgr6 <sup>+</sup> stem cells and identification of an Lgr6 isoform.<br>Experimental Dermatology, 2018, 27, 1172-1175.  | 1.4 | 1         |
| 124 | Differential expression of CXCR3 targeting chemokines CXCL10, CXCL9, and CXCL11 in different types of skin inflammation. , 2001, 194, 398.  |     | 1         |
| 125 | Antigen-Independent, Autonomous B-Cell Receptor Signaling As a Dominant Candidate Oncogenic<br>Mechanism in ABC DLBCL. Blood, 2016, 128, 778-778.   | 0.6 | 1         |
| 126 | Molecular Cloning and Neuronal Expression of a Novel Type of a G-Protein-Coupled Receptor With Ldl<br>Binding Motifs From the Pond Snail Lymnaea Stagnalis. Animal Biology, 1993, 44, 463-472.            | 0.4 | 0         |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 127 | PTPRG (protein tyrosine phosphatase, receptor type, G). Atlas of Genetics and Cytogenetics in Oncology and Haematology, 2011, , .   | 0.1 | 0         |
| 128 | Genome Wide Analysis of 41 Mycosis Fungoides Tumor Stage Using Array Comparative Genomic<br>Hybridization Technology Blood, 2008, 112, 1769-1769.   | 0.6 | 0         |
| 129 | The B-Cell Receptor of Primary Cutaneous Follicle Center Lymphoma: Implications for Pathogenesis.<br>Blood, 2016, 128, 4136-4136.   | 0.6 | 0         |
| 130 | Tumor Clone Frequency Calculation Using High-Throughput Sequencing of the TCRÎ <sup>2</sup> Gene in Patients with Folliculotropic Mycosis Fungoides. Journal of Investigative Dermatology, 2022, 142, 2544-2546.e2. | 0.3 | 0         |