

Celine Harmanus

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4946671/publications.pdf>

Version: 2024-02-01

27
papers

974
citations

516215

16
h-index

525886

27
g-index

28
all docs

28
docs citations

28
times ranked

938
citing authors

#	ARTICLE	IF	CITATIONS
1	InÂvitro anti-clostridial action and potential of the spice herbs essential oils to prevent biofilm formation of hypervirulent Clostridioides difficile strains isolated from hospitalized patients with CDI. Anaerobe, 2022, 76, 102604.	1.0	2
2	Haem is crucial for medium-dependent metronidazole resistance in clinical isolates of Clostridioides difficile. Journal of Antimicrobial Chemotherapy, 2021, 76, 1731-1740.	1.3	34
3	Detection of Clostridioides difficile in hospital environment by using C diff Banana Brothâ,,Ç. Anaerobe, 2021, , 102408.	1.0	3
4	Clostridioides difficile Ribotype 027 (RT027) Outbreak Investigation Due to the Emergence of Rifampicin Resistance Using Multilocus Variable-Number Tandem Repeat Analysis (MLVA). Infection and Drug Resistance, 2021, Volume 14, 3247-3254.	1.1	5
5	The Bacterial Gut Microbiota of Adult Patients Infected, Colonized or Noncolonized by Clostridioides difficile. Microorganisms, 2020, 8, 677.	1.6	25
6	Plasmid-mediated metronidazole resistance in Clostridioides difficile. Nature Communications, 2020, 11, 598.	5.8	79
7	Detection of Clostridium difficile in the environment in a veterinary teaching hospital. Anaerobe, 2019, 57, 55-58.	1.0	5
8	A helicase-containing module defines a family of pCD630-like plasmids in Clostridium difficile. Anaerobe, 2018, 49, 78-84.	1.0	13
9	Recreational sandboxes for children and dogs can be a source of epidemic ribotypes of Clostridium difficile. Zoonoses and Public Health, 2018, 65, 88-95.	0.9	24
10	Characterization of the virulence of a non-RT027, non-RT078 and binary toxin-positive Clostridium difficile strain associated with severe diarrhea. Emerging Microbes and Infections, 2018, 7, 1-11.	3.0	17
11	Quantification of Clostridioides (Clostridium) difficile in feces of calves of different age and determination of predominant Clostridioides difficile ribotype 033 relatedness and transmission between family dairy farms using multilocus variable-number tandem-repeat analysis. BMC Veterinary Research, 2018, 14, 298.	0.7	17
12	Distribution and tracking of Clostridium difficile and Clostridium perfringens in a free-range pig abattoir and processing plant. Food Research International, 2018, 113, 456-464.	2.9	9
13	First molecular characterisation and PCR ribotyping of Clostridium difficile strains isolated in two Algerian Hospitals. Journal of Infection in Developing Countries, 2018, 12, 015-021.	0.5	9
14	Occurrence of Clostridium difficile ribotype 027 in hospitals of Silesia, Poland. Anaerobe, 2017, 45, 106-113.	1.0	25
15	Subtyping and antimicrobial susceptibility of Clostridium difficile PCR ribotype 078/126 isolates of human and animal origin. Veterinary Microbiology, 2017, 199, 15-22.	0.8	38
16	Isolation of Clostridium difficile from dogs with digestive disorders, including stable metronidazole-resistant strains. Anaerobe, 2017, 43, 78-81.	1.0	37
17	PCR-ribotype distribution of Clostridium difficile in Irish pigs. Anaerobe, 2017, 48, 237-241.	1.0	16
18	Prevalence and characteristics of Clostridium perfringens and Clostridium difficile in dogs and cats attended in diverse veterinary clinics from the Madrid region. Anaerobe, 2017, 48, 47-55.	1.0	31

#	ARTICLE	IF	CITATIONS
19	Data from a survey of <i>Clostridium perfringens</i> and <i>Clostridium difficile</i> shedding by dogs and cats in the Madrid region (Spain), including phenotypic and genetic characteristics of recovered isolates. <i>Data in Brief</i> , 2017, 14, 88-100.	0.5	3
20	Ribotype 078 <i>Clostridium difficile</i> infection incidence in Dutch hospitals is not associated with provincial pig farming: Results from a national sentinel surveillance, 2009-2015. <i>PLoS ONE</i> , 2017, 12, e0189183.	1.1	8
21	Toxigenic <i>Clostridium difficile</i> PCR ribotypes in edible marine bivalve molluscs in Italy. <i>International Journal of Food Microbiology</i> , 2015, 208, 30-34.	2.1	32
22	Development and Validation of an Internationally-Standardized, High-Resolution Capillary Gel-Based Electrophoresis PCR-Ribotyping Protocol for <i>Clostridium difficile</i> . <i>PLoS ONE</i> , 2015, 10, e0118150.	1.1	176
23	Shedding of <i>Clostridium difficile</i> PCR ribotype 078 by zoo animals, and report of an unstable metronidazole-resistant isolate from a zebra foal (<i>Equus quagga burchellii</i>). <i>Veterinary Microbiology</i> , 2014, 169, 218-222.	0.8	32
24	Occurrence of <i>Clostridium difficile</i> PCR-ribotype 027 and it's closely related PCR-ribotype 176 in hospitals in Poland in 2008â€“2010. <i>Anaerobe</i> , 2014, 28, 13-17.	1.0	29
25	Characterization and antimicrobial susceptibility of <i>Clostridium difficile</i> strains isolated from adult patients with diarrhoea hospitalized in two university hospitals in Poland, 2004â€“2006. <i>Journal of Medical Microbiology</i> , 2011, 60, 1200-1205.	0.7	22
26	Genetic markers for <i>Clostridium difficile</i> lineages linked to hypervirulence. <i>Microbiology (United Kingdom)</i> , 2014, 158, 1052-1058.	0.7	52
27	Emergence of reduced susceptibility to metronidazole in <i>Clostridium difficile</i> . <i>Journal of Antimicrobial Chemotherapy</i> , 2008, 62, 1046-1052.	1.3	230