

# Pere Abello

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/494863/publications.pdf>

Version: 2024-02-01

117  
papers

2,713  
citations

201674  
27  
h-index

233421  
45  
g-index

119  
all docs

119  
docs citations

119  
times ranked

2465  
citing authors

#	ARTICLE	IF	CITATIONS
1	Severe, rapid and widespread impacts of an Atlantic blue crab invasion. <i>Marine Pollution Bulletin</i> , 2022, 176, 113479.	5.0	18
2	Interannual fluctuations in connectivity among crab populations ( <i>Liocarcinus depurator</i> ) along the Atlantic-Mediterranean transition. <i>Scientific Reports</i> , 2022, 12, .	3.3	4
3	Effects on faunistic composition and population characteristics of decapod crustaceans after the implementation of a fisheries no-take area in the NW Mediterranean. <i>Scientia Marina</i> , 2022, 86, e035.	0.6	2
4	Shape matters: relevance of carapace for brachyuran crab invaders. <i>Biological Invasions</i> , 2021, 23, 461-475.	2.4	6
5	Benthic marine litter in shallow fishing grounds in the NW Mediterranean Sea. <i>Waste Management</i> , 2019, 95, 620-627.	7.4	20
6	Species delimitation and multi-locus species tree solve an old taxonomic problem for European squat lobsters of the genus <i>Munida</i> Leach, 1820. <i>Marine Biodiversity</i> , 2019, 49, 1751-1773.	1.0	16
7	A new species of pea crab from south-western Europe (Crustacea, Decapoda, Brachyura): species description, geographic distribution and population structure with an identification key to European Pinnotheridae. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2019, 99, 1141-1152.	0.8	4
8	Contrasting patterns in the vertical distribution of decapod crustaceans throughout ontogeny. <i>Hydrobiologia</i> , 2018, 808, 137-152.	2.0	6
9	Updating changes in the Iberian decapod crustacean fauna (excluding crabs) after 50 years. <i>Scientia Marina</i> , 2018, 82, 207.	0.6	7
10	Possible amphi-Atlantic dispersal of <i>Scyllarus</i> lobsters (Crustacea: Scyllaridae): molecular and larval evidence. <i>Zootaxa</i> , 2017, 4306, .	0.5	10
11	Sexual dimorphism and associated population characteristics in the benthopelagic shrimp &lt;em&gt; <i>Pasiphaea sivado</i> &lt;/em&gt; (Crustacea: Caridea: Pasiphaeidae). <i>Scientia Marina</i> , 2017, 81, 57.	0.6	0
12	Temporal and spatial genetic differentiation in the crab <i>Liocarcinus depurator</i> across the Atlantic-Mediterranean transition. <i>Scientific Reports</i> , 2016, 6, 29892.	3.3	26
13	Comparative phylogeography of two symbiotic dorvilleid polychaetes (<i>Iphitime</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 267 Zoological Journal of the Linnean Society, 2016, , .	2.3	4
14	Pasiphaeid shrimps in the western Mediterranean: geographical variability in distribution and population patterns. <i>Scientia Marina</i> , 2015, 79, 199-209.	0.6	3
15	Annotated checklist of brachyuran crabs (Crustacea: Decapoda) of the Iberian Peninsula (SW Europe). <i>Scientia Marina</i> , 2015, 79, 243-256.	0.6	23
16	<i>Maja brachydactyla</i> (Brachyura: Majidae) in the western Mediterranean. <i>Marine Biodiversity Records</i> , 2014, 7, .	1.2	10
17	Reprint of “Deep epibenthic communities in two contrasting areas of the Balearic Islands (western) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 267 Zoological Journal of the Linnean Society, 2016, , .”	2.1	11
18	Body condition of the deep water demersal resources at two adjacent oligotrophic areas of the western Mediterranean and the influence of the environmental features. <i>Journal of Marine Systems</i> , 2014, 138, 194-202.	2.1	11

#	ARTICLE	IF	CITATIONS
19	Larval stages of the deep-sea lobster <i>Polycheles typhlops</i> (Decapoda, Polychelida) identified by DNA analysis: morphology, systematic, distribution and ecology. <i>Helgoland Marine Research</i> , 2014, 68, 379-397.	1.3	19
20	Vertical and temporal distribution of pelagic decapod crustaceans over the shelf-break and middle slope in two contrasting zones around Mallorca (western Mediterranean Sea). <i>Journal of Marine Systems</i> , 2014, 138, 139-149.	2.1	11
21	Deep epibenthic communities in two contrasting areas of the Balearic Islands (western) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 6	2.1	
22	The systematic position of <i>Ergasticus</i> (Decapoda, Brachyura) and allied genera, a molecular and morphological approach. <i>Zoologica Scripta</i> , 2013, 42, 427-439.	1.7	24
23	The tropical African hermit crab <i>Pagurus mbizi</i> (Crustacea, Decapoda, Paguridae) in the Western Mediterranean Sea: a new alien species or filling gaps in the knowledge of the distribution?. <i>Mediterranean Marine Science</i> , 2013, 15, 172.	1.6	9
24	Enrique Macpherson Mayol. <i>Scientia Marina</i> , 2013, 77, 9-12.	0.6	0
25	Phylogeographic patterns of decapod crustaceans at the Atlanticâ€“Mediterranean transition. <i>Molecular Phylogenetics and Evolution</i> , 2012, 62, 664-672.	2.7	59
26	Permanent Genetic Resources added to Molecular Ecology Resources Database 1 April 2012 â€“ 31 May 2012. <i>Molecular Ecology Resources</i> , 2012, 12, 972-974.	4.8	18
27	First record and evidence of an established population of the North American mud crab <i>Dyspanopeus sayi</i> (Brachyura: Heterotremata: Panopeidae) in the western Mediterranean. <i>Scientia Marina</i> , 2012, 76, 79-85.	0.6	16
28	Early juvenile development of Mediterranean <i>Liocarcinus depurator</i> (Crustacea: Decapoda: Brachyura:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 0.5	0.5	
29	Demersal Assemblages on the Soft Bottoms off the Catalan-Levante Coast of the Spanish Mediterranean. <i>Journal of Marine Biology</i> , 2011, 2011, 1-16.	1.0	11
30	Effect of oceanographic barriers and overfishing on the population genetic structure of the European spiny lobster ( <i>Palinurus elephas</i> ). <i>Biological Journal of the Linnean Society</i> , 2011, 104, 407-418.	1.6	30
31	Long-term changes in the abundance and deepening of the deep-sea shrimp <i>Aristaeomorpha foliacea</i> in the Balearic Basin: Relationships with hydrographic changes at the Levantine Intermediate Water. <i>Journal of Marine Systems</i> , 2011, 88, 516-525.	2.1	23
32	<i>Scyllarus arctus</i> (Crustacea: Decapoda: Scyllaridae) final stage phyllosoma identified by DNA analysis, with morphological description. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2011, 91, 485-492.	0.8	12
33	Faunistic assemblages of a sublittoral coarse sand habitat of the northwestern Mediterranean. <i>Scientia Marina</i> , 2011, 75, 189-196.	0.6	11
34	Genetic Diversity Levels in Fishery-Exploited Spiny Lobsters of the Genus <i>Palinurus</i> (Decapoda:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 14.0	0.8	
35	Final-stage phyllosoma of <i>Palinustus</i> A. Milne-Edwards, 1880 (Crustacea: Decapoda: Achelata:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 14.0	0.5	
36	First record of <i>Percnon gibbesi</i> (H. Milne Edwards, 1853) (Crustacea: Decapoda: Percnidae) from Egyptian waters. <i>Aquatic Invasions</i> , 2010, 5, S123-S125.	1.6	8

#	ARTICLE	IF	CITATIONS
37	Rapid radiation in spiny lobsters ( <i>Palinurus</i> spp) as revealed by classic and ABC methods using mtDNA and microsatellite data. <i>BMC Evolutionary Biology</i> , 2009, 9, 263.	3.2	31
38	Phylogenetic relationships between spiny, slipper and coral lobsters (Crustacea, Decapoda, Achelata). <i>Molecular Phylogenetics and Evolution</i> , 2009, 50, 152-162.	2.7	65
39	The true identities of the slipper lobsters <i>Nisto laevis</i> and <i>Nisto asper</i> (Crustacea: Decapoda) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 12	1.8	TF
40	Phylogeography of the European spiny lobster ( <i>Palinurus elephas</i> ): Influence of current oceanographical features and historical processes. <i>Molecular Phylogenetics and Evolution</i> , 2008, 48, 708-717.	2.7	82
41	Morphology of the final stage phyllosoma larva of <i>Scyllarus pygmaeus</i> (Crustacea: Decapoda) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 24	1.8	TF
42	Morphology of the larval stages of the spider crab <i>Herbstia condyiata</i> (Fabricius, 1787) (Brachyura:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	0.5	2
43	Occurrence of Automate branchialis Holthuis & Gottlieb, 1958 (Decapoda, Alpheidae) in the Balearic Islands (western Mediterranean Sea). <i>Crustaceana</i> , 2007, 80, 495-501.	0.3	2
44	The first phyllosoma stage of <i>Palinurus mauritanicus</i> (Crustacea: Decapoda: Palinuridae). <i>Zootaxa</i> , 2007, 1508, 49-59.	0.5	13
45	Rhythmic diel movements of pandalid shrimps in the western Mediterranean continental shelf and upper slope. <i>Journal of Zoology</i> , 2007, 273, 340-349.	1.7	11
46	The first zoea of <i>Inachus aguiarii</i> , <i>Inachus communissimus</i> and <i>Ergasticus clouei</i> (Decapoda,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 387 T	0.5	3
47	Ontogenetic changes in vertical migratory rhythms of benthopelagic shrimps <i>Pasiphaea multidentata</i> and <i>P. sivado</i> . <i>Marine Ecology - Progress Series</i> , 2007, 335, 167-174.	1.9	20
48	OCEANOGRAPHIC HABITAT OF AN ENDANGERED MEDITERRANEAN PROCELLARIIFORM: IMPLICATIONS FOR MARINE PROTECTED AREAS. , 2006, 16, 1683-1695.		117
49	Geographic and gender shape differences in the carapace of <i>Liocarcinus depurator</i> (Brachyura:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T Zoology, 2006, 269, 458-465.	1.7	36
50	Spatial and Environmental Factors Affecting the Distribution of the Main Decapod Crustacean Prey Species in the NW Mediterranean. <i>Hydrobiologia</i> , 2006, 555, 129-141.	2.0	13
51	Geostatistical analysis of densities of <i>Liocarcinus depurator</i> (Brachyura: Portunidae) on the western Mediterranean from 1994 to 2003. <i>Marine Biology</i> , 2006, 149, 855-864.	1.5	11
52	Morphology of Puerulus and Early Juvenile Stages of the Spiny Lobster <i>Palinurus Mauritanicus</i> (Decapoda: Palinuridae). <i>Journal of Crustacean Biology</i> , 2006, 26, 480-494.	0.8	7
53	Morphological variability of the megalopa of <i>Liocarcinus depurator</i> (Brachyura: Portunidae) in Mediterranean and Atlantic populations. <i>Journal of Natural History</i> , 2006, 40, 1851-1866.	0.5	8
54	The capture of the Indo-Pacific crab <i>Charybdis feriata</i> (Linnaeus, 1758) (Brachyura: Portunidae) in the Mediterranean Sea. <i>Aquatic Invasions</i> , 2006, 1, 13-16.	1.6	24

#	ARTICLE	IF	CITATIONS
55	Information, comments and thoughts on editing <i>scientia marina</i> . <i>Scientia Marina</i> , 2006, 70, 0-0.	0.6	0
56	Increased predation of juvenile European spiny lobster ( <i>&lt; i&gt; Palinurus elephas&lt;/i&gt;</i> ) in a marine protected area. <i>New Zealand Journal of Marine and Freshwater Research</i> , 2005, 39, 447-453.	2.0	31
57	The complete larval development of the crab <i>Pilumnus spinifer</i> (Brachyura: Xanthoidea: Pilumnidae) reared in the laboratory. <i>Journal of Natural History</i> , 2005, 39, 3187-3216.	0.5	0
58	The effect of methodological options on geostatistical modelling of animal distribution: A case study with <i>Liocarcinus depurator</i> (Crustacea: Brachyura) trawl survey data. <i>Fisheries Research</i> , 2005, 76, 252-265.	1.7	16
59	Small-scale distribution characteristics of <i>Munida</i> spp. populations (Decapoda: Anomura) off the Catalan coasts (western Mediterranean). <i>Journal of Sea Research</i> , 2005, 53, 283-296.	1.6	10
60	Temporal modification in cardiac rhythmicity of <i>&lt; i&gt; Nephrops norvegicus&lt;/i&gt;</i> (Crustacea: Decapoda) in relation to trawl capture stress. <i>Scientia Marina</i> , 2005, 69, 369-374.	0.6	12
61	Geographic, bathymetric and inter-annual variability in the distribution of &lt;i&gt; <i>Liocarcinus depurator</i> &lt;/i&gt; (Brachyura: Portunidae) along the Mediterranean coast of the Iberian peninsula. <i>Scientia Marina</i> , 2005, 69, 503-518.	0.6	16
62	The effect of alcohol and freezing preservation on carapace size and shape in <i>Liocarcinus depurator</i> (Crustacea, Brachyura). , 2004, , 45-53.		6
63	A review of the fisheries biology of the mantis shrimp, <i>Squilla mantis</i> (L., 1758) (Stomatopoda,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 0.3 35		10
64	Distribution of the Neustonic Isopod <i>Idotea Metallica</i> in Relation to Shelf-Slope Frontal Structures. <i>Journal of Crustacean Biology</i> , 2004, 24, 558-566.	0.8	9
65	Male and female carapace shape differences in <i>&lt; i&gt; Liocarcinus depurator&lt;/i&gt;</i> (Decapoda, Brachyura): An application of geometric morphometric analysis to crustaceans. <i>Italian Journal of Zoology</i> , 2004, 71, 79-83.	0.6	43
66	Endogenous cardiac activity rhythms of continental slope <i>Nephrops norvegicus</i> (decapoda:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 302 Td 0.9 21		
67	Locomotor Activity Rhythms of Continental Slope <i>Nephrops Norvegicus</i> (Decapoda: Nephropidae). <i>Journal of Crustacean Biology</i> , 2004, 24, 282-290.	0.8	30
68	Small-scale non-linear geostatistical analysis of <i>Liocarcinus depurator</i> (Crustacea: Brachyura) abundance and size structure in a western Mediterranean population. <i>Marine Ecology - Progress Series</i> , 2004, 276, 223-235.	1.9	18
69	CIESM Atlas of exotic species in the Mediterranean. Vol. 2. Crustaceans: decapods and stomatopods. B. Galil, C. Froglia and P. NoÃ«l. <i>Scientia Marina</i> , 2004, 68, 189-189.	0.6	2
70	Morphology of the larval and first juvenile stages of the spider crab <i>Lissa chiragra</i> (Brachyura:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 142 0.5 5		
71	Circadian Oxygen Consumption Patterns in Continental Slope <i>Nephrops Norvegicus</i> (Decapoda:) Tj ETQq1 1 0.784314 rgBT /Overlock 0.8 22		
72	Mesoscale geographical patterns in the distribution of pandalid shrimps <i>Plesionika</i> spp. in the Western Mediterranean. <i>Marine Ecology - Progress Series</i> , 2003, 247, 151-158.	1.9	15

#	ARTICLE	IF	CITATIONS
73	Diel and seasonal patterns of <i>Nephrops norvegicus</i> (Decapoda: Nephropidae) catchability in the western Mediterranean. <i>Marine Ecology - Progress Series</i> , 2003, 258, 201-211.	1.9	78
74	Geographical patterns of seabird attendance to a research trawler along the Iberian Mediterranean coast. <i>Scientia Marina</i> , 2003, 67, 69-75.	0.6	25
75	The complete larval development and first juvenile of the spider crab <i>Inachus thoracicus</i> Roux, 1830 (Brachyura: Majidae: Inachinae). <i>Journal of Plankton Research</i> , 2002, 24, 875-887.	1.8	2
76	Commercial feasibility trial for a single-warp deep-water "Maireta" (OTMS) trawl gear. <i>Fisheries Research</i> , 2002, 55, 121-130.	1.7	3
77	Geographical patterns in abundance and population structure of <math>\text{Nephrops norvegicus}</math> and <math>\text{Parapenaeus longirostris}</math> (Crustacea: Decapoda) along the European Mediterranean coasts. <i>Scientia Marina</i> , 2002, 66, 125.	0.6	74
78	Biogeography of epibenthic crustaceans on the shelf and upper slope off the Iberian Peninsula Mediterranean coasts: implications for the establishment of natural management areas. <i>Scientia Marina</i> , 2002, 66, 183.	0.6	125
79	Feeding guilds of western Mediterranean demersal fish and crustaceans: an analysis based in a spring survey. <i>Scientia Marina</i> , 2002, 66, 209.	0.6	65
80	Title is missing!. <i>Hydrobiologia</i> , 2001, 449, 187-192.	2.0	6
81	Settlement and juvenile habitat of the European spiny lobster <math>\text{i} \text{Palinurus elephas}</math> (Crustacea:) Tj ETQq1 1 0.784314 rgBT /Overlock 0.6 31		
82	The occurrence of <i>Hymenopenaeus debilis</i> (Decapoda: Aristeidae: Solenocerinae) in Mediterranean waters: a case of pseudopopulations of Atlantic origin?. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2000, 80, 549-550.	0.8	6
83	Demersal fish assemblages and habitat characteristics on the continental shelf and upper slope of the north-western Mediterranean. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2000, 80, 981-988.	0.8	96
84	Morphology of the early zoeal stages of <i>Macropipus tuberculatus</i> (Roux, 1830) (Crustacea, Brachyura,) Tj ETQq0 0 0 rgBT /Overlock 10		
85	Temporal Variability in the Vertical and Mesoscale Spatial Distribution of Crab Megalopae (Crustacea:) Tj ETQq1 1 0.784314 rgBT /Overlock 2.1 31		
86	Endogenous tidal and semilunar moulting rhythms in early juvenile shore crabs <i>Carcinus maenas</i> : implications for adaptation to a high intertidal habitat. <i>Marine Ecology - Progress Series</i> , 1999, 191, 257-266.	1.9	21
87	The first zoeal stage of <math>\text{i} \text{Parthenope macrochelos}</math> (Herbst, 1790) hatched in the laboratory (Crustacea: Brachyura: Parthenopidae). <i>Scientia Marina</i> , 1999, 63, 9-14.	0.6	2
88	Transitory hydrographic structures and distribution of fish larvae and neustonic crustaceans in the north-western Mediterranean. <i>Oceanologica Acta: European Journal of Oceanology - Revue Europeene De Oceanologie</i> , 1998, 21, 95-104.	0.7	30
89	Distribution characteristics of pandalid shrimps (Decapoda: Caridea: Pandalidae) along the western Mediterranean Sea. <i>Journal of Natural History</i> , 1998, 32, 1463-1474.	0.5	32
90	Offshore Distribution of Seabirds in the Northwestern Mediterranean in June 1995. <i>Waterbirds</i> , 1998, 21, 422.	0.4	10

#	ARTICLE	IF	CITATIONS
91	Circatidal Moulting Rhythm in The Shore Crab <i>Carcinus Maenas</i>. Journal of the Marine Biological Association of the United Kingdom, 1997, 77, 277-279.	0.8	15
92	Larval Development of the Spider Crab Macropodia Longipes (Brachyura: Majidae: Inachinae). Journal of Crustacean Biology, 1997, 17, 459-471.	0.8	7
93	Carapace Colour, Inter-moult Duration and the Behavioural and Physiological Ecology of the Shore Crab <i>Carcinus maenas</i>. Estuarine, Coastal and Shelf Science, 1997, 44, 203-211.	2.1	98
94	Endogenous control of timing of metamorphosis in megalopae of the shore crab <i>Carcinus maenas</i>. Marine Biology, 1997, 128, 299-305.	1.5	26
95	Spatial variability in the population structure of the shore crab <i>Carcinus maenas</i> (Crustacea:Brachyura) in a shallow-water, weakly tidal fjord. Marine Ecology - Progress Series, 1997, 147, 97-103.	1.9	24
96	Morphology of the larval stages of the deep-sea crab <i>Geryon longipes</i>(Brachyura: Geryonidae). Journal of Natural History, 1996, 30, 505-521.	0.5	8
97	Larval development of <i>Acanthonyx lunulatus</i> (Brachyura, Majidae, Epialtinae). Journal of Plankton Research, 1996, 18, 1185-1200.	1.8	5
98	Description of the first larval stage of <i>Polycheles typhlops</i> (Decapoda: Eryonidea: Polychelidae). Journal of Natural History, 1996, 30, 1179-1184.	0.5	4
99	Population structure and growth of <i>Donax trunculus</i> (Bivalvia: Donacidae) in the western Mediterranean. Marine Biology, 1995, 121, 665-671.	1.5	44
100	Larval development of <i>Brachynotus gemmellari</i> (Rizza, 1839) (Brachyura, Grapsidae) reared under laboratory conditions. Journal of Plankton Research, 1995, 17, 1143-1161.	1.8	10
101	Note on the diet of <i>Lithodes ferox</i> (Anomura: Lithodidae) off Namibia. African Journal of Marine Science, 1995, 15, 273-277.	0.6	3
102	Chela loss in the shore crab <i>Carcinus maenas</i> (Crustacea: Brachyura) and its effect on mating success. Marine Biology, 1994, 121, 247-252.	1.5	66
103	Size-related mating success in the shore crab <i>Carcinus maenas</i> (Crustacea: Brachyura). Journal of Zoology, 1994, 232, 397-407.	1.7	43
104	Fishery dynamics of the mantis shrimp <i>Squilla mantis</i> (Crustacea: Stomatopoda) population off the Ebro delta (northwestern Mediterranean). Fisheries Research, 1993, 16, 131-145.	1.7	25
105	Epibionts and intermoult duration in the crab <i> Bathynectes piperitus</i>. Marine Ecology - Progress Series, 1993, 98, 107-113.	1.9	69
106	Epibiosis and Rhizocephalan Infestation Patterns in Relation to the Reproductive Biology of <i>Lithodes ferox</i> (Filhol, 1885) (Anomura: Lithodidae). Journal of Crustacean Biology, 1992, 12, 561.	0.8	28
107	Distribution Patterns and Migration of <i>Lithodes ferox</i> (Filhol) (Anomura: Lithodidae) off Namibia. Journal of Crustacean Biology, 1991, 11, 261.	0.8	10
108	A new species of <i>Ophryotrocha</i> (Polychaeta: Dorvilleidae) commensal in <i>Geryon longipes</i> (Crustacea: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 0.5		

#	ARTICLE	IF	CITATIONS
109	Comparative locomotor activity patterns in the portunid crabs <i>Liocarcinus holsatus</i> and <i>L. depurator</i> . Journal of the Marine Biological Association of the United Kingdom, 1991, 71, 1-10.	0.8	29
110	Epibiosis in deep-sea crab populations as indicator of biological and behavioural characteristics of the host. Journal of the Marine Biological Association of the United Kingdom, 1990, 70, 687-695.	0.8	69
111	Influence of environmental conditions on the distribution of <i>i&gt;Pterygosquilla armata capensis&lt;/i&gt;</i> (Crustacea: Stomatopoda) off Namibia. African Journal of Marine Science, 1990, 9, 169-175.	0.6	3
112	Reproductive biology of <i>i&gt;Macropipus tuberculatus&lt;/i&gt;</i> (Roux, 1830) (Brachyura: Portunidae) in the Northwestern Mediterranean. Ophelia, 1989, 30, 47-53.	0.3	14
113	Distribution of <i>Bathynectes piperitus</i> (Brachyura: Portunidae) in the Benguela Upwelling Region and Its Relationship with Some Environmental Parameters. Journal of Crustacean Biology, 1989, 9, 373.	0.8	2
114	Analysis of the structure of decapod crustacean assemblages off the Catalan coast (North-West) Tj ETQq0 0 0 rgBT <sub>1.5</sub> /Overlock <sub>215</sub> 10 Tf 50 5		
115	The Fecundity of the Norway Lobster ( <i>Nephrops Norvegicus</i> (L.)) Off the Catalan and Portuguese Coasts. Crustaceana, 1982, 43, 13-20.	0.3	13
116	Updating hosts and distribution range of the pea crab <i>Pinnotheres bicristatus</i> (Brachyura:) Tj ETQq0 0 0 rgBT /Overlock <sub>1.6</sub> 10 Tf <sub>2</sub> 50 462 Td		
117	Detection of a population of <i>Pseudosquillopsis cerisii</i> (Roux, 1828) (Crustacea, Stomatopoda,) Tj ETQq1 1 0.784314 rgBT <sub>0.5</sub> /Overlock <sub>10</sub>		