

Katsuhiko Maeda

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4953221/katsuhiko-maeda-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

130
papers

8,674
citations

42
h-index

92
g-index

147
ext. papers

9,741
ext. citations

8.6
avg, IF

6.26
L-index

#	Paper	IF	Citations
130	Solvent-dependent helix inversion in optically active poly(diphenylacetylene)s and their chiral recognition abilities as chiral stationary phases for high-performance liquid chromatography.. <i>Chirality</i> , 2022 ,	2.1	1
129	Speciation analysis of inorganic selenium in wastewater using a highly selective cellulose-based adsorbent via liquid electrode plasma optical emission spectrometry. <i>Journal of Hazardous Materials</i> , 2022 , 424, 127250	12.8	1
128	Synthesis of Pentaarylcyclobutenylrhodium(I) Complexes and Their Reactivity and Initiation Mechanism in Polymerization of Monosubstituted Acetylenes. <i>Organometallics</i> , 2022 , 41, 472-479	3.8	
127	Enantioseparation on Helical Poly(diphenylacetylene)s Bearing Optically-Active Pendants: Effects of Differences in Higher-Order Structures of Kinetically-Trapped and Thermodynamically-Stable States on Chiral Recognition Ability. <i>Journal of Chromatography A</i> , 2022 , 463164	4.5	0
126	Visualisation of helical structures of poly(diphenylacetylene)s bearing chiral amide pendants by atomic force microscopy. <i>Chemical Communications</i> , 2021 , 57, 12266-12269	5.8	3
125	Helical springs as a color indicator for determining chirality and enantiomeric excess. <i>Science Advances</i> , 2021 , 7,	14.3	5
124	Selective recovery of silver and palladium from acidic waste solutions using dithiocarbamate-functionalized cellulose. <i>Chemical Engineering Journal</i> , 2021 , 407, 127225	14.7	12
123	Highly selective and straightforward recovery of gold and platinum from acidic waste effluents using cellulose-based bio-adsorbent. <i>Journal of Hazardous Materials</i> , 2021 , 410, 124569	12.8	17
122	Racemic Monomer-Based One-Handed Helical Polymer Recognizes Enantiomers through Auto-Evolution of Its Helical Handedness Excess. <i>Angewandte Chemie - International Edition</i> , 2021 , 60, 4625-4632	16.4	10
121	Racemic Monomer-Based One-Handed Helical Polymer Recognizes Enantiomers through Auto-Evolution of Its Helical Handedness Excess. <i>Angewandte Chemie</i> , 2021 , 133, 4675-4682	3.6	2
120	Synthesis of Stereoregular Telechelic Poly(phenylacetylene)s: Facile Terminal Chain-End Functionalization of Poly(phenylacetylene)s by Terminative Coupling with Acrylates and Acrylamides in Rhodium-Catalyzed Living Polymerization of Phenylacetylenes. <i>Journal of the American Chemical Society</i> , 2021 , 143, 3664-3672	16.4	7
119	Emergence of Highly Enantioselective Catalytic Activity in a Helical Polymer Mediated by Deracemization of Racemic Pendants. <i>Journal of the American Chemical Society</i> , 2021 , 143, 12725-12735	16.4	9
118	Rhodium(I) Complexes Bearing an Aryl-Substituted 1,3,5-Hexatriene Chain: Catalysts for Living Polymerization of Phenylacetylene and Potential Helical Chirality of 1,3,5-Hexatrienes. <i>Angewandte Chemie - International Edition</i> , 2021 , 60, 22201-22206	16.4	1
117	Rhodium(I) Complexes Bearing an Aryl-Substituted 1,3,5-Hexatriene Chain: Catalysts for Living Polymerization of Phenylacetylene and Potential Helical Chirality of 1,3,5-Hexatrienes. <i>Angewandte Chemie</i> , 2021 , 133, 22375-22380	3.6	0
116	Understanding the Polymerization of Diphenylacetylenes with Tantalum(V) Chloride and Cocatalysts: Production of Cyclic Poly(diphenylacetylene)s by Low-Valent Tantalum Species Generated in Situ. <i>Journal of the American Chemical Society</i> , 2021 , 143, 16136-16146	16.4	0
115	Comparative evaluation of dithiocarbamate-modified cellulose and commercial resins for recovery of precious metals from aqueous matrices. <i>Journal of Hazardous Materials</i> , 2021 , 418, 126308	12.8	5
114	Radical -Hydroboration of Substituted 1,3-Diyne)s with an -Heterocyclic Carbene Borane. <i>Organic Letters</i> , 2021 , 23, 1071-1075	6.2	6

113	Revisiting the Polymerization of Diphenylacetylenes with Tungsten(VI) Chloride and Tetraphenyltin: An Alternative Mechanism by a Metathesis Catalytic System. <i>Angewandte Chemie</i> , 2020 , 132, 14882-14890	3.6	2
112	Spin Filtering Along Chiral Polymers. <i>Angewandte Chemie</i> , 2020 , 132, 14779-14784	3.6	0
111	Spin Filtering Along Chiral Polymers. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 14671-14676	16.4	20
110	Synthesis of a poly(diphenylacetylene) bearing optically active anilide pendants and its application to a chiral stationary phase for high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2020 , 1622, 461173	4.5	9
109	Facile and Versatile Synthesis of End-Functionalized Poly(phenylacetylene)s: A Multicomponent Catalytic System for Well-Controlled Living Polymerization of Phenylacetylenes. <i>Angewandte Chemie</i> , 2020 , 132, 8748-8758	3.6	6
108	Helix-Sense-Selective Synthesis of Right- and Left-Handed Helical Luminescent Poly(diphenylacetylene)s with Memory of the Macromolecular Helicity and Their Helical Structures. <i>Journal of the American Chemical Society</i> , 2020 , 142, 7668-7682	16.4	35
107	Revisiting Polyfluoroarenes as Radical Acceptors: Radical C-F Bond Borylation of Polyfluoroarenes with N-Heterocyclic Carbene Boranes and Synthesis of Borane-Containing Liquid Crystals. <i>Organic Letters</i> , 2020 , 22, 2054-2059	6.2	8
106	Facile and Versatile Synthesis of End-Functionalized Poly(phenylacetylene)s: A Multicomponent Catalytic System for Well-Controlled Living Polymerization of Phenylacetylenes. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 8670-8680	16.4	12
105	Disclosing chirality in consecutive supramolecular polymerizations: chiral induction by light in N-annulated perylenetetracarboxamides. <i>Chemical Communications</i> , 2020 , 56, 2244-2247	5.8	12
104	Chiral/Achiral Copolymers of Biphenylacetylenes Bearing Various Substituents: Chiral Amplification through Copolymerization, Followed by Enhancement/Inversion and Memory of the Macromolecular Helicity. <i>Macromolecules</i> , 2020 , 53, 973-981	5.5	15
103	The Thermal Rearrangement of an NHC-Ligated 3-Benzoborepin to an NHC-Boranorcaradiene. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 903-909	16.4	8
102	The Thermal Rearrangement of an NHC-Ligated 3-Benzoborepin to an NHC-Boranorcaradiene. <i>Angewandte Chemie</i> , 2020 , 132, 913-919	3.6	5
101	Catalytic one-handed helix-induction and memory of amphiphilic poly(biphenylacetylene)s in water. <i>Giant</i> , 2020 , 2, 100016	5.6	12
100	Dithiocarbamate-modified cellulose-based sorbents with high storage stability for selective removal of arsenite and hazardous heavy metals.. <i>RSC Advances</i> , 2020 , 10, 30238-30244	3.7	3
99	Revisiting the Polymerization of Diphenylacetylenes with Tungsten(VI) Chloride and Tetraphenyltin: An Alternative Mechanism by a Metathesis Catalytic System. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 14772-14780	16.4	8
98	Macromolecular helicity control of poly(phenyl isocyanate)s with a single stimuli-responsive chiral switch. <i>Chemical Communications</i> , 2019 , 55, 7906-7909	5.8	13
97	Three-State Switchable Chiral Stationary Phase Based on Helicity Control of an Optically Active Poly(phenylacetylene) Derivative by Using Metal Cations in the Solid State. <i>Journal of the American Chemical Society</i> , 2019 , 141, 8592-8598	16.4	47
96	Unexpectedly Strong Chiral Amplification of Chiral/Achiral and Chiral/Chiral Copolymers of Biphenylacetylenes and Further Enhancement/Inversion and Memory of the Macromolecular Helicity. <i>Journal of the American Chemical Society</i> , 2019 , 141, 7605-7614	16.4	57

95	Esters as Radical Acceptors: π NHC-Borylalkenyl Radicals Induce Lactonization by C-C Bond Formation/Cleavage on Esters. <i>Angewandte Chemie - International Edition</i> , 2019 , 58, 6357-6361	16.4	28
94	Esters as Radical Acceptors: π NHC-Borylalkenyl Radicals Induce Lactonization by C \equiv C Bond Formation/Cleavage on Esters. <i>Angewandte Chemie</i> , 2019 , 131, 6423-6427	3.6	13
93	Dithiocarbamate-modified cellulose resins: A novel adsorbent for selective removal of arsenite from aqueous media. <i>Journal of Hazardous Materials</i> , 2019 , 380, 120816	12.8	17
92	Helicity Induction and Its Static Memory of Poly(biphenylacetylene)s Bearing Pyridine N-Oxide Groups and Their Use as Asymmetric Organocatalysts. <i>Journal of Polymer Science Part A</i> , 2019 , 57, 2481-2490	2.5	12
91	Helicity induction and memory effect in poly(biphenylacetylene)s bearing various functional groups and their use as switchable chiral stationary phases for HPLC. <i>Polymer Chemistry</i> , 2019 , 10, 6260-6268	4.9	22
90	Circularly Polarized Luminescent Triptycene-Based Polymers. <i>ACS Macro Letters</i> , 2018 , 7, 364-369	6.6	39
89	Direct Detection of Hardly Detectable Hidden Chirality of Hydrocarbons and Deuterated Isotopomers by a Helical Polyacetylene through Chiral Amplification and Memory. <i>Journal of the American Chemical Society</i> , 2018 , 140, 3270-3276	16.4	70
88	Radical trans-Hydroboration of Alkynes with N-Heterocyclic Carbene Boranes. <i>Angewandte Chemie</i> , 2018 , 130, 9629-9634	3.6	20
87	Optically active distorted cyclic triptycenes: chiral stationary phases for HPLC.. <i>RSC Advances</i> , 2018 , 8, 20483-20487	3.7	10
86	Radical trans-Hydroboration of Alkynes with N-Heterocyclic Carbene Boranes. <i>Angewandte Chemie - International Edition</i> , 2018 , 57, 9485-9490	16.4	55
85	A mechanistic insight into the organocatalytic properties of imidazolium-based ionic liquids and a positive co-solvent effect on cellulose modification reactions in an ionic liquid. <i>RSC Advances</i> , 2017 , 7, 9423-9430	3.7	31
84	Chromatographic enantioseparation by poly(biphenylacetylene) derivatives with memory of both axial chirality and macromolecular helicity. <i>Chirality</i> , 2017 , 29, 120-129	2.1	16
83	Cellulose derivatives bearing pyrene-based π conjugated pendants with circularly polarized luminescence in molecularly dispersed state. <i>Polymer</i> , 2017 , 117, 220-224	3.9	19
82	Static Memory of Enantiomeric Helices Induced in a Poly(biphenylacetylene) by a Single Enantiomer Assisted by Temperature- and Solvent-Driven Helix Inversion. <i>Macromolecules</i> , 2017 , 50, 7801-7806	5.5	13
81	Chiral triptycene-pyrene π conjugated chromophores with circularly polarized luminescence. <i>Organic and Biomolecular Chemistry</i> , 2017 , 15, 8440-8447	3.9	24
80	Helical Polyacetylenes Induced via Noncovalent Chiral Interactions and Their Applications as Chiral Materials. <i>Topics in Current Chemistry</i> , 2017 , 375, 72	7.2	56
79	Chiral stationary phases consisting of π conjugated polymers bearing glucose-linked biphenyl units: reversible switching of resolution abilities based on a coil-to-helix transition. <i>Polymer Chemistry</i> , 2017 , 8, 4190-4198	4.9	18
78	Helical Polyacetylenes Induced via Noncovalent Chiral Interactions and Their Applications as Chiral Materials. <i>Topics in Current Chemistry Collections</i> , 2017 , 1-31	1.8	2

77	Supramolecular Helical Systems: Helical Assemblies of Small Molecules, Foldamers, and Polymers with Chiral Amplification and Their Functions. <i>Chemical Reviews</i> , 2016 , 116, 13752-13990	68.1	1029
76	Chiral Recognition Ability of an Optically Active Poly(diphenylacetylene) as a Chiral Stationary Phase for HPLC. <i>Chemistry Letters</i> , 2016 , 45, 1063-1065	1.7	31
75	Synthesis and chiroptical properties of a π -conjugated polymer containing glucose-linked biphenyl units in the main chain capable of folding into a helical conformation. <i>Polymer Chemistry</i> , 2016 , 7, 7522-7529	4.9	23
74	Synthesis of Optically Active Poly(diphenylacetylene)s Using Polymer Reactions and an Evaluation of Their Chiral Recognition Abilities as Chiral Stationary Phases for HPLC. <i>Molecules</i> , 2016 , 21,	4.8	27
73	Synthesis of Thieno[3,4-b]thiophene-Based Donor Molecules with Phenyl Ester Pendants for Organic Solar Cells: Control of Photovoltaic Properties via Single Substituent Replacement. <i>ChemistrySelect</i> , 2016 , 1, 703-709	1.8	8
72	Dual Memory of Enantiomeric Helices in Poly(phenylacetylene)s Induced by a Single Enantiomer through Helix Inversion and Dual Storage of the Enantiomeric Helicity Memories. <i>Macromolecules</i> , 2015 , 48, 4281-4293	5.5	40
71	Influence of 4-fluorophenyl pendants in thieno[3,4-b]thiophene-benzo[1,2-b:4,5-b']dithiophene-based polymers on the performance of photovoltaics. <i>Journal of Polymer Science Part A</i> , 2015 , 53, 1586-1593	2.5	2
70	Macromolecular Helicity Induction and Memory in a Poly(biphenylacetylene) Bearing an Ester Group and Its Application to a Chiral Stationary Phase for High-performance Liquid Chromatography. <i>Chemistry Letters</i> , 2015 , 44, 946-948	1.7	15
69	Efficient and rapid direct transesterification reactions of cellulose with isopropenyl acetate in ionic liquids. <i>RSC Advances</i> , 2015 , 5, 72071-72074	3.7	46
68	Switchable enantioseparation based on macromolecular memory of a helical polyacetylene in the solid state. <i>Nature Chemistry</i> , 2014 , 6, 429-34	17.6	259
67	Chiral Amplification in Polymer Brushes Consisting of Dynamic Helical Polymer Chains through the Long-Range Communication of Stereochemical Information. <i>Macromolecules</i> , 2014 , 47, 6540-6546	5.5	42
66	Helical polymer brushes with a preferred-handed helix-sense triggered by a terminal optically active group in the pendant. <i>Chemical Communications</i> , 2012 , 48, 3342-4	5.8	53
65	Amplification of macromolecular helicity of dynamic helical poly(phenylacetylene)s bearing non-racemic alanine pendants in dilute solution, liquid crystal and two-dimensional crystal. <i>Polymer Journal</i> , 2012 , 44, 42-50	2.7	23
64	Effect of Polyelectrolyte Function on Helical Structures of Optically Active Poly(phenylacetylene) Derivatives Bearing Basic or Acidic Functional Pendant Groups. <i>Macromolecules</i> , 2011 , 44, 8343-8349	5.5	21
63	Synthesis of polysaccharide derivatives bearing pyridine N-oxide groups and their use as asymmetric organocatalysts. <i>Reactive and Functional Polymers</i> , 2011 , 71, 1055-1058	4.6	20
62	Hierarchical amplification of macromolecular helicity of dynamic helical poly(phenylacetylene)s composed of chiral and achiral phenylacetylenes in dilute solution, liquid crystal, and two-dimensional crystal. <i>Journal of the American Chemical Society</i> , 2011 , 133, 108-14	16.4	57
61	Enantioselective Adsorption of Chiral Amines on an Induced Helical Poly(bis(4-carboxyphenoxy)phosphazene): Chiral Filter Effect. <i>Macromolecules</i> , 2011 , 44, 2457-2464	5.5	21
60	Synthesis of functional poly(phenyl isocyanide)s with macromolecular helicity memory and their use as asymmetric organocatalysts. <i>Chirality</i> , 2009 , 21, 44-50	2.1	70

59	Mechanism of helix induction in poly(4-carboxyphenyl isocyanide) with chiral amines and memory of the macromolecular helicity and its helical structures. <i>Journal of the American Chemical Society</i> , 2009 , 131, 10719-32	16.4	91
58	Helical polymers: synthesis, structures, and functions. <i>Chemical Reviews</i> , 2009 , 109, 6102-211	68.1	1277
57	Chirality-Responsive Helical Polymers. <i>Macromolecules</i> , 2008 , 41, 3-12	5.5	380
56	Single- and double-stranded helical polymers: synthesis, structures, and functions. <i>Accounts of Chemical Research</i> , 2008 , 41, 1166-80	24.3	400
55	Chemical Modification of a Luminescent Poly(phenylenevinylene)-Amylose Composite. <i>Macromolecules</i> , 2008 , 41, 5065-5069	5.5	7
54	Helicity induction and memory of the macromolecular helicity in a polyacetylene bearing a biphenyl pendant. <i>Chemistry - an Asian Journal</i> , 2008 , 3, 614-624	4.5	18
53	Application of Soluble Poly(phenylenevinylene) Wrapped in Amylose to Organic Light-Emitting Diodes. <i>Molecular Crystals and Liquid Crystals</i> , 2007 , 471, 29-38	0.5	6
52	Synthesis of Optically Active Helical Poly(phenylacetylene)s Bearing Oligopeptide Pendants and Their Use as Polymeric Organocatalysts for Asymmetric Epoxidation. <i>Macromolecules</i> , 2007 , 40, 6783-6785	5.5	58
51	Syntheses and Chiroptical Properties of Optically Active Helical Poly(phenylacetylene)s Bearing [60]Fullerene Pendants. <i>Macromolecules</i> , 2007 , 40, 9244-9251	5.5	29
50	Temperature-induced chiroptical changes in a helical poly(phenylacetylene) bearing N,N-diisopropylaminomethyl groups with chiral acids in water. <i>Chemistry - an Asian Journal</i> , 2007 , 2, 1314-21	4.5	23
49	A Luminescent Poly(phenylenevinylene)-Amylose Composite with Supramolecular Liquid Crystallinity. <i>Angewandte Chemie</i> , 2006 , 118, 6641-6645	3.6	11
48	Helicity Induction in a Poly(4-carboxyphenyl isocyanide) with Chiral Amines and Memory of the Macromolecular Helicity in Aqueous Solution. <i>Macromolecules</i> , 2006 , 39, 6003-6008	5.5	28
47	Hierarchical Amplification of Macromolecular Helicity in a Lyotropic Liquid Crystalline Charged Poly(phenylacetylene) by Nonracemic Dopants in Water and Its Helical Structure. <i>Macromolecules</i> , 2006 , 39, 5371-5380	5.5	65
46	Switching of macromolecular helicity of optically active poly(phenylacetylene)s bearing cyclodextrin pendants induced by various external stimuli. <i>Journal of the American Chemical Society</i> , 2006 , 128, 7639-50	16.4	166
45	Helicity Induction on a Poly (phenylacetylene) Bearing Carboxy Groups at the Meta-Position. <i>Kobunshi Ronbunshu</i> , 2006 , 63, 325-330	0	4
44	Chirality sensing of various biomolecules with helical poly(phenylacetylene)s bearing acidic functional groups in water. <i>Journal of Polymer Science Part A</i> , 2006 , 44, 5039-5048	2.5	26
43	Helicity Induction on a Poly(phenylacetylene) Derivative Bearing a Sulfonic Acid Pendant with Chiral Amines and Memory of the Macromolecular Helicity in Dimethyl Sulfoxide. <i>Polymer Journal</i> , 2006 , 38, 912-919	2.7	22
42	Helicity Induction in Charged Poly(phenylacetylene)s Bearing Various Acidic Functional Groups in Water and Its Mechanism. <i>Macromolecules</i> , 2005 , 38, 8625-8633	5.5	51

41	Dual memory of enantiomeric helices in a polyacetylene induced by a single enantiomer. <i>Journal of the American Chemical Society</i> , 2005 , 127, 5018-9	16.4	106
40	Layer-by-layer assembly of charged poly(phenylacetylene)s with induced macromolecular helicity. <i>Chemical Communications</i> , 2005 , 4152-4	5.8	12
39	Helicity Induction and Chiral Amplification in a Poly(phenylacetylene) Bearing N,N-Diisopropylaminomethyl Groups with Chiral Acids in Water. <i>Macromolecules</i> , 2005 , 38, 5444-5451	5.5	42
38	Enantioselective esterification of prochiral phosphonate pendants of a polyphenylacetylene assisted by macromolecular helicity: storage of a dynamic macromolecular helicity memory. <i>Journal of the American Chemical Society</i> , 2005 , 127, 2960-5	16.4	91
37	Helix induction in an optically inactive poly[(4-carboxyphenyl)acetylene] film with chiral amines. <i>Mendeleev Communications</i> , 2004 , 14, 231-233	1.9	8
36	Convenient synthesis of fully and partially deuterated stereoregular poly(phenylacetylene)s bearing a carboxy pendant and helicity induction on the polymers with chiral amines and its memory. <i>Journal of Polymer Science Part A</i> , 2004 , 42, 4711-4722	2.5	11
35	Helicity induction on a poly(phenylacetylene) bearing a phosphonate residue by chiral dendrons. <i>Journal of Polymer Science Part A</i> , 2004 , 42, 4580-4586	2.5	39
34	Detection and amplification of chirality by helical polymers. <i>Chemistry - A European Journal</i> , 2004 , 10, 42-51	4.8	504
33	Poly(phenylacetylene)s bearing a peptide pendant: helical conformational changes of the polymer backbone stimulated by the pendant conformational change. <i>Chemistry - A European Journal</i> , 2004 , 10, 4000-10	4.8	90
32	A helical array of pendant fullerenes on a helical poly(phenylacetylene) induced by non-covalent chiral interactions. <i>Chemical Communications</i> , 2004 , 646-7	5.8	24
31	Macromolecular helicity induction on a poly(phenylacetylene) with C ₂ -symmetric chiral [60]fullerene-bisadducts. <i>Journal of the American Chemical Society</i> , 2004 , 126, 11711-7	16.4	80
30	Macromolecular helicity induction in a cationic polyacetylene assisted by an anionic polyisocyanide with helicity memory in water: replication of macromolecular helicity. <i>Journal of the American Chemical Society</i> , 2004 , 126, 15161-6	16.4	52
29	Mechanism of helix induction on a stereoregular Poly((4-carboxyphenyl)acetylene) with chiral amines and memory of the macromolecular helicity assisted by interaction with achiral amines. <i>Journal of the American Chemical Society</i> , 2004 , 126, 4329-42	16.4	146
28	An unprecedented memory of macromolecular helicity induced in an achiral polyisocyanide in water. <i>Journal of the American Chemical Society</i> , 2004 , 126, 732-3	16.4	111
27	Helicity Induction on Poly(phenylacetylene)s Bearing Phosphonic Acid Pendants with Chiral Amines and Memory of the Macromolecular Helicity Assisted by Interaction with Achiral Amines in Dimethyl Sulfoxide. <i>Macromolecules</i> , 2004 , 37, 5495-5503	5.5	60
26	Nonracemic dopant-mediated hierarchical amplification of macromolecular helicity in a charged polyacetylene leading to a cholesteric liquid crystal in water. <i>Journal of the American Chemical Society</i> , 2004 , 126, 16284-5	16.4	59
25	Macromolecular helicity inversion of poly(phenylacetylene) derivatives induced by various external stimuli. <i>Macromolecular Symposia</i> , 2003 , 201, 135-142	0.8	13
24	Cu(II)-assisted Helicity Induction on a Poly(phenylacetylene) Derivative Bearing an Achiral Glycine Residue with Amino Acids in Water. <i>Chemistry Letters</i> , 2003 , 32, 1086-1087	1.7	7

23	Solvent-induced switching of the macromolecular helicity of poly[(4-carboxyphenyl)acetylene] induced by a single chiral amino alcohol. <i>Journal of Polymer Science Part A</i> , 2003 , 41, 3625-3631	2.5	25
22	Helix-Sense Inversion of Poly(phenylacetylene) Derivatives Bearing an Optically Active Substituent Induced by External Chiral and Achiral Stimuli. <i>Macromolecules</i> , 2003 , 36, 1480-1486	5.5	88
21	A Helical Array of Pendant Fullerenes on an Optically Active Polyphenylacetylene. <i>Angewandte Chemie</i> , 2002 , 114, 3754-3756	3.6	12
20	Temperature dependence of helical structures of poly(phenylacetylene) derivatives bearing an optically active substituent. <i>Chemistry - A European Journal</i> , 2002 , 8, 5112-20	4.8	77
19	Helicity Induction on Optically Inactive Polyacetylenes and Polyphosphazenes. <i>ACS Symposium Series</i> , 2002 , 41-53	0.4	2
18	Helicity Induction on Macromolecules.. <i>Yuki Gosei Kagaku Kyokaiishi/Journal of Synthetic Organic Chemistry</i> , 2002 , 60, 878-890	0.2	10
17	Molecular Recognition of Nucleosides and Nucleotides Based on Circular Dichroism Induced by Helix Formation of Poly[(4-dihydroxyborophenyl)acetylene]. <i>Chemistry Letters</i> , 2001 , 30, 58-59	1.7	21
16	Helix formation of poly(phenylacetylene) derivatives bearing amino groups at the meta position induced by optically active carboxylic acids. <i>Journal of Polymer Science Part A</i> , 2001 , 39, 3180-3189	2.5	27
15	A helical polyelectrolyte induced by specific interactions with biomolecules in water. <i>Journal of the American Chemical Society</i> , 2001 , 123, 7441-2	16.4	112
14	Switching of a macromolecular helicity for visual distinction of molecular recognition events. <i>Journal of the American Chemical Society</i> , 2001 , 123, 8159-60	16.4	185
13	Stereospecific Polymerization of Propiolic Acid with Rhodium Complexes in the Presence of Bases and Helix Induction on the Polymer in Water. <i>Macromolecules</i> , 2001 , 34, 1160-1164	5.5	88
12	Synthesis and structure of poly(phenyl isocyanate)s bearing an optically active alkoxyl group. <i>Journal of Physical Organic Chemistry</i> , 2000 , 13, 361-367	2.1	14
11	Synthesis and Macromolecular Helicity Induction of a Stereoregular Polyacetylene Bearing a Carboxy Group with Natural Amino Acids in Water. <i>Macromolecules</i> , 2000 , 33, 4616-4618	5.5	85
10	Helicity Induction and Conformational Dynamics of Poly(bis(4-carboxyphenoxy)phosphazene) with Optically Active Amines. <i>Journal of the American Chemical Society</i> , 2000 , 122, 7813-7814	16.4	71
9	Memory of macromolecular helicity assisted by interaction with achiral small molecules. <i>Nature</i> , 1999 , 399, 449-451	50.4	668
8	Synthesis and Conformation of Optically Active Poly(phenyl isocyanate)s Bearing an ((S)- β Methylbenzyl)carbamoyl) Group. <i>Macromolecules</i> , 1998 , 31, 1046-1052	5.5	58
7	Unusual Conformational Change of Optically Active Poly(3-((S)-sec-butoxycarbonyl)phenyl isocyanate). <i>Macromolecules</i> , 1998 , 31, 5164-6	5.5	64
6	Helical Structure of Oligo- and Poly(m-substituted phenyl isocyanate)s Bearing an Optically Active End-Group. <i>Polymer Journal</i> , 1998 , 30, 100-105	2.7	36

5	Synthesis and Chiral Recognition of Helical Polymers. <i>Journal of Macromolecular Science - Pure and Applied Chemistry</i> , 1997 , 34, 1771-1783	2.2	21
4	Chiroptical Properties of Oligomers of m-Methylphenyl Isocyanate Bearing an Optically Active End-Group. <i>Polymer Journal</i> , 1995 , 27, 141-146	2.7	43
3	Helically Folding Polymers331-366		29
2	Dynamic Helical Structures: Detection and Amplification of Chirality47-88		205
1	Helical Polymers with Dynamic and Static Macromolecular Helicity Memory: The Power of Helicity Memory for Helical Polymer Synthesis and Applications. <i>Bulletin of the Chemical Society of Japan</i> ,	5.1	8