

AurÃ©lie Goineau

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4954273/publications.pdf>

Version: 2024-02-01

18
papers

806
citations

623734

14
h-index

839539

18
g-index

18
all docs

18
docs citations

18
times ranked

854
citing authors

#	ARTICLE	IF	CITATIONS
1	Insights into the abundance and diversity of abyssal megafauna in a polymetallic-nodule region in the eastern Clarion-Clipperton Zone. <i>Scientific Reports</i> , 2016, 6, 30492.	3.3	173
2	Characteristics of meiofauna in extreme marine ecosystems: a review. <i>Marine Biodiversity</i> , 2018, 48, 35-71.	1.0	153
3	Live (stained) benthic foraminifera from the Rhône prodelta (Gulf of Lion, NW Mediterranean): Environmental controls on a river-dominated shelf. <i>Journal of Sea Research</i> , 2011, 65, 58-75.	1.6	89
4	Giant protists (xenophyophores, Foraminifera) are exceptionally diverse in parts of the abyssal eastern Pacific licensed for polymetallic nodule exploration. <i>Biological Conservation</i> , 2017, 207, 106-116.	4.1	60
5	Temporal variability of live (stained) benthic foraminiferal faunas in a river-dominated shelf – Faunal response to rapid changes of the river influence (Rhône prodelta, NW Mediterranean). <i>Biogeosciences</i> , 2012, 9, 1367-1388.	3.3	39
6	Abyssal foraminifera attached to polymetallic nodules from the eastern Clarion Clipperton Fracture Zone: a preliminary description and comparison with North Atlantic dropstone assemblages. <i>Marine Biodiversity</i> , 2015, 45, 391-412.	1.0	39
7	Live–dead comparison of benthic foraminiferal faunas from the Rhône prodelta (Gulf of Lions, NW) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	1.2	38
8	Environment, ecology, and potential effectiveness of an area protected from deep-sea mining (Clarion) Tj ETQq0 0 0 rgBT /Overlock 10 T	3.2	36
9	Novel benthic foraminifera are abundant and diverse in an area of the abyssal equatorial Pacific licensed for polymetallic nodule exploration. <i>Scientific Reports</i> , 2017, 7, 45288.	3.3	35
10	Historical evolution and extreme climate events during the last 400years on the Rhone prodelta (NW) Tj ETQq0 0 0 rgBT /Overlock 10 T	2.1	30
11	Five new species and two new genera of xenophyophores (Foraminifera: Rhizaria) from part of the abyssal equatorial Pacific licensed for polymetallic nodule exploration. <i>Zoological Journal of the Linnean Society</i> , 2018, 183, 723-748.	2.3	20
12	Diversity and spatial patterns of foraminiferal assemblages in the eastern Clarion–Clipperton zone (abyssal eastern equatorial Pacific). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2019, 149, 103036.	1.4	18
13	The Biodiversity and Distribution of Abyssal Benthic Foraminifera and Their Possible Ecological Roles: A Synthesis Across the Clarion-Clipperton Zone. <i>Frontiers in Marine Science</i> , 2021, 8, .	2.5	18
14	The Contribution of Fine Sieve Fractions (63–150 µm) to Foraminiferal Abundance and Diversity in an Area of the Eastern Pacific Ocean Licensed for Polymetallic Nodule Exploration. <i>Frontiers in Marine Science</i> , 2019, 6, .	2.5	17
15	New species of the xenophyophore genus <i>Aschemonella</i> (Rhizaria: Foraminifera) from areas of the abyssal eastern Pacific licensed for polymetallic nodule exploration. <i>Zoological Journal of the Linnean Society</i> , 2018, 182, 479-499.	2.3	14
16	Radiolarian tests as microhabitats for novel benthic foraminifera: Observations from the abyssal eastern equatorial Pacific (Clarion–Clipperton Fracture Zone). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2015, 103, 73-85.	1.4	13
17	Xenophyophores (Rhizaria, Foraminifera) from the Eastern Clarion-Clipperton Zone (equatorial) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	1.5	9
18	Loricifera inhabiting spherical agglutinated structures in the abyssal eastern equatorial Pacific nodule fields. <i>Marine Biodiversity</i> , 2019, 49, 2455-2466.	1.0	5