

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4958197/publications.pdf>

Version: 2024-02-01

11  
papers

81  
citations

1937685  
4  
h-index

1588992  
8  
g-index

11  
all docs

11  
docs citations

11  
times ranked

41  
citing authors

#	ARTICLE	IF	CITATIONS
1	Patients with end-stage renal disease requiring hemodialysis benefit from percutaneous coronary intervention after non-ST-segment elevation myocardial infarction. <i>Internal and Emergency Medicine</i> , 2022, , 1.	2.0	2
2	Distal Transradial Access: a Safe and Feasible Approach for Coronary Catheterization in Cases of Total Radial Artery Occlusion. <i>Journal of Cardiovascular Translational Research</i> , 2022, , 1.	2.4	6
3	Dysbiosis of Gut Microbiota and Metabolite Phenylacetylglutamine in Coronary Artery Disease Patients With Stent Stenosis. <i>Frontiers in Cardiovascular Medicine</i> , 2022, 9, 832092.	2.4	19
4	A risk score model to predict in-hospital mortality of patients with end-stage renal disease and acute myocardial infarction. <i>Internal and Emergency Medicine</i> , 2021, 16, 905-912.	2.0	5
5	The potential regulatory role of hsa_circ_0004104 in the persistency of atrial fibrillation by promoting cardiac fibrosis via TGF- $\beta$ 2 pathway. <i>BMC Cardiovascular Disorders</i> , 2021, 21, 25.	1.7	17
6	Predictive value of CHA2DS2-VASc score combined with hs-CRP for new-onset atrial fibrillation in elderly patients with acute myocardial infarction. <i>BMC Cardiovascular Disorders</i> , 2021, 21, 175.	1.7	10
7	Plasma Galectin-3 is associated with progression from paroxysmal to persistent atrial fibrillation. <i>BMC Cardiovascular Disorders</i> , 2021, 21, 226.	1.7	6
8	Coronary catheterization via distal transradial access in patient with superficial radial artery: a case report. <i>BMC Cardiovascular Disorders</i> , 2021, 21, 616.	1.7	1
9	Blood group A: a risk factor for heart rupture after acute myocardial infarction. <i>BMC Cardiovascular Disorders</i> , 2020, 20, 471.	1.7	4
10	A novel risk score for predicting left atrial and left atrial appendage thrombogenic milieu in patients with non-valvular atrial fibrillation. <i>Thrombosis Research</i> , 2020, 192, 161-166.	1.7	6
11	ABO blood groups: A risk factor for left atrial and left atrial appendage thrombogenic milieu in patients with non-valvular atrial fibrillation. <i>Thrombosis Research</i> , 2017, 156, 45-50.	1.7	5