

# Delphine Muths

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4965119/publications.pdf>

Version: 2024-02-01

15  
papers

305  
citations

933447

10  
h-index

996975

15  
g-index

15  
all docs

15  
docs citations

15  
times ranked

403  
citing authors

#	ARTICLE	IF	CITATIONS
1	Large-scale genetic patchiness among NE Atlantic populations of the brittle star <i>Ophiothrix fragilis</i> . <i>Aquatic Biology</i> , 2009, 5, 117-132.	1.4	38
2	Restricted dispersal of the reef fish <i>Myripristis berndti</i> at the scale of the SW Indian Ocean. <i>Marine Ecology - Progress Series</i> , 2011, 443, 167-180.	1.9	38
3	Incomplete cryptic speciation between intertidal and subtidal morphs of <i>Acrocnida brachiata</i> (Echinodermata: Ophiuroidea) in the Northeast Atlantic. <i>Molecular Ecology</i> , 2006, 15, 3303-3318.	3.9	36
4	Genetic structure of the reef grouper <i>Epinephelus merra</i> in the West Indian Ocean appears congruent with biogeographic and oceanographic boundaries. <i>Marine Ecology</i> , 2015, 36, 447-461.	1.1	36
5	Comparative phylogeography of the western Indian Ocean reef fauna. <i>Acta Oecologica</i> , 2016, 72, 72-86.	1.1	35
6	Genetic population structure of the Swordfish ( <i>Xiphias gladius</i> ) in the southwest Indian Ocean: Sex-biased differentiation, congruency between markers and its incidence in a way of stock assessment. <i>Fisheries Research</i> , 2009, 97, 263-269.	1.7	26
7	Genetic connectivity of the reef fish <i>Lutjanus kasmira</i> at the scale of the western Indian Ocean. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 2012, 69, 842-853.	1.4	25
8	Multi-Genetic Marker Approach and Spatio-Temporal Analysis Suggest There Is a Single Panmictic Population of Swordfish <i>Xiphias gladius</i> in the Indian Ocean. <i>PLoS ONE</i> , 2013, 8, e63558.	2.5	19
9	Population structure enhances perspectives on regional management of the western Indian Ocean green turtle. <i>Conservation Genetics</i> , 2015, 16, 1069-1083.	1.5	18
10	Morphological diagnosis of the two genetic lineages of <i>Acrocnida brachiata</i> (Echinodermata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 United Kingdom, 2010, 90, 831-843.	0.8	13
11	Permanent Genetic Resources added to Molecular Ecology Resources Database 1 April 2011-31 May 2011. <i>Molecular Ecology Resources</i> , 2011, 11, 935-936.	4.8	8
12	Local scale connectivity in the cave-dwelling brooding fish <i>Apogon imberbis</i> . <i>Journal of Sea Research</i> , 2015, 95, 70-74.	1.6	6
13	Characterization of thirteen new polymorphic microsatellite markers from the honeycomb grouper <i>Epinephelus merra</i> . <i>Conservation Genetics Resources</i> , 2011, 3, 629-631.	0.8	4
14	Characterisation of 22 polymorphic microsatellite loci for the Broadbill Swordfish, <i>Xiphias gladius</i> . <i>Conservation Genetics Resources</i> , 2011, 3, 263-266.	0.8	2
15	Genetic differentiation in Eurasian Woodcock ( <i>Scolopax rusticola</i> ) from the Azores. <i>Ibis</i> , 2022, 164, 313-319.	1.9	1