Nadim W Alkharouf

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/496666/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Expanding the RNA virome of nematodes and other soil-inhabiting organisms. Virus Evolution, 2022, 8, veac019.	4.9	4
2	The conserved oligomeric Golgi (COG) complex, a window into plant-pathogen interactions. Journal of Plant Interactions, 2022, 17, 344-360.	2.1	1
3	The impact of pRAP vectors on plant genetic transformation and pathogenesis studies including an analysis of <i>BRI1-ASSOCIATED RECEPTOR KINASE 1 (BAK1)</i> Interactions, 2021, 16, 270-283.	2.1	5
4	Conserved oligomeric Golgi (COG) complex genes functioning in defense are expressed in root cells undergoing a defense response to a pathogenic infection and exhibit regulation my MAPKs. PLoS ONE, 2021, 16, e0256472.	2.5	5
5	Exocyst components promote an incompatible interaction between Glycine max (soybean) and Heterodera glycines (the soybean cyst nematode). Scientific Reports, 2020, 10, 15003.	3.3	18
6	Mitogen activated protein kinase (MAPK)-regulated genes with predicted signal peptides function in the Glycine max defense response to the root pathogenic nematode Heterodera glycines. PLoS ONE, 2020, 15, e0241678.	2.5	10
7	LUX ARRHYTHMO mediates crosstalk between the circadian clock and defense in Arabidopsis. Nature Communications, 2019, 10, 2543.	12.8	47
8	Transcriptome analysis identifies genes related to the waxy coating on blueberry fruit in two northern-adapted rabbiteye breeding populations. BMC Plant Biology, 2019, 19, 460.	3.6	22
9	MAPKDB: A MAP kinase database for signal transduction element identification. Bioinformation, 2019, 15, 338-341.	0.5	6
10	Compare and Contrast of Differential Gene Expression Software Packages of RNA-Seq. , 2018, , .		1
11	An eFP browser for visualizing strawberry fruit and flower transcriptomes. Horticulture Research, 2017, 4, 17029.	6.3	63
12	Analysis of the genome sequence of Phomopsis longicolla: a fungal pathogen causing Phomopsis seed decay in soybean. BMC Genomics, 2017, 18, 688.	2.8	40
13	Proteomic Investigation of <i>Rhizoctonia solani</i> AG 4 Identifies Secretome and Mycelial Proteins with Roles in Plant Cell Wall Degradation and Virulence. Journal of Agricultural and Food Chemistry, 2016, 64, 3101-3110.	5.2	18
14	Genome-wide functional annotation of Phomopsis longicolla isolate MSPL 10-6. Genomics Data, 2016, 8, 67-69.	1.3	2
15	Data mining of genomic data generated from soybean treated with different phytohormones. , 2016, , .		0
16	A searchable database for the genome of Phomopsis longicolla (isolate MSPL 10-6). Bioinformation, 2016, 12, 233-236.	0.5	0
17	Draft genome sequence of Phomopsis longicolla isolate MSPL 10-6. Genomics Data, 2015, 3, 55-56.	1.3	20
18	Re-annotation of the woodland strawberry (Fragaria vesca) genome. BMC Genomics, 2015, 16, 29.	2.8	60

NADIM W ALKHAROUF

#	Article	IF	CITATIONS
19	SoyProLow: A protein database enriched in low abundant soybean proteins. Bioinformation, 2014, 10, 599-601.	0.5	13
20	Floral Transcriptomes in Woodland Strawberry Uncover Developing Receptacle and Anther Gene Networks. Plant Physiology, 2014, 165, 1062-1075.	4.8	167
21	SCNProDB: A database for the identification of soybean cyst nematode proteins. Bioinformation, 2014, 10, 387-389.	0.5	4
22	Generating Genomic Tools for Blueberry Improvement. International Journal of Fruit Science, 2012, 12, 276-287.	2.4	19
23	Differences in gene expression amplitude overlie a conserved transcriptomic program occurring between the rapid and potent localized resistant reaction at the syncytium of the Glycine max genotype Peking (PI 548402) as compared to the prolonged and potent resistant reaction of PI 88788. Plant Molecular Biology, 2011, 75, 141-165.	3.9	48
24	Mapping cell fate decisions that occur during soybean defense responses. Plant Molecular Biology, 2011, 77, 513-528.	3.9	59
25	Syncytium gene expression in Glycine max[Pl 88788] roots undergoing a resistant reaction to the parasitic nematode Heterodera glycines. Plant Physiology and Biochemistry, 2010, 48, 176-193.	5.8	74
26	Microarray Detection Call Methodology as a Means to Identify and Compare Transcripts Expressed within Syncytial Cells from Soybean (<i>Glycine max</i>) Roots Undergoing Resistant and Susceptible Reactions to the Soybean Cyst Nematode (<i>Heterodera glycines</i>). Journal of Biomedicine and Biotechnology, 2010, 2010, 1-30.	3.0	37
27	A gene expression analysis of syncytia laser microdissected from the roots of the Glycine max (soybean) genotype PI 548402 (Peking) undergoing a resistant reaction after infection by Heterodera glycines (soybean cyst nematode). Plant Molecular Biology, 2009, 71, 525-567.	3.9	99
28	BBGD: an online database for blueberry genomic data. BMC Plant Biology, 2007, 7, 5.	3.6	17
29	Laser capture microdissection (LCM) and comparative microarray expression analysis of syncytial cells isolated from incompatible and compatible soybean (Glycine max) roots infected by the soybean cyst nematode (Heterodera glycines). Planta, 2007, 226, 1389-1409.	3.2	154
30	Laser Capture Microdissection (LCM) and Expression Analyses of Glycine max (Soybean) Syncytium Containing Root Regions Formed by the Plant Pathogen Heterodera glycines (Soybean Cyst Nematode). Plant Molecular Biology, 2005, 59, 965-979.	3.9	114
31	The heterologous expression of conserved Glycine max (soybean) mitogen activated protein kinase 3 (MAPK3) paralogs suppresses Meloidogyne incognita parasitism in Gossypium hirsutum (upland) Tj ETQq1 1 0	.784 3.1 4 rg	BT Øverlock