

Silvio Aime

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4968472/publications.pdf>

Version: 2024-02-01

737
papers

31,300
citations

6124

83
h-index

13635

134
g-index

839
all docs

839
docs citations

839
times ranked

20891
citing authors

#	ARTICLE	IF	CITATIONS
1	Prospects and limitations of paramagnetic chemical exchange saturation transfer agents serving as biological reporters in vivo. <i>NMR in Biomedicine</i> , 2023, 36, e4698.	1.6	10
2	GlucoCEST MRI for the Evaluation Response to Chemotherapeutic and Metabolic Treatments in a Murine Triple-Negative Breast Cancer: A Comparison with [18F]F-FDG-PET. <i>Molecular Imaging and Biology</i> , 2022, 24, 126-134.	1.3	9
3	Effect of the hydrogenation solvent in the PHIP-SAH hyperpolarization of [1-13C]pyruvate. <i>Catalysis Today</i> , 2022, 397-399, 94-102.	2.2	10
4	How the catalysis of the prototropic exchange affects the properties of lanthanide(III) complexes in their applications as MRI contrast agents. <i>Inorganica Chimica Acta</i> , 2022, 532, 120730.	1.2	6
5	What do we know about dynamic glucose-enhanced (DGE) MRI and how close is it to the clinics? Horizon 2020 GLINT consortium report. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2022, 35, 87-104.	1.1	7
6	XNAT-PIC: Extending XNAT to Preclinical Imaging Centers. <i>Journal of Digital Imaging</i> , 2022, 35, 860-875.	1.6	3
7	The interaction between iodinated X-ray contrast agents and macrocyclic GBCAs provides a signal enhancement in T ₁ -weighted MR images: Insights into the renal excretion pathways of GdHPDO3A and iodixanol in healthy mice. <i>Magnetic Resonance in Medicine</i> , 2022, 88, 357-364.	1.9	4
8	Highly Sensitive Off/On-EPR Probes to Monitor Enzymatic Activity. <i>Chemistry - A European Journal</i> , 2022, 28, .	1.7	3
9	New tools to investigate tumor metabolism by NMR/MRI. <i>Journal of Magnetic Resonance</i> , 2022, 338, 107198.	1.2	5
10	Effects of Cations on HPTS Fluorescence and Quantification of Free Gadolinium Ions in Solution; Assessment of Intracellular Release of Gd ³⁺ from Gd-Based MRI Contrast Agents. <i>Molecules</i> , 2022, 27, 2490.	1.7	1
11	Review and consensus recommendations on clinical APT-weighted imaging approaches at 3T: Application to brain tumors. <i>Magnetic Resonance in Medicine</i> , 2022, 88, 546-574.	1.9	79
12	Studies of the hydrophobic interaction between a pyrene-containing dye and a tetra-aza macrocyclic gadolinium complex. <i>Inorganic Chemistry Frontiers</i> , 2022, 9, 3494-3504.	3.0	1
13	[Gd(AAZTA)] ⁿ⁺ Derivatives with n-Alkyl Acid Side Chains Show Improved Properties for Their Application as MRI Contrast Agents**. <i>Chemistry - A European Journal</i> , 2021, 27, 1849-1859.	1.7	4
14	Supramolecular adducts between macrocyclic Gd(III) complexes and polyaromatic systems: a route to enhance the relaxivity through the formation of hydrophobic interactions. <i>Chemical Science</i> , 2021, 12, 1368-1377.	3.7	7
15	A Novel Class of 1 H-MRI Contrast Agents Based on the Relaxation Enhancement Induced on Water Protons by 14 N-Containing Imidazole Moieties. <i>Angewandte Chemie - International Edition</i> , 2021, 60, 4208-4214.	7.2	8
16	A Novel Class of 1 H-MRI Contrast Agents Based on the Relaxation Enhancement Induced on Water Protons by 14 N-Containing Imidazole Moieties. <i>Angewandte Chemie</i> , 2021, 133, 4254-4260.	1.6	1
17	Singulett-Kontrast-Magnetresonanztomographie: Freisetzung der Hyperpolarisation durch den Metabolismus**. <i>Angewandte Chemie</i> , 2021, 133, 6866-6873.	1.6	3
18	Singlet-Contrast Magnetic Resonance Imaging: Unlocking Hyperpolarization with Metabolism**. <i>Angewandte Chemie - International Edition</i> , 2021, 60, 6791-6798.	7.2	28

#	ARTICLE	IF	CITATIONS
19	LipHosomes: Reporters for Ligand/Anti-Ligand Assays Based On pH Readout. <i>Analysis & Sensing</i> , 2021, 1, 48-53.	1.1	1
20	Chemistry of Molecular Imaging: An Overview. , 2021, , 423-443.		2
21	An albumin-binding Gd-HPDO3A contrast agent for improved intravascular retention. <i>Inorganic Chemistry Frontiers</i> , 2021, 8, 4014-4025.	3.0	4
22	Enhanced relaxivity of Gd ^{III} -complexes with HP-DO3A-like ligands upon the activation of the intramolecular catalysis of the prototropic exchange. <i>Inorganic Chemistry Frontiers</i> , 2021, 8, 1500-1510.	3.0	9
23	Hydrogenative-PHIP polarized metabolites for biological studies. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2021, 34, 25-47.	1.1	34
24	Rapid hyperpolarization and purification of the metabolite fumarate in aqueous solution. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	54
25	Targeting Chronic Myeloid Leukemia Stem/Progenitor Cells Using Venetoclax-Loaded Immunoliposome. <i>Cancers</i> , 2021, 13, 1311.	1.7	21
26	A global view of standards for open image data formats and repositories. <i>Nature Methods</i> , 2021, 18, 1440-1446.	9.0	36
27	ParaHydrogen Polarized Ethyl ¹³ C pyruvate in Water, a Key Substrate for Fostering the PHIP-5AH Approach to Metabolic Imaging. <i>ChemPhysChem</i> , 2021, 22, 1042-1048.	1.0	10
28	In vitro and in vivo comparison of MRI chemical exchange saturation transfer (CEST) properties between native glucose and 3-Methyl-D-glucose in a murine tumor model. <i>NMR in Biomedicine</i> , 2021, 34, 1.6 e4602.	1.6	9
29	Fe(deferasirox) ₂ : An Iron(III)-Based Magnetic Resonance Imaging Contrast Agent Endowed with Remarkable Molecular and Functional Characteristics. <i>Journal of the American Chemical Society</i> , 2021, 143, 14178-14188.	6.6	22
30	Low-Field NMR Relaxometry for Intraoperative Tumour Margin Assessment in Breast-Conserving Surgery. <i>Cancers</i> , 2021, 13, 4141.	1.7	3
31	Analysis of the Gadolinium retention in the Experimental Autoimmune Encephalomyelitis (EAE) murine model of Multiple Sclerosis. <i>Journal of Trace Elements in Medicine and Biology</i> , 2021, 68, 126831.	1.5	2
32	Monitoring tissue implants by field-cycling 1H-MRI via the detection of changes in the 14N-quadrupolar-peak from imidazole moieties incorporated in a smart scaffold material. <i>Journal of Materials Chemistry B</i> , 2021, 9, 4863-4872.	2.9	5
33	H-Bonding and intramolecular catalysis of proton exchange affect the CEST properties of Eu ^{III} complexes with HP-DO3A-like ligands. <i>Chemical Communications</i> , 2021, 57, 3287-3290.	2.2	3
34	Detection of U-87 Tumor Cells by RGD-Functionalized/Gd-Containing Giant Unilamellar Vesicles in Magnetization Transfer Contrast Magnetic Resonance Images. <i>Investigative Radiology</i> , 2021, 56, 301-312.	3.5	8
35	Imaging of Dysfunctional Elastogenesis in Atherosclerosis Using an Improved Gadolinium-Based Tetrameric MRI Probe Targeted to Tropoelastin. <i>Journal of Medicinal Chemistry</i> , 2021, 64, 15250-15261.	2.9	2
36	Intracellular Water Lifetime as a Tumor Biomarker to Monitor Doxorubicin Treatment via FFC-Relaxometry in a Breast Cancer Model. <i>Frontiers in Oncology</i> , 2021, 11, 778823.	1.3	5

#	ARTICLE	IF	CITATIONS
37	Novel Gastrin-Releasing Peptide Receptor Targeted Near-Infrared Fluorescence Dye for Image-Guided Surgery of Prostate Cancer. <i>Molecular Imaging and Biology</i> , 2020, 22, 85-93.	1.3	16
38	Gadolinium Retention in Erythrocytes and Leukocytes From Human and Murine Blood Upon Treatment With Gadolinium-Based Contrast Agents for Magnetic Resonance Imaging. <i>Investigative Radiology</i> , 2020, 55, 30-37.	3.5	13
39	Contaminations in (meta)genome data: An open issue for the scientific community. <i>IUBMB Life</i> , 2020, 72, 698-705.	1.5	13
40	Insights into Interfacial Water Structuring at the Nafion Surface by T_1 -Weighted Magnetic Resonance Imaging. <i>Langmuir</i> , 2020, 36, 540-545.	1.6	5
41	L-ferritin: A theranostic agent of natural origin for MRI visualization and treatment of breast cancer. <i>Journal of Controlled Release</i> , 2020, 319, 300-310.	4.8	19
42	Multilamellar LipoCEST Agents Obtained from Osmotic Shrinkage of Paramagnetically Loaded Giant Unilamellar Vesicles (GUVs). <i>Angewandte Chemie - International Edition</i> , 2020, 59, 2279-2283.	7.2	5
43	Liposome-Based Bioassays. <i>Biology</i> , 2020, 9, 202.	1.3	15
44	Rare earth elements (REE) in biology and medicine. <i>Rendiconti Lincei</i> , 2020, 31, 821-833.	1.0	33
45	Manganese-enhanced MRI (MEMRI) in breast and prostate cancers: Preliminary results exploring the potential role of calcium receptors. <i>PLoS ONE</i> , 2020, 15, e0224414.	1.1	4
46	An Improved Biocompatible Probe for Photoacoustic Tumor Imaging Based on the Conjugation of Melanin to Bovine Serum Albumin. <i>Applied Sciences (Switzerland)</i> , 2020, 10, 8313.	1.3	0
47	In-vitro NMR Studies of Prostate Tumor Cell Metabolism by Means of Hyperpolarized $[1-^{13}C]$ Pyruvate Obtained Using the PHIP-SAH Method. <i>Frontiers in Oncology</i> , 2020, 10, 497.	1.3	25
48	Towards an Improved Design of MRI Contrast Agents: Synthesis and Relaxometric Characterisation of Gd ϵ HPDO3A Analogues. <i>Chemistry - A European Journal</i> , 2020, 26, 6056-6063.	1.7	6
49	Relaxometric studies of erythrocyte suspensions infected by <i>Plasmodium falciparum</i> : a tool for staging infection and testing anti-malarial drugs. <i>Magnetic Resonance in Medicine</i> , 2020, 84, 3366-3378.	1.9	13
50	Relaxometric Studies of Gd-Chelate Conjugated on the Surface of Differently Shaped Gold Nanoparticles. <i>Nanomaterials</i> , 2020, 10, 1115.	1.9	4
51	An innovative approach for the synthesis of dual modality peptide imaging probes based on the native chemical ligation approach. <i>Chemical Communications</i> , 2020, 56, 3500-3503.	2.2	8
52	Nanocarriers as Magic Bullets in the Treatment of Leukemia. <i>Nanomaterials</i> , 2020, 10, 276.	1.9	38
53	In vivo assessment of tumour associated macrophages in murine melanoma obtained by low-field relaxometry in the presence of iron oxide particles. <i>Biomaterials</i> , 2020, 236, 119805.	5.7	16
54	A Simple and Fast Assay Based on Carboxyfluorescein-Loaded Liposome for Quantitative DNA Detection. <i>ACS Omega</i> , 2020, 5, 1764-1772.	1.6	7

#	ARTICLE	IF	CITATIONS
55	Multilamellar LipoCEST Agents Obtained from Osmotic Shrinkage of Paramagnetically Loaded Giant Unilamellar Vesicles (GUVs). <i>Angewandte Chemie</i> , 2020, 132, 2299-2303.	1.6	2
56	Quantification of hydroxyl exchange of D ² O-Glucose at physiological conditions for optimization of glucoCEST MRI at 3, 7 and 9.4 Tesla. <i>NMR in Biomedicine</i> , 2019, 32, e4113.	1.6	49
57	Synthesis and Relaxometric Characterization of New Poly[<i>N</i> , <i>N</i> '-bis(3-aminopropyl)glycine] (PAPGly) Dendrons Gd ³⁺ -Based Contrast Agents and Their <i>In Vivo</i> Study by Using the Dynamic Contrast-Enhanced MRI Technique at Low Field (1 T). <i>Chemistry and Biodiversity</i> , 2019, 16, e1900322.	1.0	3
58	Absence of dentate nucleus resting-state functional connectivity changes in nonneurological patients with gadolinium-related hyperintensity on T ₁ -weighted images. <i>Journal of Magnetic Resonance Imaging</i> , 2019, 50, 445-455.	1.9	23
59	Exploring the tumour extracellular matrix by <i>in vivo</i> Fast Field Cycling relaxometry after the administration of a Gadolinium-based MRI contrast agent. <i>Magnetic Resonance in Chemistry</i> , 2019, 57, 845-851.	1.1	7
60	Magnetic Resonance Imaging Reveals Distinct Roles for Tissue Transglutaminase and Factor XIII in Maternal Angiogenesis During Early Mouse Pregnancy. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2019, 39, 1602-1613.	1.1	4
61	Polydopamine-decorated tobacco mosaic virus for photoacoustic/magnetic resonance bimodal imaging and photothermal cancer therapy. <i>Nanoscale</i> , 2019, 11, 9760-9768.	2.8	37
62	A Gadolinium(III) Zeolite-like Metal-Organic-Framework-Based Magnetic Resonance Thermometer. <i>Chem</i> , 2019, 5, 1609-1618.	5.8	38
63	Use of FCC-NMRD relaxometry for early detection and characterization of <i>ex-vivo</i> murine breast cancer. <i>Scientific Reports</i> , 2019, 9, 4624.	1.6	8
64	Development and characterization of lanthanide-HPDO3A-C16-based micelles as CEST-MRI contrast agents. <i>Dalton Transactions</i> , 2019, 48, 5343-5351.	1.6	6
65	Differences in Molecular Structure Markedly Affect GBCA Elimination Behavior. <i>Radiology</i> , 2019, 291, 267-268.	3.6	8
66	Characterization of a Manganese-Containing Nanoparticle as an MRI Contrast Agent. <i>European Journal of Inorganic Chemistry</i> , 2019, 2019, 1759-1766.	1.0	6
67	Real-Time Nuclear Magnetic Resonance Detection of Fumarase Activity Using Parahydrogen-Hyperpolarized [¹³ C]Fumarate. <i>Journal of the American Chemical Society</i> , 2019, 141, 20209-20214.	6.6	50
68	Characterisation of magnetic resonance imaging (MRI) contrast agents using NMR relaxometry. <i>Molecular Physics</i> , 2019, 117, 898-909.	0.8	50
69	Modifying LnHPDO3A Chelates for Improved T ₁ and CEST MRI Applications. <i>Chemistry - A European Journal</i> , 2019, 25, 4184-4193.	1.7	8
70	Gadolinium presence, MRI hyperintensities, and glucose uptake in the hypoperfused rat brain after repeated administrations of gadodiamide. <i>Neuroradiology</i> , 2019, 61, 163-173.	1.1	14
71	Chemical Insights into the Issues of Gd Retention in the Brain and Other Tissues Upon the Administration of Gd-Containing MRI Contrast Agents. <i>European Journal of Inorganic Chemistry</i> , 2019, 2019, 137-151.	1.0	32
72	Emerging Technologies to Image Tissue Metabolism. <i>Cell Metabolism</i> , 2019, 29, 518-538.	7.2	47

#	ARTICLE	IF	CITATIONS
73	An efficient MRI agent targeting extracellular markers in prostate adenocarcinoma. <i>Magnetic Resonance in Medicine</i> , 2019, 81, 1935-1946.	1.9	6
74	Relaxometric investigations addressing the determination of intracellular water lifetime: a novel tumour biomarker of general applicability. <i>Molecular Physics</i> , 2019, 117, 968-974.	0.8	12
75	Metabolic Studies of Tumor Cells Using [¹³ C] Pyruvate Hyperpolarized by Means of PHIP Side Arm Hydrogenation. <i>ChemPhysChem</i> , 2019, 20, 318-325.	1.0	49
76	Tumor Targeting via Sialic Acid: [68Ga]DOTA-en-pba as a New Tool for Molecular Imaging of Cancer with PET. <i>Molecular Imaging and Biology</i> , 2018, 20, 798-807.	1.3	10
77	Synthesis of High Relaxivity Gadolinium AAZTA Tetramers as Building Blocks for Bioconjugation. <i>Bioconjugate Chemistry</i> , 2018, 29, 1428-1437.	1.8	18
78	CEST MRI studies of cells loaded with lanthanide shift reagents. <i>Magnetic Resonance in Medicine</i> , 2018, 80, 1626-1637.	1.9	15
79	Gd accumulation in tissues of healthy mice upon repeated administrations of Gadodiamide and Gadoteridol. <i>Journal of Trace Elements in Medicine and Biology</i> , 2018, 48, 239-245.	1.5	23
80	Exploiting the Proton Exchange as an Additional Route to Enhance the Relaxivity of Paramagnetic MRI Contrast Agents. <i>Inorganic Chemistry</i> , 2018, 57, 5567-5574.	1.9	23
81	Evidence for the Role of Intracellular Water Lifetime as a Tumour Biomarker Obtained by In Vivo Field Cycling Relaxometry. <i>Angewandte Chemie - International Edition</i> , 2018, 57, 7468-7472.	7.2	44
82	Studies to enhance the hyperpolarization level in PHIP-SAH-produced C13-pyruvate. <i>Journal of Magnetic Resonance</i> , 2018, 289, 12-17.	1.2	70
83	[Yb(AAZTA)(H ₂ O)] ⁺ : an unconventional ParaCEST MRI probe. <i>Chemical Communications</i> , 2018, 54, 2004-2007.	2.2	11
84	Evidence for the Role of Intracellular Water Lifetime as a Tumour Biomarker Obtained by In Vivo Field Cycling Relaxometry. <i>Angewandte Chemie</i> , 2018, 130, 7590-7594.	1.6	4
85	A generalized ratiometric chemical exchange saturation transfer (CEST) MRI approach for mapping renal pH using iopamidol. <i>Magnetic Resonance in Medicine</i> , 2018, 79, 1553-1558.	1.9	57
86	Inner-sphere water and hydrogen bonds in lanthanide DOTAM complexes. A neutron diffraction study. <i>Inorganica Chimica Acta</i> , 2018, 470, 433-438.	1.2	6
87	No Lanthanide-based Catalysis in Eukaryotes. <i>IUBMB Life</i> , 2018, 71, 398-399.	1.5	4
88	Lanthanide-based catalysis in eukaryotes. <i>IUBMB Life</i> , 2018, 70, 1067-1075.	1.5	5
89	Cancer cell death induced by ferritins and the peculiar role of their labile iron pool. <i>Oncotarget</i> , 2018, 9, 27974-27984.	0.8	12
90	1H-NMR and MALDI-TOF MS as metabolomic quality control tests to classify platelet derived medium additives for GMP compliant cell expansion procedures. <i>PLoS ONE</i> , 2018, 13, e0203048.	1.1	2

#	ARTICLE	IF	CITATIONS
91	CESTâ€MRI for glioma pH quantification in mouse model: Validation by immunohistochemistry. NMR in Biomedicine, 2018, 31, e4005.	1.6	26
92	Complete on/off responsive ParaCEST MRI contrast agents for copper and zinc. Dalton Transactions, 2018, 47, 11346-11357.	1.6	19
93	Generation of multiparametric MRI maps by using Gd-labelled- RBCs reveals phenotypes and stages of murine prostate cancer. Scientific Reports, 2018, 8, 10567.	1.6	7
94	Efficient Route to Label Mesenchymal Stromal Cell-Derived Extracellular Vesicles. ACS Omega, 2018, 3, 8097-8103.	1.6	15
95	An innovative therapeutic approach for malignant mesothelioma treatment based on the use of Gd/boron multimodal probes for MRI guided BNCT. Journal of Controlled Release, 2018, 280, 31-38.	4.8	27
96	Exploring the intramolecular catalysis of the proton exchange process to modulate the relaxivity of Gd(ⁱⁱⁱ)-complexes of HP-DO3A-like ligands. Chemical Communications, 2018, 54, 10056-10059.	2.2	13
97	The ¹³ C hyperpolarized pyruvate generated by ParaHydrogen detects the response of the heart to altered metabolism in real time. Scientific Reports, 2018, 8, 8366.	1.6	119
98	Modulation of the Prototropic Exchange Rate in pHâ€Responsive Ybâ€HPDO3A Derivatives as ParaCEST Agents. ChemistrySelect, 2018, 3, 6035-6041.	0.7	11
99	The Issue of Gadolinium Retained in Tissues. Investigative Radiology, 2018, 53, 167-172.	3.5	44
100	AAZTA: An Ideal Chelating Agent for the Development of ⁴⁴ Sc PET Imaging Agents. Angewandte Chemie - International Edition, 2017, 56, 2118-2122.	7.2	53
101	AAZTA: An Ideal Chelating Agent for the Development of ⁴⁴ Sc PET Imaging Agents. Angewandte Chemie, 2017, 129, 2150-2154.	1.6	11
102	Theranostic Nanoparticles Loaded with Imaging Probes and Rubrocurcumin for Combined Cancer Therapy by Folate Receptor Targeting. ChemMedChem, 2017, 12, 502-509.	1.6	40
103	EXCI-CEST: Exploiting pharmaceutical excipients as MRI-CEST contrast agents for tumor imaging. International Journal of Pharmaceutics, 2017, 525, 275-281.	2.6	25
104	Unsaturated Longâ€Chain Fatty Acids Are Preferred Ferritin Ligands That Enhance Iron Biom mineralization. Chemistry - A European Journal, 2017, 23, 9879-9887.	1.7	10
105	Iron oxide/PLGA nanoparticles for magnetically controlled drug release. International Journal of Applied Electromagnetics and Mechanics, 2017, 53, S53-S60.	0.3	7
106	Noninvasive evaluation of renal pH homeostasis after ischemia reperfusion injury by CESTâ€MRI. NMR in Biomedicine, 2017, 30, e3720.	1.6	41
107	MRI. , 2017, , 227-324.		2
108	Ferritin Decorated PLGA/Paclitaxel Loaded Nanoparticles Endowed with an Enhanced Toxicity Toward MCF-7 Breast Tumor Cells. Bioconjugate Chemistry, 2017, 28, 1283-1290.	1.8	31

#	ARTICLE	IF	CITATIONS
109	Assessing tumor vascularization as a potential biomarker of imatinib resistance in gastrointestinal stromal tumors by dynamic contrast-enhanced magnetic resonance imaging. <i>Gastric Cancer</i> , 2017, 20, 629-639.	2.7	22
110	Eight-Coordinate, Stable Fe(II) Complex as a Dual ¹⁹ F and CEST Contrast Agent for Ratiometric pH Imaging. <i>Inorganic Chemistry</i> , 2017, 56, 12206-12213.	1.9	41
111	Gadolinium Retention in the Rat Brain: Assessment of the Amounts of Insoluble Gadolinium-containing Species and Intact Gadolinium Complexes after Repeated Administration of Gadolinium-based Contrast Agents. <i>Radiology</i> , 2017, 285, 839-849.	3.6	92
112	Enzyme-Responsive LipoCEST Agents: Assessment of MMP-2 Activity by Measuring the Intra-liposomal Water ¹ H-NMR Shift. <i>Angewandte Chemie - International Edition</i> , 2017, 56, 12170-12173.	7.2	19
113	Enzyme-Responsive LipoCEST Agents: Assessment of MMP-2 Activity by Measuring the Intra-liposomal Water ¹ H-NMR Shift. <i>Angewandte Chemie</i> , 2017, 129, 12338-12341.	1.6	7
114	Nano-sized and other improved reporters for magnetic resonance imaging of angiogenesis. <i>Advanced Drug Delivery Reviews</i> , 2017, 119, 61-72.	6.6	28
115	Water Soluble Melanin Derivatives for Dynamic Contrast Enhanced Photoacoustic Imaging of Tumor Vasculature and Response to Antiangiogenic Therapy. <i>Advanced Healthcare Materials</i> , 2017, 6, 1600550.	3.9	31
116	Macrocyclic paramagnetic agents for MRI: Determinants of relaxivity and strategies for their improvement. <i>Magnetic Resonance in Medicine</i> , 2017, 78, 1523-1532.	1.9	21
117	¹³ C-MR Hyperpolarization of Lactate by Using ParaHydrogen and Metabolic Transformation in Vitro. <i>Chemistry - A European Journal</i> , 2017, 23, 1200-1204.	1.7	56
118	Melanin-Based Contrast Agents for Biomedical Optoacoustic Imaging and Theranostic Applications. <i>International Journal of Molecular Sciences</i> , 2017, 18, 1719.	1.8	43
119	In vivo evaluation of tumour acidosis for assessing the early metabolic response and onset of resistance to dichloroacetate by using magnetic resonance pH imaging. <i>International Journal of Oncology</i> , 2017, 51, 498-506.	1.4	57
120	Chemical Shift and Relaxation Reagents in NMR \hat{t} . , 2017, , 195-202.		0
121	Chapter 3. Chemical Exchange Saturation Transfer (CEST) Contrast Agents. <i>New Developments in NMR</i> , 2017, , 243-317.	0.1	7
122	Chapter 19 Iodinated Contrast Media as pH-Responsive CEST Agents. , 2017, , 447-466.		8
123	The RNA-binding protein ESRP1 promotes human colorectal cancer progression. <i>Oncotarget</i> , 2017, 8, 10007-10024.	0.8	57
124	Endogenous glutamine decrease is associated with pancreatic cancer progression. <i>Oncotarget</i> , 2017, 8, 95361-95376.	0.8	41
125	Chapter 5 Birth of CEST Agents in Torino. , 2017, , 47-54.		0
126	Chapter 14 Saturating Compartmentalized Water Protons: Liposome- and Cell-Based CEST Agents. , 2017, , 311-344.		1

#	ARTICLE	IF	CITATIONS
127	Assessing the transport rate of hyperpolarized pyruvate and lactate from the intra- to the extracellular space. <i>NMR in Biomedicine</i> , 2016, 29, 1022-1027.	1.6	17
128	LipoCEST and cellCEST imaging agents: opportunities and challenges. <i>Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology</i> , 2016, 8, 602-618.	3.3	40
129	In Vitro and In Vivo Assessment of Nonionic Iodinated Radiographic Molecules as Chemical Exchange Saturation Transfer Magnetic Resonance Imaging Tumor Perfusion Agents. <i>Investigative Radiology</i> , 2016, 51, 155-162.	3.5	41
130	<i>Contrast Media</i> , 2016, , 59-70.		1
131	The release of Doxorubicin from liposomes monitored by MRI and triggered by a combination of US stimuli led to a complete tumor regression in a breast cancer mouse model. <i>Journal of Controlled Release</i> , 2016, 230, 57-63.	4.8	39
132	High kinetic inertness of a bis-hydrated Gd-complex with a constrained AAZTA-like ligand. <i>Chemical Communications</i> , 2016, 52, 11235-11238.	2.2	29
133	Functional imaging of the angiogenic switch in a transgenic mouse model of human breast cancer by dynamic contrast enhanced magnetic resonance imaging. <i>International Journal of Cancer</i> , 2016, 139, 404-413.	2.3	9
134	Mn^{2+} MRI-CEST at 1T with large Mn^{2+} complexes: An efficient MRI pH reporter. <i>Magnetic Resonance in Medicine</i> , 2016, 75, 329-336.	1.9	19
135	<i>In Vivo</i> Imaging of Tumor Metabolism and Acidosis by Combining PET and MRI-CEST pH Imaging. <i>Cancer Research</i> , 2016, 76, 6463-6470.	0.4	134
136	Probing lactate secretion in tumours with hyperpolarised NMR. <i>NMR in Biomedicine</i> , 2016, 29, 1079-1087.	1.6	6
137	Paramagnetic Phospholipid-Based Micelles Targeting VCAM-1 Receptors for MRI Visualization of Inflammation. <i>Bioconjugate Chemistry</i> , 2016, 27, 1921-1930.	1.8	21
138	Diolefin Based Nanostructures as Targeted Theranostics. <i>Journal of Biomedical Nanotechnology</i> , 2016, 12, 1076-1088.	0.5	3
139	^{15}N -labeled hyperpolarized perfusion marker. <i>Magnetic Resonance in Medicine</i> , 2016, 76, 1900-1904.	1.9	28
140	A relaxometric method for the assessment of intestinal permeability based on the oral administration of gadolinium-based MRI contrast agents. <i>NMR in Biomedicine</i> , 2016, 29, 475-482.	1.6	1
141	Re-evaluation of the water exchange lifetime value across red blood cell membrane. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2016, 1858, 627-631.	1.4	33
142	A rigidified AAZTA-like ligand as efficient chelator for ^{68}Ga radiopharmaceuticals. <i>ChemistrySelect</i> , 2016, 1, 163-171.	0.7	14
143	Mesoporous silica nanoparticles functionalized with fluorescent and MRI reporters for the visualization of murine tumors overexpressing $\text{V}\beta_3$ receptors. <i>Nanoscale</i> , 2016, 8, 7094-7104.	2.8	26
144	Gd-AAZTA-MADEC, an improved blood pool agent for DCE-MRI studies on mice on 1T scanners. <i>Biomaterials</i> , 2016, 75, 47-57.	5.7	41

#	ARTICLE	IF	CITATIONS
145	L-Ferritin targets breast cancer stem cells and delivers therapeutic and imaging agents. <i>Oncotarget</i> , 2016, 7, 66713-66727.	0.8	54
146	Simultaneous MR imaging for tissue engineering in a rat model of stroke. <i>Scientific Reports</i> , 2015, 5, 14597.	1.6	26
147	Sensitive MRI detection of internalized T_1 contrast agents using magnetization transfer contrast. <i>NMR in Biomedicine</i> , 2015, 28, 1663-1670.	1.6	11
148	MEMRI and tumors: a method for the evaluation of the contribution of Mn(II) ions in the extracellular compartment. <i>NMR in Biomedicine</i> , 2015, 28, 1104-1110.	1.6	6
149	Glucan Particles Loaded with Fluorinated Emulsions: A Sensitivity Improvement for the Visualization of Phagocytic Cells by ^{19}F -MRI. <i>Current Molecular Imaging</i> , 2015, 4, 29-34.	0.7	3
150	MRI Contrast Agents for Pharmacological Research. <i>Frontiers in Pharmacology</i> , 2015, 6, 290.	1.6	13
151	The hydroboration reaction as a key for a straightforward synthesis of new MRI-NCT agents. <i>Organic and Biomolecular Chemistry</i> , 2015, 13, 3288-3297.	1.5	7
152	Evaluation of the dose enhancement of combined $^{10}B + ^{157}Gd$ neutron capture therapy (NCT). <i>Radiation Protection Dosimetry</i> , 2015, 166, 369-373.	0.4	10
153	ParaHydrogen Induced Polarization of ^{13}C carboxylate resonance in acetate and pyruvate. <i>Nature Communications</i> , 2015, 6, 5858.	5.8	206
154	The Role of Equilibrium and Kinetic Properties in the Dissociation of Gd[DTPA α -bis(methylamide)] (Omniscan) at near to Physiological Conditions. <i>Chemistry - A European Journal</i> , 2015, 21, 4789-4799.	1.7	40
155	Sonosensitive theranostic liposomes for preclinical in vivo MRI-guided visualization of doxorubicin release stimulated by pulsed low intensity non-focused ultrasound. <i>Journal of Controlled Release</i> , 2015, 202, 21-30.	4.8	52
156	Insights on the relaxation of liposomes encapsulating paramagnetic Ln α -based complexes. <i>Magnetic Resonance in Medicine</i> , 2015, 74, 468-473.	1.9	15
157	A theranostic approach based on the use of a dual boron/Gd agent to improve the efficacy of Boron Neutron Capture Therapy in the lung cancer treatment. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2015, 11, 741-750.	1.7	51
158	Targeting ferritin receptors for the selective delivery of imaging and therapeutic agents to breast cancer cells. <i>Nanoscale</i> , 2015, 7, 6527-6533.	2.8	67
159	Cluster analysis of quantitative parametric maps from DCE-MRI: application in evaluating heterogeneity of tumor response to antiangiogenic treatment. <i>Magnetic Resonance Imaging</i> , 2015, 33, 725-736.	1.0	34
160	An MRI Method To Map Tumor Hypoxia Using Red Blood Cells Loaded with a pO_2 -Responsive Gd-Agent. <i>ACS Nano</i> , 2015, 9, 8239-8248.	7.3	36
161	A novel patient-derived tumorgraft model with TRAF1-ALK anaplastic large-cell lymphoma translocation. <i>Leukemia</i> , 2015, 29, 1390-1401.	3.3	42
162	Cardio-Chemical Exchange Saturation Transfer Magnetic Resonance Imaging Reveals Molecular Signatures of Endogenous Fibrosis and Exogenous Contrast Media. <i>Circulation: Cardiovascular Imaging</i> , 2015, 8, .	1.3	27

#	ARTICLE	IF	CITATIONS
163	Optimizing the high-field relaxivity by self-assembling of macrocyclic Gd(III) complexes. Dalton Transactions, 2015, 44, 4910-4917.	1.6	9
164	Preparation and in vitro characterization of chitosan nanobubbles as theranostic agents. Colloids and Surfaces B: Biointerfaces, 2015, 129, 39-46.	2.5	62
165	In Vivo MR Imaging of Fibrin in a Neuroblastoma Tumor Model by Means of a Targeting Gd-Containing Peptide. Molecular Imaging and Biology, 2015, 17, 819-828.	1.3	10
166	Quantitative description of radiofrequency (RF) power-based ratiometric chemical exchange saturation transfer (CEST) pH imaging. NMR in Biomedicine, 2015, 28, 555-565.	1.6	53
167	Gd-loaded-RBCs for the assessment of tumor vascular volume by contrast-enhanced-MRI. Biomaterials, 2015, 58, 82-92.	5.7	29
168	Synthesis of phosphonic analogues of AAZTA=6-Amino-6-methylperhydro-1,4-diazepine-N,N ² ,N ³ ,N ³ -tetraacetic acid. and relaxometric evaluation of the corresponding Gd(III) complexes as potential MRI contrast agents. Tetrahedron Letters, 2015, 56, 1994-1997.	0.7	13
169	Polyhydroxylated GdDTPA-derivatives as high relaxivity magnetic resonance imaging contrast agents. RSC Advances, 2015, 5, 74734-74743.	1.7	6
170	The Role of the Amino Protecting Group during Parahydrogenation of Protected Dehydroamino Acids. Journal of Physical Chemistry A, 2015, 119, 11271-11279.	1.1	2
171	Effects of Magnetic Field Cycle on the Polarization Transfer from Parahydrogen to Heteronuclei through Long-Range J-Couplings. Journal of Physical Chemistry B, 2015, 119, 10035-10041.	1.2	66
172	Societ� Chimica Italiana Prizes 2014. Angewandte Chemie - International Edition, 2015, 54, 1070-1071.	7.2	0
173	Asparagine in plums detected by CEST-MRI. Food Chemistry, 2015, 169, 1-4.	4.2	6
174	A Quantitative Relaxometric Version of the ELISA Test for the Measurement of Cell Surface Biomarkers. Angewandte Chemie - International Edition, 2014, 53, 3488-3491.	7.2	48
175	Polymeric Vesicles Loaded with Gadoteridol as Reversible and Concentration-Independent Magnetic Resonance Imaging Thermometers. Journal of Biomedical Nanotechnology, 2014, 10, 1620-1626.	0.5	6
176	Highly Shifted Proton MR Imaging: Cell Tracking by Using Direct Detection of Paramagnetic Compounds. Radiology, 2014, 272, 785-795.	3.6	30
177	Role of the reaction intermediates in determining PHIP (parahydrogen induced polarization) effect in the hydrogenation of acetylene dicarboxylic acid with the complex [Rh (dppb)] ⁺ (dppb): Tj ETQq1 1 0.784314 rgBTi, @verlock140 Tf 50		
178	Combined inhibition of PI3K ² and PI3K ³ reduces fat mass by enhancing β -MSH-dependent sympathetic drive. Science Signaling, 2014, 7, ra110.	1.6	31
179	Towards a Successful Clinical Implementation of Fluorescence-Guided Surgery. Molecular Imaging and Biology, 2014, 16, 147-151.	1.3	23
180	B25716/1: a novel albumin-binding Gd-AAZTA MRI contrast agent with improved properties in tumor imaging. Journal of Biological Inorganic Chemistry, 2014, 19, 715-726.	1.1	34

#	ARTICLE	IF	CITATIONS
181	Design and testing of paramagnetic liposome-based CEST agents for MRI visualization of payload release on pH-induced and ultrasound stimulation. <i>Journal of Biological Inorganic Chemistry</i> , 2014, 19, 207-214.	1.1	20
182	Dual-modality gene reporter for in vivo imaging. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 415-420.	3.3	91
183	Lanthanide-Loaded Erythrocytes As Highly Sensitive Chemical Exchange Saturation Transfer MRI Contrast Agents. <i>Journal of the American Chemical Society</i> , 2014, 136, 638-641.	6.6	47
184	Gadolinium-binding cyclic hexapeptoids: synthesis and relaxometric properties. <i>Organic and Biomolecular Chemistry</i> , 2014, 12, 424-431.	1.5	44
185	Design of PLGA Based Nanoparticles for Imaging Guided Applications. <i>Molecular Pharmaceutics</i> , 2014, 11, 4100-4106.	2.3	45
186	Platinum(II)–Gadolinium(III) Complexes as Potential Single-Molecular Theranostic Agents for Cancer Treatment. <i>Angewandte Chemie - International Edition</i> , 2014, 53, 13225-13228.	7.2	77
187	Chemical exchange saturation transfer (CEST): an efficient tool for detecting molecular information on proteins' behaviour. <i>Analyst</i> , 2014, 139, 2687-2690.	1.7	26
188	MRI nanoprobe based on chemical exchange saturation transfer: Ln ^{III} chelates anchored on the surface of mesoporous silica nanoparticles. <i>Nanoscale</i> , 2014, 6, 9604-9607.	2.8	19
189	Frequency-Encoded MRI-CEST Agents Based on Paramagnetic Liposomes/RBC Aggregates. <i>Nano Letters</i> , 2014, 14, 6857-6862.	4.5	24
190	Successful Entrapping of Liposomes in Glucan Particles: An Innovative Micron-Sized Carrier to Deliver Water-Soluble Molecules. <i>Molecular Pharmaceutics</i> , 2014, 11, 3760-3765.	2.3	22
191	Preliminary in vitro and in vivo evaluation of liposomal nanoparticles for passive and active tumour targeting by scintigraphic and MRI imaging. <i>EJNMMI Physics</i> , 2014, 1, A82.	1.3	0
192	Synthesis of a carborane-containing cholesterol derivative and evaluation as a potential dual agent for MRI/BNCT applications. <i>Organic and Biomolecular Chemistry</i> , 2014, 12, 2457-2467.	1.5	41
193	Solution Structures, Stabilities, Kinetics, and Dynamics of DO3A and DO3A–Sulphonamide Complexes. <i>Inorganic Chemistry</i> , 2014, 53, 2858-2872.	1.9	50
194	Probing Protein Conformation in Cells by EPR Distance Measurements using Gd ³⁺ Spin Labeling. <i>Journal of the American Chemical Society</i> , 2014, 136, 13458-13465.	6.6	187
195	A General MRI-CEST Ratiometric Approach for pH Imaging: Demonstration of <i>In Vivo</i> pH Mapping with lobitridol. <i>Journal of the American Chemical Society</i> , 2014, 136, 14333-14336.	6.6	155
196	In vivo MRI visualization of release from liposomes triggered by local application of pulsed low-intensity non-focused ultrasound. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2014, 10, e901-e904.	1.7	18
197	In vivo maps of extracellular pH in murine melanoma by CEST–MRI. <i>Magnetic Resonance in Medicine</i> , 2014, 71, 326-332.	1.9	98
198	In vivo MRI visualization of different cell populations labeled with PARACEST agents. <i>Magnetic Resonance in Medicine</i> , 2013, 69, 1703-1711.	1.9	58

#	ARTICLE	IF	CITATIONS
199	Design and Synthesis of a β -Cyclodextrin Oligomer: A New Platform with Potential Application as a Dendrimeric MulticARRIER. <i>Chemistry - A European Journal</i> , 2013, 19, 12086-12092.	1.7	17
200	MRI Tracking of Macrophages Labeled with Glucan Particles Entrapping a Water Insoluble Paramagnetic Gd-Based Agent. <i>Molecular Imaging and Biology</i> , 2013, 15, 307-315.	1.3	16
201	Evaluation of ^{111}In -Labeled EPep and FibPep as Tracers for Fibrin SPECT Imaging. <i>Molecular Pharmaceutics</i> , 2013, 10, 4309-4321.	2.3	26
202	Picture of a chelate in exchange: the crystal structure of NaHoDOTMA, a semi-hydrated chelate. <i>Chemical Communications</i> , 2013, 49, 2320.	2.2	7
203	Nanoparticle-based chemical exchange saturation transfer (CEST) agents. <i>NMR in Biomedicine</i> , 2013, 26, 839-849.	1.6	62
204	^{15}N -methylated amino acids as efficient probes for MRI-DNP applications. <i>Contrast Media and Molecular Imaging</i> , 2013, 8, 417-421.	0.4	23
205	Controlled mixing of lanthanide(III) ions in coacervate core micelles. <i>Chemical Communications</i> , 2013, 49, 3736.	2.2	57
206	MRI Visualization of Melanoma Cells by Targeting Overexpressed Sialic Acid with a Gd(III)-DOTA- β Imaging Reporter. <i>Angewandte Chemie - International Edition</i> , 2013, 52, 1161-1164.	7.2	81
207	A Carborane Derivative Click-Reaction under Heterogeneous Conditions for the Synthesis of a Promising Lipophilic MRI/GdBNCT Agent. <i>Chemistry - A European Journal</i> , 2013, 19, 721-728.	1.7	32
208	SPECT imaging of fibrin using fibrin-binding peptides. <i>Contrast Media and Molecular Imaging</i> , 2013, 8, 229-237.	0.4	21
209	Equilibrium, Kinetic and Structural Studies of AAZTA Complexes with Ga^{3+} , In^{3+} and Cu^{2+} . <i>European Journal of Inorganic Chemistry</i> , 2013, 2013, 147-162.	1.0	49
210	Synthesis and characterization of an MRI Gd-based probe designed to target the translocator protein. <i>Magnetic Resonance in Chemistry</i> , 2013, 51, 116-122.	1.1	7
211	Lipid-Based Nanoparticles in Cardiovascular Molecular Imaging. <i>Current Cardiovascular Imaging Reports</i> , 2013, 6, 69-75.	0.4	1
212	Synthesis and Relaxometric Characterization of a MRI Gd-Based Probe Responsive to Glutamic Acid Decarboxylase Enzymatic Activity. <i>Journal of Medicinal Chemistry</i> , 2013, 56, 2466-2477.	2.9	21
213	Combined High Resolution NMR and ^1H and ^{17}O Relaxometric Study Sheds Light on the Solution Structure and Dynamics of the Lanthanide(III) Complexes of HPDO3A. <i>Inorganic Chemistry</i> , 2013, 52, 7130-7138.	1.9	52
214	On the Fate of MRI Gd-Based Contrast Agents in Cells. Evidence for Extensive Degradation of Linear Complexes upon Endosomal Internalization. <i>Analytical Chemistry</i> , 2013, 85, 5627-5631.	3.2	47
215	Curcumin/Gd Loaded Apoferritin: A Novel Theranostic Agent To Prevent Hepatocellular Damage in Toxic Induced Acute Hepatitis. <i>Molecular Pharmaceutics</i> , 2013, 10, 2079-2085.	2.3	61
216	Release of a Paramagnetic Magnetic Resonance Imaging Agent from Liposomes Triggered by Low Intensity Non-Focused Ultrasound. <i>Journal of Medical Imaging and Health Informatics</i> , 2013, 3, 356-366.	0.2	14

#	ARTICLE	IF	CITATIONS
217	NMR investigation of the equilibrium partitioning of a water-soluble bile salt protein carrier to phospholipid vesicles. <i>Proteins: Structure, Function and Bioinformatics</i> , 2013, 81, 1776-1791.	1.5	32
218	Gd loading by hypotonic swelling: an efficient and safe route for cellular labeling. <i>Contrast Media and Molecular Imaging</i> , 2013, 8, 475-486.	0.4	37
219	In vivo Magnetic Resonance Imaging Detection of Paramagnetic Liposomes Loaded with Amphiphilic Gadolinium(III) Complexes: Impact of Molecular Structure on Relaxivity and Excretion Efficiency. <i>ChemPlusChem</i> , 2013, 78, 712-722.	1.3	16
220	A Ratiometric Procedure to Assess Matrix Metalloproteinase Activity by Magnetic Resonance Imaging. <i>Angewandte Chemie - International Edition</i> , 2013, 52, 3926-3930.	7.2	30
221	Imaging the pH evolution of an acute kidney injury model by means of iopamidol, a MRI-CEST pH-responsive contrast agent. <i>Magnetic Resonance in Medicine</i> , 2013, 70, 859-864.	1.9	114
222	Lack of Plasma Protein Hemopexin Results in Increased Duodenal Iron Uptake. <i>PLoS ONE</i> , 2013, 8, e68146.	1.1	11
223	Boronated Compounds for Imaging Guided BNCT Applications. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2012, 12, 543-553.	0.9	25
224	Structure-Function Relationships of Iodinated Contrast Media and Risk of Nephrotoxicity. <i>Current Medicinal Chemistry</i> , 2012, 19, 736-743.	1.2	17
225	Dissociation Kinetics of Open Chain and Macrocyclic Gadolinium(III) Aminopolycarboxylate Complexes Related to Magnetic Resonance Imaging: Catalytic Effect of Endogenous Ligands. <i>Chemistry - A European Journal</i> , 2012, 18, 16426-16435.	1.7	87
226	Mn loaded apoferritin as an MRI sensor of melanin formation in melanoma cells. <i>Chemical Communications</i> , 2012, 48, 2436.	2.2	25
227	Nature-inspired particles as carriers for multimodal molecular imaging applications. , 2012, , .		0
228	¹⁵ N Magnetic Resonance Hyperpolarization via the Reaction of Parahydrogen with ¹⁵ N-Propargylcholine. <i>Journal of the American Chemical Society</i> , 2012, 134, 11146-11152.	6.6	38
229	Assessment of iron absorption in mice by ICP-MS measurements of ⁵⁷ Fe levels. <i>European Journal of Nutrition</i> , 2012, 51, 783-789.	1.8	19
230	Hyperpolarized ¹³ C-pyruvate magnetic resonance imaging in cancer diagnostics. <i>Expert Opinion on Medical Diagnostics</i> , 2012, 6, 335-345.	1.6	10
231	In vivo imaging of inhibitory, GABAergic neurons by MRI. <i>NeuroImage</i> , 2012, 62, 1685-1693.	2.1	17
232	Para-hydrogen induced polarization of Si-29 NMR resonances as a potentially useful tool for analytical applications. <i>Magnetic Resonance in Chemistry</i> , 2012, 50, 529-533.	1.1	8
233	Supramolecular Adducts of Negatively Charged Lanthanide(III) DOTP Chelates and Cyclodextrins Functionalized with Ammonium Groups: Mass Spectrometry and Nuclear Magnetic Resonance Studies. <i>European Journal of Inorganic Chemistry</i> , 2012, 2012, 2087-2098.	1.0	6
234	MRI Paramagnetic Probes for Cellular Labeling. <i>European Journal of Inorganic Chemistry</i> , 2012, 2012, 1934-1944.	1.0	24

#	ARTICLE	IF	CITATIONS
235	High Relaxivity Supramolecular Adducts Between Human Liver Fatty Acid Binding Protein and Amphiphilic Gd ^{III} Complexes: Structural Basis for the Design of Intracellular Targeting MRI Probes. <i>Chemistry - A European Journal</i> , 2012, 18, 9919-9928.	1.7	25
236	Synthesis and Characterization of a Hypoxia Sensitive MRI Probe. <i>Chemistry - A European Journal</i> , 2012, 18, 9669-9676.	1.7	47
237	Hyperpolarized ¹³ C-labelled anhydrides as DNP precursors of metabolic MRI agents. <i>Contrast Media and Molecular Imaging</i> , 2012, 7, 469-477.	0.4	10
238	Novel Gd(III)-based probes for MR molecular imaging of matrix metalloproteinases. <i>Contrast Media and Molecular Imaging</i> , 2012, 7, 175-184.	0.4	23
239	Mn-loaded apoferritin: a highly sensitive MRI imaging probe for the detection and characterization of hepatocarcinoma lesions in a transgenic mouse model. <i>Contrast Media and Molecular Imaging</i> , 2012, 7, 281-288.	0.4	34
240	Paramagnetic self-assembled nanoparticles as supramolecular MRI contrast agents. <i>Contrast Media and Molecular Imaging</i> , 2012, 7, 356-361.	0.4	57
241	Relaxometric Investigations and MRI Evaluation of a Liposome-Loaded pH-Responsive Gadolinium(III) Complex. <i>Inorganic Chemistry</i> , 2012, 51, 7210-7217.	1.9	35
242	A vaccine targeting angiominin induces an antibody response which alters tumor vessel permeability and hampers the growth of established tumors. <i>Angiogenesis</i> , 2012, 15, 305-316.	3.7	35
243	MRI evaluation of the antitumor activity of paramagnetic liposomes loaded with prednisolone phosphate. <i>European Journal of Pharmaceutical Sciences</i> , 2012, 45, 436-441.	1.9	34
244	Image guided therapy: The advent of theranostic agents. <i>Journal of Controlled Release</i> , 2012, 161, 328-337.	4.8	136
245	Yeast cell wall particles: a promising class of nature-inspired microcarriers for multimodal imaging. <i>Chemical Communications</i> , 2011, 47, 10635.	2.2	31
246	Gadolinium-doped LipoCEST agents: a potential novel class of dual 1H-MRI probes. <i>Chemical Communications</i> , 2011, 47, 4667.	2.2	31
247	Synthesis and Physicochemical Characterization of Carbon Backbone Modified [Gd(TTDA)(H ₂ O)] ₂ Derivatives. <i>Inorganic Chemistry</i> , 2011, 50, 1275-1287.	1.9	16
248	Properties, Solution State Behavior, and Crystal Structures of Chelates of DOTMA. <i>Inorganic Chemistry</i> , 2011, 50, 7955-7965.	1.9	86
249	Contrast agents and mechanisms. <i>Drug Discovery Today: Technologies</i> , 2011, 8, e109-e115.	4.0	14
250	Dual MRI-SPECT agent for pH-mapping. <i>Chemical Communications</i> , 2011, 47, 1539-1541.	2.2	49
251	In Vivo Labeling of B16 Melanoma Tumor Xenograft with a Thiol-Reactive Gadolinium Based MRI Contrast Agent. <i>Molecular Pharmaceutics</i> , 2011, 8, 1750-1756.	2.3	17
252	MRI Contrast Agents: State of the Art and New Trends. , 2011, , 223-251.		2

#	ARTICLE	IF	CITATIONS
253	I^{2} -Gal Gene Expression MRI Reporter in Melanoma Tumor Cells. Design, Synthesis, and <i>in Vitro</i> and <i>in Vivo</i> Testing of a Gd(III) Containing Probe Forming a High Relaxivity, Melanin-Like Structure upon I^{2} -Gal Enzymatic Activation. <i>Bioconjugate Chemistry</i> , 2011, 22, 2625-2635.	1.8	51
254	Theranostic Nanomedicine. <i>Accounts of Chemical Research</i> , 2011, 44, 1029-1038.	7.6	765
255	Society Spotlight: European Commission-supported development of targeted nanomedicines: did MediTrans make a difference?. <i>Therapeutic Delivery</i> , 2011, 2, 861-864.	1.2	0
256	Improved paramagnetic liposomes for MRI visualization of pH triggered release. <i>Journal of Controlled Release</i> , 2011, 154, 196-202.	4.8	58
257	Dose estimation in B16 tumour bearing mice for future irradiation in the thermal column of the TRIGA reactor after B/Gd/LDL adduct infusion. <i>Applied Radiation and Isotopes</i> , 2011, 69, 1842-1845.	0.7	0
258	Supramolecular protamine/Gd-loaded liposomes adducts as relaxometric protease responsive probes. <i>Bioorganic and Medicinal Chemistry</i> , 2011, 19, 1131-1135.	1.4	14
259	Synthesis and testing of a 2 hyperpolarized ^{13}C probe based on the pyrazolo[1,5- <i>a</i>]pyrimidineacetamide DPA-713, an MRI vector to target the peripheral benzodiazepine receptors. <i>Magnetic Resonance in Chemistry</i> , 2011, 49, 795-800.	1.1	4
260	Photochemical activation of endosomal escape of MRI-Gd agents in tumor cells. <i>Magnetic Resonance in Medicine</i> , 2011, 65, 212-219.	1.9	25
261	Iopamidol as a responsive MRI-chemical exchange saturation transfer contrast agent for pH mapping of kidneys: In vivo studies in mice at 7 T. <i>Magnetic Resonance in Medicine</i> , 2011, 65, 202-211.	1.9	178
262	Relaxometric Study of a Series of Monoaquated Gd^{III} Complexes of Rigidified EGTA-Like Chelators and Their Noncovalent Interaction with Human Serum Albumin. <i>European Journal of Inorganic Chemistry</i> , 2011, 2011, 802-810.	1.0	25
263	How to design ^{13}C para-hydrogen-induced polarization experiments for MRI applications. <i>Contrast Media and Molecular Imaging</i> , 2011, 6, 77-84.	0.4	23
264	MRI of cells and mice at 1 and 7% Tesla with Gd-targeting agents: when the low field is better!. <i>Contrast Media and Molecular Imaging</i> , 2011, 6, 421-425.	0.4	22
265	Yb^{III} -HPDO3A: A Dual pH- and Temperature-Responsive CEST Agent. <i>Angewandte Chemie - International Edition</i> , 2011, 50, 1798-1800.	7.2	103
266	Use of Labile Precursors for the Generation of Hyperpolarized Molecules from Hydrogenation with Parahydrogen and Aqueous-Phase Extraction. <i>Angewandte Chemie - International Edition</i> , 2011, 50, 7350-7353.	7.2	55
267	MRI-Guided Neutron Capture Therapy by Use of a Dual Gadolinium/Boron Agent Targeted at Tumour Cells through Upregulated Low-Density Lipoprotein Transporters. <i>Chemistry - A European Journal</i> , 2011, 17, 8479-8486.	1.7	56
268	Imaging probes-Introduction. <i>Bioorganic and Medicinal Chemistry</i> , 2011, 19, 1022.	1.4	1
269	Advances in bio-imaging: A survey from WWMR 2010. <i>Journal of Magnetic Resonance</i> , 2011, 209, 109-115.	1.2	4
270	The use of contrast agents with fast field-cycling magnetic resonance imaging. <i>Physics in Medicine and Biology</i> , 2011, 56, 105-115.	1.6	12

#	ARTICLE	IF	CITATIONS
271	MR Contrast Agents. , 2011, , 165-193.		0
272	Fast field-cycling magnetic resonance imaging. <i>Comptes Rendus Physique</i> , 2010, 11, 136-148.	0.3	63
273	In vivo MRI multicontrast kinetic analysis of the uptake and intracellular trafficking of paramagnetically labeled liposomes. <i>Journal of Controlled Release</i> , 2010, 144, 271-279.	4.8	64
274	Encoding the frequency dependence in MRI contrast media: the emerging class of CEST agents. <i>Contrast Media and Molecular Imaging</i> , 2010, 5, 78-98.	0.4	113
275	Novel MRI and fluorescent probes responsive to the Factor XIII transglutaminase activity. <i>Contrast Media and Molecular Imaging</i> , 2010, 5, 213-222.	0.4	22
276	Target Visualization by MRI Using the Avidin/Biotin Amplification Route: Synthesis and Testing of a Biotin- ⁶⁷ Gd-DOTA Monoamide Trimer. <i>Chemistry - A European Journal</i> , 2010, 16, 8080-8087.	1.7	22
277	Reduction/Dissolution of a ⁵⁵ MnOOH Nanophase in the Ferritin Cavity To Yield a Highly Sensitive, Biologically Compatible Magnetic Resonance Imaging Agent. <i>Angewandte Chemie - International Edition</i> , 2010, 49, 612-615.	7.2	36
278	Current concepts on hyperpolarized molecules in MRI. <i>Current Opinion in Chemical Biology</i> , 2010, 14, 90-96.	2.8	73
279	Improving the relaxivity by dimerizing Gd-AAZTA: Insights for enhancing the sensitivity of MRI contrast agents. <i>Inorganic Chemistry Communication</i> , 2010, 13, 663-665.	1.8	16
280	Synthesis and characterization of a novel macrocyclic ligand containing nine donor atoms. <i>Recueil Des Travaux Chimiques Des Pays-Bas</i> , 2010, 115, 94-98.	0.0	2
281	Combined Delivery and Magnetic Resonance Imaging of Neural Cell Adhesion Molecule-Targeted Doxorubicin-Containing Liposomes in Experimentally Induced Kaposi's Sarcoma. <i>Cancer Research</i> , 2010, 70, 2180-2190.	0.4	90
282	Advances in Metal-Based Probes for MR Molecular Imaging Applications. <i>Current Medicinal Chemistry</i> , 2010, 17, 3684-3700.	1.2	47
283	Block copolymer vesicles containing paramagnetic lanthanide complexes: a novel class of T1- and CEST MRI contrast agents. <i>Soft Matter</i> , 2010, 6, 4847.	1.2	24
284	Challenges for Molecular Magnetic Resonance Imaging. <i>Chemical Reviews</i> , 2010, 110, 3019-3042.	23.0	728
285	Para-hydrogenated Glucose Derivatives as Potential ¹³ C-Hyperpolarized Probes for Magnetic Resonance Imaging. <i>Journal of the American Chemical Society</i> , 2010, 132, 7186-7193.	6.6	55
286	A copper-activated magnetic resonance imaging contrast agent with improved turn-on relaxivity response and anion compatibility. <i>Dalton Transactions</i> , 2010, 39, 469-476.	1.6	51
287	Exofacial Protein Thiols as a Route for the Internalization of Gd(III)-Based Complexes for Magnetic Resonance Imaging Cell Labeling. <i>Journal of Medicinal Chemistry</i> , 2010, 53, 4877-4890.	2.9	46
288	Tuning Glutamine Binding Modes in Gd-DOTA-Based Probes for an Improved MRI Visualization of Tumor Cells. <i>Chemistry - A European Journal</i> , 2009, 15, 76-85.	1.7	19

#	ARTICLE	IF	CITATIONS
289	Osmotically Shrunken LIPOCEST Agents: An Innovative Class of Magnetic Resonance Imaging Contrast Media Based on Chemical Exchange Saturation Transfer. <i>Chemistry - A European Journal</i> , 2009, 15, 1440-1448.	1.7	50
290	Equilibrium and Kinetic Properties of the Lanthanoids(III) and Various Divalent Metal Complexes of the Heptadentate Ligand AAZTA. <i>Chemistry - A European Journal</i> , 2009, 15, 1696-1705.	1.7	90
291	Release of Toxic Gd ³⁺ Ions to Tumour Cells by Vitamin B ₁₂ Bioconjugates. <i>Chemistry - A European Journal</i> , 2009, 15, 7980-7989.	1.7	33
292	Lanthanide(III) Complexes of Tris(amide) PCTA Derivatives as Potential Bimodal Magnetic Resonance and Optical Imaging Agents. <i>Chemistry - A European Journal</i> , 2009, 15, 13188-13200.	1.7	38
293	Synthesis and Relaxometric Properties of Gadolinium(III) Complexes of New Triazine-Based Polydentate Ligands. <i>Helvetica Chimica Acta</i> , 2009, 92, 2414-2426.	1.0	8
294	Effect of a mesitylene-based ligand cap on the relaxometric properties of Gd(III) hydroxypyridonate MRI contrast agents. <i>Contrast Media and Molecular Imaging</i> , 2009, 4, 220-229.	0.4	13
295	Methods for an improved detection of the MRI-CEST effect. <i>Contrast Media and Molecular Imaging</i> , 2009, 4, 237-247.	0.4	54
296	CMR2009: 5.09: B25716/1: a novel Gd-AAZTA-based MRI agent with albumin binding properties. <i>Contrast Media and Molecular Imaging</i> , 2009, 4, 276-277.	0.4	0
297	CMR2009: 11.02: Evaluating iopamidol as pH-responsive CEST agent at 3 and 7 T. <i>Contrast Media and Molecular Imaging</i> , 2009, 4, 294-295.	0.4	5
298	Biodistribution of gadolinium-based contrast agents, including gadolinium deposition. <i>Journal of Magnetic Resonance Imaging</i> , 2009, 30, 1259-1267.	1.9	444
299	Evidence for <i>in vivo</i> macrophage mediated tumor uptake of paramagnetic/fluorescent liposomes. <i>NMR in Biomedicine</i> , 2009, 22, 1084-1092.	1.6	36
300	Towards improved boron neutron capture therapy agents: evaluation of <i>in vitro</i> cellular uptake of a glutamine-functionalized carborane. <i>Journal of Biological Inorganic Chemistry</i> , 2009, 14, 883-890.	1.1	9
301	Effect of low and zero magnetic field on the hyperpolarization lifetime in parahydrogenated perdeuterated molecules. <i>Journal of Magnetic Resonance</i> , 2009, 200, 15-20.	1.2	10
302	Copper-Responsive Magnetic Resonance Imaging Contrast Agents. <i>Journal of the American Chemical Society</i> , 2009, 131, 8527-8536.	6.6	139
303	Isolation of a ¹⁷⁷ Hf Complex Formed by ^β -Decay of a ¹⁷⁷ Lu-Labeled Radiotherapeutic Compound and NMR Structural Elucidation of the Ligand and its Lu and Hf Complexes. <i>Inorganic Chemistry</i> , 2009, 48, 3114-3124.	1.9	4
304	Chemistry of Molecular Imaging. <i>Accounts of Chemical Research</i> , 2009, 42, 821-821.	7.6	23
305	High-Relaxivity Gadolinium-Modified High-Density Lipoproteins as Magnetic Resonance Imaging Contrast Agents. <i>Journal of Physical Chemistry B</i> , 2009, 113, 6283-6289.	1.2	62
306	1,2-Hydroxypyridonate/Terephthalamide Complexes of Gadolinium(III): Synthesis, Stability, Relaxivity, and Water Exchange Properties. <i>Inorganic Chemistry</i> , 2009, 48, 277-286.	1.9	40

#	ARTICLE	IF	CITATIONS
307	Relaxometric Studies for Food Characterization: The Case of Balsamic and Traditional Balsamic Vinegars. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 3028-3032.	2.4	34
308	Thermodynamic analysis of hydration in human serum heme- α -albumin. <i>Biochemical and Biophysical Research Communications</i> , 2009, 385, 385-389.	1.0	6
309	Poly- β -cyclodextrin based platform for pH mapping via a ratiometric $^{19}\text{F}/^1\text{H}$ MRI method. <i>Chemical Communications</i> , 2009, , 6044.	2.2	60
310	Well-Defined, Multifunctional Nanostructures of a Paramagnetic Lipid and a Lipopeptide for Macrophage Imaging. <i>Journal of the American Chemical Society</i> , 2009, 131, 406-407.	6.6	28
311	Pushing the Sensitivity Envelope of Lanthanide-Based Magnetic Resonance Imaging (MRI) Contrast Agents for Molecular Imaging Applications. <i>Accounts of Chemical Research</i> , 2009, 42, 822-831.	7.6	327
312	Lanthanide-Loaded Paramagnetic Liposomes as Switchable Magnetically Oriented Nanovesicles. <i>Methods in Enzymology</i> , 2009, 464, 193-210.	0.4	21
313	Targeting exofacial protein thiols with Gd(III) complexes. An efficient procedure for MRI cell labelling. <i>Chemical Communications</i> , 2009, , 893.	2.2	20
314	New cyclodextrin dimers and trimers capable of forming supramolecular adducts with shape-specific ligands. <i>Organic and Biomolecular Chemistry</i> , 2009, 7, 370-379.	1.5	42
315	Variation of water exchange dynamics with ligand structure and stereochemistry in lanthanide complexes based on 1,4-diazepine derivatives. <i>Organic and Biomolecular Chemistry</i> , 2009, 7, 1120.	1.5	34
316	Hyperpolarized agents for advanced MRI investigations. <i>Quarterly Journal of Nuclear Medicine and Molecular Imaging</i> , 2009, 53, 604-17.	0.4	37
317	Contrast-enhanced MRI of murine sponge model for progressive angiogenesis assessed with gadoteridol (ProHance) and gadocoletic acid trisodium salt (B22956/1). <i>Journal of Magnetic Resonance Imaging</i> , 2008, 27, 872-880.	1.9	6
318	Characterization of human hair melanin and its degradation products by means of magnetic resonance techniques. <i>Magnetic Resonance in Chemistry</i> , 2008, 46, 471-479.	1.1	33
319	^1H and ^{17}O NMR relaxometric study in aqueous solution of Gd(III) complexes of EGTA-like derivatives bearing methylenephosphonic groups. <i>Magnetic Resonance in Chemistry</i> , 2008, 46, S86-S93.	1.1	14
320	High-Relaxivity Magnetic Resonance Imaging (MRI) Contrast Agent Based on Supramolecular Assembly between a Gadolinium Chelate, a Modified Dextran, and Poly- β -Cyclodextrin. <i>Chemistry - A European Journal</i> , 2008, 14, 4551-4561.	1.7	99
321	A Novel Method of Cellular Labeling: Anchoring MR-Imaging Reporter Particles on the Outer Cell Surface. <i>ChemMedChem</i> , 2008, 3, 60-62.	1.6	24
322	Efficient Cellular Labeling by CD44 Receptor-Mediated Uptake of Cationic Liposomes Functionalized with Hyaluronic Acid and Loaded with MRI Contrast Agents. <i>ChemMedChem</i> , 2008, 3, 1858-1862.	1.6	31
323	First <i>in vivo</i> MRI co-localization of two LIPOCEST agents. <i>Contrast Media and Molecular Imaging</i> , 2008, 3, 38-43.	0.4	46
324	Development and validation of a smoothing-splines-based correction method for improving the analysis of CEST-MR images. <i>Contrast Media and Molecular Imaging</i> , 2008, 3, 136-149.	0.4	102

#	ARTICLE	IF	CITATIONS
325	Intramolecular Host-Guest Dynamics of FeCp(CO)2X (X = I and CH3) and Mo2Cp2(CO)6 Included in β - or γ -Cyclodextrin. <i>European Journal of Inorganic Chemistry</i> , 2008, 2008, 152-157.	1.0	8
326	The Detection of PHIP Effects Allows New Insights into the Mechanism of Olefin Isomerisation during Catalytic Hydrogenation. <i>European Journal of Inorganic Chemistry</i> , 2008, 2008, 4348-4351.	1.0	9
327	Paramagnetic Liposomes as Innovative Contrast Agents for Magnetic Resonance (MR) Molecular Imaging Applications. <i>Chemistry and Biodiversity</i> , 2008, 5, 1901-1912.	1.0	76
328	NorDATA: An original ligand based on the norbornane skeleton. Synthesis and thermodynamic characterization of metal complexes. <i>Polyhedron</i> , 2008, 27, 3683-3687.	1.0	3
329	Determination of water permeability of paramagnetic liposomes of interest in MRI field. <i>Journal of Inorganic Biochemistry</i> , 2008, 102, 1112-1119.	1.5	70
330	Metal containing nanosized systems for MR-Molecular Imaging applications. <i>Coordination Chemistry Reviews</i> , 2008, 252, 2424-2443.	9.5	116
331	An unusual gadolinium ten-coordinated dimeric complex in the series of MRI contrast agents: Na[Gd(H2O)AAZTA]·3H2O. <i>Inorganica Chimica Acta</i> , 2008, 361, 1534-1541.	1.2	29
332	High Relaxivity Gadolinium Hydroxypyridonate-Viral Capsid Conjugates: Nanosized MRI Contrast Agents. <i>Journal of the American Chemical Society</i> , 2008, 130, 2546-2552.	6.6	165
333	New Hyperpolarized Contrast Agents for ¹³ C-MRI from Para-Hydrogenation of Oligooxyethylene Alkynes. <i>Journal of the American Chemical Society</i> , 2008, 130, 15047-15053.	6.6	58
334	Functionalized Nanocontainers as Dual Magnetic and Optical Probes for Molecular Imaging Applications. <i>Chemistry of Materials</i> , 2008, 20, 5888-5893.	3.2	73
335	Synthesis and solution thermodynamic study of rigidified and functionalised EGTA derivatives. <i>Organic and Biomolecular Chemistry</i> , 2008, 6, 2361.	1.5	23
336	Bifunctional ligands based on the DOTA-monoamide cage. <i>Organic and Biomolecular Chemistry</i> , 2008, 6, 1176.	1.5	49
337	Carbon coated microshells containing nanosized Gd(III) oxides for multiple bio-medical applications. <i>Chemical Communications</i> , 2008, , 5936.	2.2	5
338	Synthesis of Gd(III)-C-palmitamidomethyl- ϵ -DOTAMA-C6-o-carborane: a new dual agent for innovative MRI/BNCT applications. <i>Organic and Biomolecular Chemistry</i> , 2008, 6, 4460.	1.5	33
339	Lanthanide-Loaded Paramagnetic Liposomes as Switchable Magnetically Oriented Nanovesicles. <i>Inorganic Chemistry</i> , 2008, 47, 2928-2930.	1.9	26
340	Highly shifted LIPOCEST agents based on the encapsulation of neutral polynuclear paramagnetic shift reagents. <i>Chemical Communications</i> , 2008, , 600-602.	2.2	38
341	Synthesis, Potentiometric, Kinetic, and NMR Studies of 1,4,7,10-Tetraazacyclododecane-1,7-bis(acetic) Lanthanide(III) Ions. <i>Inorganic Chemistry</i> , 2008, 47, 3851-3862.	1.9	65
342	Solution Structure of the Supramolecular Adduct between a Liver Cytosolic Bile Acid Binding Protein and a Bile Acid-Based Gadolinium(III)-Chelate, a Potential Hepatospecific Magnetic Resonance Imaging Contrast Agent. <i>Journal of Medicinal Chemistry</i> , 2008, 51, 6782-6792.	2.9	18

#	ARTICLE	IF	CITATIONS
343	A New, Easy Access to the 6-Aminoperhydro-1,4-diazepine Scaffold under Ultrasound and Microwave Irradiation. <i>Synthesis</i> , 2008, 2008, 1879-1882.	1.2	1
344	Gadolinium-Enhanced Magnetic Resonance Imaging, Renal Failure and Nephrogenic Systemic Fibrosis /Nephrogenic Fibrosing Dermopathy. <i>Current Medicinal Chemistry</i> , 2008, 15, 1229-1235.	1.2	55
345	Response to "Advisory about gadolinium calls for caution in the treatment of uremic patients with lanthanum carbonate"™. <i>Kidney International</i> , 2008, 74, 536-537.	2.6	0
346	Agents for Polarization Enhancement in MRI. <i>Handbook of Experimental Pharmacology</i> , 2008, , 247-272.	0.9	25
347	Advisory about gadolinium calls for caution in the treatment of uremic patients with lanthanum carbonate. <i>Kidney International</i> , 2007, 72, 1162-1163.	2.6	17
348	Maximizing the relaxivity of HSA-bound gadolinium complexes by simultaneous optimization of rotation and water exchange. <i>Chemical Communications</i> , 2007, , 4726.	2.2	49
349	Synthesis and characterization of a Gd(III) based contrast agent responsive to thiol containing compounds. <i>Dalton Transactions</i> , 2007, , 4980.	1.6	36
350	Gd-Loaded Liposomes as T1, Susceptibility, and CEST Agents, All in One. <i>Journal of the American Chemical Society</i> , 2007, 129, 2430-2431.	6.6	83
351	1,2-Hydroxypyridonates as Contrast Agents for Magnetic Resonance Imaging: " TREN-1,2-HOPO. <i>Inorganic Chemistry</i> , 2007, 46, 9182-9191.	1.9	58
352	Highly Soluble Tris-hydroxypyridonate Gd(III) Complexes with Increased Hydration Number, Fast Water Exchange, Slow Electronic Relaxation, and High Relaxivity1. <i>Journal of the American Chemical Society</i> , 2007, 129, 1870-1871.	6.6	97
353	Optimized Relaxivity and Stability of [Gd(H(2,2)-1,2-HOPO)(H2O)]-for Use as an MRI Contrast Agent1. <i>Inorganic Chemistry</i> , 2007, 46, 4796-4798.	1.9	39
354	NMR Structural Studies of the Supramolecular Adducts between a Liver Cytosolic Bile Acid Binding Protein and Gadolinium(III)-Chelates Bearing Bile Acids Residues: " Molecular Determinants of the Binding of a Hepatospecific Magnetic Resonance Imaging Contrast Agent. <i>Journal of Medicinal Chemistry</i> , 2007, 50, 5257-5268.	2.9	17
355	Magnetic Resonance Imaging Detection of Tumor Cells by Targeting Low-Density Lipoprotein Receptors with Gd-Loaded Low-Density Lipoprotein Particles. <i>Neoplasia</i> , 2007, 9, 1046-1056.	2.3	59
356	Magnetic resonance imaging of gadolinium-labeled pancreatic islets for experimental transplantation. <i>NMR in Biomedicine</i> , 2007, 20, 40-48.	1.6	85
357	Peptide Derivatized Lamellar Aggregates as Target-Specific MRI Contrast Agents. <i>ChemBioChem</i> , 2007, 8, 950-955.	1.3	23
358	Relaxometric and Modelling Studies of the Binding of a Lipophilic Gd-AAZTA Complex to Fatted and Defatted Human Serum Albumin. <i>Chemistry - A European Journal</i> , 2007, 13, 5785-5797.	1.7	93
359	From Spherical to Osmotically Shrunken Paramagnetic Liposomes: An Improved Generation of LIPOCEST MRI Agents with Highly Shifted Water Protons. <i>Angewandte Chemie - International Edition</i> , 2007, 46, 966-968.	7.2	87
360	The effects of intramolecular H-bond formation on the stability constant and water exchange rate of the Gd(III)-diethylenetriamine-Na ²⁻ -(3-amino-1,1-propylenephosphonic)-N,N,N ³⁻ ,N ³⁻ -tetraacetate complex. <i>Contrast Media and Molecular Imaging</i> , 2007, 2, 94-102.	0.4	17

#	ARTICLE	IF	CITATIONS
361	Molecular recognition of sugars by lanthanide (III) complexes of a conjugate of $\langle i \rangle N, N \langle /i \rangle$ -bis[2- $\langle i \rangle$ bis[2-(1,1-dimethylethoxy)- $\langle i \rangle$ -2-oxoethyl]amino]ethyl]glycine and phenylboronic acid. <i>Contrast Media and Molecular Imaging</i> , 2007, 2, 163-171.	1.4	17
362	CMR 2007: 7.05: Gd-loaded nanosized systems for MR visualization of tumor cells. <i>Contrast Media and Molecular Imaging</i> , 2007, 2, 290-291.	0.4	0
363	Targeting of human renal tumor-derived endothelial cells with peptides obtained by phage display. <i>Journal of Molecular Medicine</i> , 2007, 85, 897-906.	1.7	17
364	New CD derivatives as self-assembling contrast agents for magnetic resonance imaging (MRI). <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2007, 57, 489-495.	1.6	19
365	An iron-based T1 contrast agent made of iron-phosphate complexes: In vitro and in vivo studies. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2007, 20, 27-37.	1.1	15
366	High-relaxivity supramolecular aggregates containing peptides and Gd complexes as contrast agents in MRI. <i>Journal of Biological Inorganic Chemistry</i> , 2007, 12, 267-276.	1.1	39
367	On the role of the counter-ion in defining water structure and dynamics: order, structure and dynamics in hydrophilic and hydrophobic gadolinium salt complexes. <i>Dalton Transactions</i> , 2006, , 5605.	1.6	54
368	Relaxometric and solution NMR structural studies on ditopic lanthanide(iii) complexes of a phosphinate analogue of DOTA with a fast rate of water exchange. <i>Dalton Transactions</i> , 2006, , 2323.	1.6	44
369	Glycoconjugates of gadolinium complexes for MRI applications. <i>Chemical Communications</i> , 2006, , 1064.	2.2	84
370	A responsive MRI contrast agent to monitor functional cell status. <i>NeuroImage</i> , 2006, 32, 1142-1149.	2.1	63
371	Neural-cell adhesion molecule (NCAM) expression by immature and tumor-derived endothelial cells favors cell organization into capillary-like structures. <i>Experimental Cell Research</i> , 2006, 312, 913-924.	1.2	46
372	A ^{13}C CP/MAS NMR Study of the Structure and Dynamics of $[(\beta\text{-5-C5H5})_2\text{Fe}_2(\text{CO})_4]$ Included in $\beta\text{-Cyclodextrin}$: Evidence for Terminal Bridging Exchange in the cis Isomer. <i>Organometallics</i> , 2006, 25, 2248-2252.	1.1	7
373	Improved syntheses of bis($\beta\text{-cyclodextrin}$) derivatives, new carriers for gadolinium complexes. <i>Organic and Biomolecular Chemistry</i> , 2006, 4, 1124.	1.5	29
374	A R2/R1 Ratio Procedure for a Concentration-Independent, pH-Responsive, Gd(III)-Based MRI Agent. <i>Journal of the American Chemical Society</i> , 2006, 128, 11326-11327.	6.6	92
375	Substituent Effects on Gd(III)-Based MRI Contrast Agents: Optimizing the Stability and Selectivity of the Complex and the Number of Coordinated Water Molecules. <i>Inorganic Chemistry</i> , 2006, 45, 8355-8364.	1.9	82
376	In Vitro and in Vivo Magnetic Resonance Detection of Tumor Cells by Targeting Glutamine Transporters with Gd-Based Probes. <i>Journal of Medicinal Chemistry</i> , 2006, 49, 4926-4936.	2.9	69
377	Fe(III)-Templated Gd(III) Self-Assemblies A New Route toward Macromolecular MRI Contrast Agents. <i>Journal of the American Chemical Society</i> , 2006, 128, 9272-9273.	6.6	46
378	PAMAM Dendrimeric Conjugates with a Gd-DOTA Phosphinate Derivative and Their Adducts with Polyaminoacids: The Interplay of Global Motion, Internal Rotation, and Fast Water Exchange. <i>Bioconjugate Chemistry</i> , 2006, 17, 975-987.	1.8	108

#	ARTICLE	IF	CITATIONS
379	Tuning the Coordination Number of Hydroxypyridonate-Based Gadolinium Complexes: Implications for MRI Contrast Agents. <i>Journal of the American Chemical Society</i> , 2006, 128, 5344-5345.	6.6	49
380	High sensitivity lanthanide(III) based probes for MR-medical imaging. <i>Coordination Chemistry Reviews</i> , 2006, 250, 1562-1579.	9.5	284
381	The crystallized solvent could influence the lanthanide water bonding?. <i>Inorganica Chimica Acta</i> , 2006, 359, 3405-3411.	1.2	25
382	Polarization transfer from para-hydrogen to heteronuclei: Effect of H/D substitution. The case of AA ² X and spin systems. <i>Journal of Magnetic Resonance</i> , 2006, 178, 184-192.	1.2	42
383	Magnetic resonance imaging of tumor cells by targeting the amino acid transport system. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2006, 16, 4111-4114.	1.0	14
384	New paramagnetic supramolecular adducts for MRI applications based on non-covalent interactions between Gd(III)-complexes and β - or γ -cyclodextrin units anchored to chitosan. <i>Journal of Inorganic Biochemistry</i> , 2006, 100, 931-938.	1.5	31
385	CMR 2005: 13.07: Novel applications in the field of MRI CEST agents. <i>Contrast Media and Molecular Imaging</i> , 2006, 1, 89-89.	0.4	0
386	Cellular labeling with Gd(III) chelates: only high thermodynamic stabilities prevent the cells acting as "sponges" of Gd ³⁺ ions. <i>Contrast Media and Molecular Imaging</i> , 2006, 1, 23-29.	0.4	89
387	Catalytic effects of apoferritin interior surface residues on water proton exchange in lanthanide complexes. <i>Contrast Media and Molecular Imaging</i> , 2006, 1, 10-14.	0.4	32
388	Gd-enhanced MR images of substrates other than water. <i>Contrast Media and Molecular Imaging</i> , 2006, 1, 101-105.	0.4	14
389	Para-hydrogen enrichment and hyperpolarization. <i>Concepts in Magnetic Resonance Part A: Bridging Education and Research</i> , 2006, 28A, 321-330.	0.2	73
390	Effect of the intracellular localization of a Gd-based imaging probe on the relaxation enhancement of water protons. <i>Magnetic Resonance in Medicine</i> , 2006, 55, 491-497.	1.9	158
391	Quantification of the expression level of integrin receptor $\alpha_5\beta_3$ in cell lines and MR imaging with antibody-coated iron oxide particles. <i>Magnetic Resonance in Medicine</i> , 2006, 56, 711-716.	1.9	51
392	Water exchange across the erythrocyte plasma membrane studied by HR-MAS NMR spectroscopy. <i>Magnetic Resonance in Medicine</i> , 2006, 56, 978-985.	1.9	13
393	Magnetic Resonance Visualization of Tumor Angiogenesis by Targeting Neural Cell Adhesion Molecules with the Highly Sensitive Gadolinium-Loaded Apoferritin Probe. <i>Cancer Research</i> , 2006, 66, 9196-9201.	0.4	132
394	Paramagnetic Metal Complexes As Contrast Agents for Magnetic Resonance Imaging. , 2005, , 541-560.		1
395	Gadolinium(III) Complexes of dota-Derived N-Sulfonylacetamides (H ₄ (dota-NHSO ₂ R)=10-{2-[(R)sulfonylamino]-2-oxoethyl}-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic) <i>Inorganica Chimica Acta</i> , 2005, 88, 588-603.	1.0	12
396	Tunable Imaging of Cells Labeled with MRI-PARACEST Agents. <i>Angewandte Chemie - International Edition</i> , 2005, 44, 1813-1815.	7.2	170

#	ARTICLE	IF	CITATIONS
397	Highly Sensitive MRI Chemical Exchange Saturation Transfer Agents Using Liposomes. <i>Angewandte Chemie - International Edition</i> , 2005, 44, 5513-5515.	7.2	185
398	lopamidol: Exploring the potential use of a well-established x-ray contrast agent for MRI. <i>Magnetic Resonance in Medicine</i> , 2005, 53, 830-834.	1.9	115
399	HR-MAS of cells: A cellular water shift due to water-protein interactions?. <i>Magnetic Resonance in Medicine</i> , 2005, 54, 1547-1552.	1.9	19
400	New Insights for Pursuing High Relaxivity MRI Agents from Modelling the Binding Interaction of GdIII Chelates to HSA. <i>ChemBioChem</i> , 2005, 6, 818-820.	1.3	39
401	A Multinuclear NMR Relaxometry Study of Ternary Adducts Formed between Heptadentate GdIII Chelates and L-Lactate. <i>Chemistry - A European Journal</i> , 2005, 11, 5531-5537.	1.7	50
402	Magnetic resonance imaging visualization of targeted cells by the internalization of supramolecular adducts formed between avidin and biotinylated Gd ³⁺ chelates. <i>Journal of Biological Inorganic Chemistry</i> , 2005, 10, 78-86.	1.1	32
403	Visualization through Magnetic Resonance Imaging of DNA Internalized Following <i>In Vivo</i> Electroporation. <i>Molecular Imaging</i> , 2005, 4, 153535002005041.	0.7	5
404	Blast from the Past: The Aluminum's Ghost on the Lanthanum Salts. <i>Current Medicinal Chemistry</i> , 2005, 12, 1631-1636.	1.2	19
405	High Relaxivity Contrast Agents for MRI and Molecular Imaging. , 2005, , 99-121.		9
406	Gd(III)-BASED CONTRAST AGENTS FOR MRI. <i>Advances in Inorganic Chemistry</i> , 2005, 57, 173-237.	0.4	307
407	Efficient relaxivity enhancement in dendritic gadolinium complexes: effective motional coupling in medium molecular weight conjugates. <i>Chemical Communications</i> , 2005, , 474.	2.2	48
408	Dendrimeric Gd(III) complex of a monophosphinated DOTA analogue: optimizing relaxivity by reducing internal motion. <i>Chemical Communications</i> , 2005, , 2390.	2.2	57
409	para-Hydrogenation of unsaturated moieties on poly(lysine) derived substrates for the development of novel hyperpolarized MRI contrast agents. <i>Organic and Biomolecular Chemistry</i> , 2005, 3, 3948.	1.5	19
410	Synthesis of a bifunctional monophosphinic acid DOTA analogue ligand and its lanthanide(III) complexes. A gadolinium(III) complex endowed with an optimal water exchange rate for MRI applications. <i>Organic and Biomolecular Chemistry</i> , 2005, 3, 112-117.	1.5	84
411	Visualization through magnetic resonance imaging of DNA internalized following "in vivo" electroporation. <i>Molecular Imaging</i> , 2005, 4, 7-17.	0.7	9
412	Proteomics as a tool to improve investigation of substantial equivalence in genetically modified organisms: The case of a virus-resistant tomato. <i>Proteomics</i> , 2004, 4, 193-200.	1.3	90
413	Improved route for the visualization of stem cells labeled with a Gd-/Eu-Chelate as dual (MRI and) Tj ETQq1 1 0.784314 rgBT /Overlock 1	1.9	151
414	Towards Targeted MRI: New MRI Contrast Agents for Sialic Acid Detection. <i>Chemistry - A European Journal</i> , 2004, 10, 5205-5217.	1.7	62

#	ARTICLE	IF	CITATIONS
415	Synthesis and characterization of a macrocyclic Schiff base Gd(III) complex as a relaxation agent for a faster acquisition of ² H NMR spectra of ethanol. <i>Inorganica Chimica Acta</i> , 2004, 357, 1374-1380.	1.2	4
416	Synthesis of new polyoxapolycarboxylic ligands for lanthanide(III) ions complexation. <i>Tetrahedron Letters</i> , 2004, 45, 5901-5903.	0.7	6
417	[Gd-AAZTA]-: A New Structural Entry for an Improved Generation of MRI Contrast Agents. <i>Inorganic Chemistry</i> , 2004, 43, 7588-7590.	1.9	217
418	Mannich Reaction as a New Route to Pyridine-Based Polyaminocarboxylic Ligands. <i>Organic Letters</i> , 2004, 6, 1201-1204.	2.4	18
419	Detection and Quantification of Lanthanide Complexes in Cell Lysates by Matrix-Assisted Laser Desorption/Ionization Time-of-Flight Mass Spectrometry. <i>Analytical Chemistry</i> , 2004, 76, 6012-6016.	3.2	13
420	Hetero-Tripodal Hydroxypyridonate Gadolinium Complexes: Syntheses, Relaxometric Properties, Water Exchange Dynamics, and Human Serum Albumin Binding. <i>Inorganic Chemistry</i> , 2004, 43, 8577-8586.	1.9	39
421	Ln(III)-DOTAMGly Complexes: A Versatile Series to Assess the Determinants of the Efficacy of Paramagnetic Chemical Exchange Saturation Transfer Agents for Magnetic Resonance Imaging Applications. <i>Investigative Radiology</i> , 2004, 39, 235-243.	3.5	112
422	Targeting Cells with MR Imaging Probes Based on Paramagnetic Gd(III) Chelates. <i>Current Pharmaceutical Biotechnology</i> , 2004, 5, 509-518.	0.9	97
423	Designing Novel Contrast Agents for Magnetic Resonance Imaging. Synthesis and Relaxometric Characterization of three Gadolinium(III) Complexes Based on Functionalized Pyridine-Containing Macrocyclic Ligands. <i>Helvetica Chimica Acta</i> , 2003, 86, 615-632.	1.0	75
424	Modulation of the Prototropic Exchange Rate at the Water Molecule Coordinated to a Gd(III) Ion. <i>European Journal of Inorganic Chemistry</i> , 2003, 2003, 2045-2048.	1.0	8
425	The Water-Exchange Rate in Neutral Heptadentate DO3A-Like Gd(III) Complexes: Effect of the Basicity at the Macrocyclic Nitrogen Site. <i>European Journal of Inorganic Chemistry</i> , 2003, 2003, 3530-3533.	1.0	25
426	Novel radical-responsive MRI contrast agent based on paramagnetic liposomes. <i>Magnetic Resonance in Chemistry</i> , 2003, 41, 585-588.	1.1	37
427	β-Cyclodextrin adducts of Gd(III) chelates: useful models for investigating the structural and dynamic determinants of the relaxivity of gadolinium-based systems. <i>Magnetic Resonance in Chemistry</i> , 2003, 41, 800-805.	1.1	19
428	Ternary Complexes between Cationic Gd(III) Chelates and Anionic Metabolites in Aqueous Solution: An NMR Relaxometric Study. <i>Chemistry - A European Journal</i> , 2003, 9, 2102-2109.	1.7	87
429	The Structure of the Sugar Residue in Glycated Human Serum Albumin and Its Molecular Recognition by Phenylboronate. <i>Chemistry - A European Journal</i> , 2003, 9, 2193-2199.	1.7	52
430	Supramolecular Adducts between Poly-L-arginine and [Tm(III)DOTP]: A Route to Sensitivity-Enhanced Magnetic Resonance Imaging Contrast Agents. <i>Angewandte Chemie - International Edition</i> , 2003, 42, 4527-4529.	7.2	74
431	Intramolecular hydrogen bonding in transition metal clusters: NMR evidence for an Os ^{IV} -H ₂ S interaction in H ₂ (¹⁷ O) ₃ (CO) ₁₀ (HSR) (R=ethyl, cyclopentyl). <i>Inorganica Chimica Acta</i> , 2003, 351, 251-255.	1.2	5
432	Determination of ferric heme-human serum albumin by ¹ H NMR relaxometry. <i>Journal of Inorganic Biochemistry</i> , 2003, 95, 64-67.	1.5	4

#	ARTICLE	IF	CITATIONS
433	The influence of strong acidic proton donors on the reactivity of H ₂ Ir(CO)Cl(PPh ₃) ₂ with D ₂ . <i>Journal of Molecular Catalysis A</i> , 2003, 204-205, 371-379.	4.8	1
434	Where have all the lanthanum salts gone, long time passing?. <i>Kidney International</i> , 2003, 64, 2327-2328.	2.6	8
435	Enantioselective Recognition between Chiral $\hat{\pm}$ -Hydroxy $\hat{\sim}$ Carboxylates and Macrocyclic Heptadentate Lanthanide(III) Chelates. <i>Inorganic Chemistry</i> , 2003, 42, 4891-4897.	1.9	24
436	Structural Variations Across the Lanthanide Series of Macrocyclic DOTA Complexes: $\hat{\sim}$ Insights into the Design of Contrast Agents for Magnetic Resonance Imaging. <i>Inorganic Chemistry</i> , 2003, 42, 148-157.	1.9	140
437	The Effect of Ligand Scaffold Size on the Stability of Tripodal Hydroxypyridonate Gadolinium Complexes. <i>Inorganic Chemistry</i> , 2003, 42, 2577-2583.	1.9	32
438	In Vitro and In Vivo Study of Solid Lipid Nanoparticles Loaded with Superparamagnetic Iron Oxide. <i>Journal of Drug Targeting</i> , 2003, 11, 19-24.	2.1	100
439	A Highly Stable Gadolinium Complex with a Fast, Associative Mechanism of Water Exchange. <i>Journal of the American Chemical Society</i> , 2003, 125, 14274-14275.	6.6	81
440	Modulation of the antioxidant activity of HO scavengers by albumin binding: a ¹⁹ F-NMR study. <i>Biochemical and Biophysical Research Communications</i> , 2003, 307, 962-966.	1.0	11
441	NMR Structure of Two Novel Polyethylene Glycol Conjugates of the Human Growth Hormone-Releasing Factor, hGRF(1 $\hat{\sim}$ 29) $\hat{\sim}$ NH ₂ . <i>Journal of the American Chemical Society</i> , 2003, 125, 3458-3470.	6.6	42
442	Hyperpolarization transfer from parahydrogen to deuterium via carbon-13. <i>Journal of Chemical Physics</i> , 2003, 119, 8890-8896.	1.2	47
443	Structural and Spectroscopic Study of the Dihydrogen Bond in an Imine Triosmium Complex. <i>Organometallics</i> , 2002, 21, 50-57.	1.1	27
444	NMR Conformational Analysis of Antide, a Potent Antagonist of the Gonadotropin Releasing Hormone. <i>Journal of the American Chemical Society</i> , 2002, 124, 3431-3442.	6.6	13
445	A Paramagnetic MRI-CEST Agent Responsive to Lactate Concentration. <i>Journal of the American Chemical Society</i> , 2002, 124, 9364-9365.	6.6	182
446	Reactivity Studies of Para-Hydrogen with $\hat{\sim}$ ¹³ C-Quinolyl Triosmium Clusters. <i>Organometallics</i> , 2002, 21, 1508-1511.	1.1	23
447	Controlling the variation of axial water exchange rates in macrocyclic lanthanide(III) complexes. Electronic supplementary information (ESI) available: experimental section. See http://www.rsc.org/suppdata/cc/b2/b202862j/ . <i>Chemical Communications</i> , 2002, , 1120-1121.	2.2	69
448	Binding and Relaxometric Properties of Heme Complexes with Cyanogen Bromide Fragments of Human Serum Albumin. <i>Biophysical Journal</i> , 2002, 83, 2248-2258.	0.2	17
449	Novel pH-Reporter MRI Contrast Agents. <i>Angewandte Chemie</i> , 2002, 114, 4510-4512.	1.6	49
450	Molecular Dynamics Simulation of [Gd(egta)(H ₂ O)] $\hat{\sim}$ in Aqueous Solution: Internal Motions of the Poly(amino carboxylate) and Water Ligands, and Rotational Correlation Times. <i>Chemistry - A European Journal</i> , 2002, 8, 1031.	1.7	49

#	ARTICLE	IF	CITATIONS
451	Heterodinuclear LnIII;Na Complexes with an Asymmetric Macrocyclic Compartmental Schiff Base. <i>Chemistry - A European Journal</i> , 2002, 8, 3917-3926.	1.7	21
452	Compartmentalization of a Gadolinium Complex in the Apoferritin Cavity: A Route To Obtain High Relaxivity Contrast Agents for Magnetic Resonance Imaging. <i>Angewandte Chemie - International Edition</i> , 2002, 41, 1017-1019.	7.2	265
453	Novel pH-Reporter MRI Contrast Agents. <i>Angewandte Chemie - International Edition</i> , 2002, 41, 4334-4336.	7.2	198
454	Insights into the use of paramagnetic Gd(III) complexes in MR-molecular imaging investigations. <i>Journal of Magnetic Resonance Imaging</i> , 2002, 16, 394-406.	1.9	340
455	187Os subspectra in 1H and 31P{1H} spectra of triosmium carbonyl clusters. <i>Magnetic Resonance in Chemistry</i> , 2002, 40, 107-113.	1.1	13
456	Coordination equilibrium in an Ln(III) macrocyclic chelate modulated by a reversible interaction with a weakly donor substituent. <i>Magnetic Resonance in Chemistry</i> , 2002, 40, 87-92.	1.1	10
457	1H and 17O relaxometric investigations of the binding of Mn(II) ion to human serum albumin. <i>Magnetic Resonance in Chemistry</i> , 2002, 40, 41-48.	1.1	33
458	Separation of intra- and extracellular lactate NMR signals using a lanthanide shift reagent. <i>Magnetic Resonance in Medicine</i> , 2002, 47, 10-13.	1.9	17
459	Paramagnetic Lanthanide(III) complexes as pH-sensitive chemical exchange saturation transfer (CEST) contrast agents for MRI applications. <i>Magnetic Resonance in Medicine</i> , 2002, 47, 639-648.	1.9	365
460	Innovative magnetic resonance imaging diagnostic agents based on paramagnetic Gd(III) complexes. <i>Biopolymers</i> , 2002, 66, 419-428.	1.2	35
461	Relaxometric evaluation of novel manganese(II) complexes for application as contrast agents in magnetic resonance imaging. <i>Journal of Biological Inorganic Chemistry</i> , 2002, 7, 58-67.	1.1	98
462	Synthesis of the Gd(III) complex with a tetrazole-armed macrocyclic ligand as a potential MRI contrast agent. <i>Tetrahedron Letters</i> , 2002, 43, 783-786.	0.7	20
463	One-step synthesis of a new eight-membered cyclic ligand from glycine, formaldehyde and hypophosphorous acid. <i>Tetrahedron Letters</i> , 2002, 43, 8387-8389.	0.7	14
464	The effect of ligand basicity on the unconventional hydrogen-bond in H(1/4-H)Os3(CO)10L (L=amine) derivatives. <i>Inorganica Chimica Acta</i> , 2002, 334, 448-454.	1.2	4
465	Structural, Luminescence, and NMR Studies of the Reversible Binding of Acetate, Lactate, Citrate, and Selected Amino Acids to Chiral Diaqua Ytterbium, Gadolinium, and Europium Complexes. <i>Journal of the American Chemical Society</i> , 2002, 124, 12697-12705.	6.6	246
466	A paramagnetic probe to localize residues next to carboxylates on protein surfaces. <i>Journal of Biological Inorganic Chemistry</i> , 2002, 7, 617-622.	1.1	24
467	NOE (Nuclear Overhauser Effect) Transfers from para-H2 Enhanced Magnetizations in Alkene Moieties at Rh(I) Centers. <i>Journal of Physical Chemistry A</i> , 2001, 105, 6305-6310.	1.1	25
468	Investigating Pathways of Molecular H2 Exchange in (1/4-H)2Os3(CO)10. <i>Organometallics</i> , 2001, 20, 2924-2927.	1.1	20

#	ARTICLE	IF	CITATIONS
469	Optimization of the Relaxivity of MRI Contrast Agents: Effect of Poly(ethylene glycol) Chains on the Water-Exchange Rates of GdIII Complexes. <i>Journal of the American Chemical Society</i> , 2001, 123, 10758-10759.	6.6	87
470	pH-Dependent Modulation of Relaxivity and Luminescence in Macrocyclic Gadolinium and Europium Complexes Based on Reversible Intramolecular Sulfonamide Ligation. <i>Journal of the American Chemical Society</i> , 2001, 123, 7601-7609.	6.6	269
471	Modulation of the water exchange rates in [Gd(III)DO3A] complex by formation of ternary complexes with carboxylate ligands. <i>Chemical Communications</i> , 2001, , 115-116.	2.2	49
472	Synthesis and Characterization of Triosmium Clusters Containing the Bidentate Ligand Ph ₂ PCH ₂ CH ₂ SMe: Detection of an Isomerization Reaction Involving Bridging and Chelating Ligand Coordination Modes. <i>Organometallics</i> , 2001, 20, 4150-4160.	1.1	21
473	6-Carboxamido-5,4-Hydroxypyrimidinones: A New Class of Heterocyclic Ligands and Their Evaluation as Gadolinium Chelating Agents. <i>Inorganic Chemistry</i> , 2001, 40, 6746-6756.	1.9	44
474	Effect of ibuprofen and warfarin on the allosteric properties of haem-human serum albumin. <i>FEBS Journal</i> , 2001, 268, 6214-6220.	0.2	123
475	Slow clearance gadolinium-based extracellular and intravascular contrast media for three-dimensional MR angiography. <i>Journal of Magnetic Resonance Imaging</i> , 2001, 13, 588-593.	1.9	13
476	{DOTA-bis(amide)}lanthanide Complexes: NMR Evidence for Differences in Water-Molecule Exchange Rates for Coordination Isomers. <i>Chemistry - A European Journal</i> , 2001, 7, 288-296.	1.7	65
477	High-Relaxivity Contrast Agents for Magnetic Resonance Imaging Based on Multisite Interactions between α -2-Cyclodextrin Oligomer and Suitably Functionalized GdIII Chelates. <i>Chemistry - A European Journal</i> , 2001, 7, 5261-5269.	1.7	69
478	A Calix[4]arene GdIII Complex Endowed with High Stability, Relaxivity, and Binding Affinity to Serum Albumin. <i>Angewandte Chemie - International Edition</i> , 2001, 40, 4737-4739.	7.2	41
479	Ring-opening polymerisation of norbornene by (1/4-H)2Os3(CO)10 complex. <i>Journal of Organometallic Chemistry</i> , 2001, 622, 43-46.	0.8	14
480	Relaxometric characterization of human hemalbumin. <i>Journal of Biological Inorganic Chemistry</i> , 2001, 6, 650-658.	1.1	33
481	NMR Relaxation Studies of Polynuclear Hydrides Derivatives. , 2001, , 351-374.		2
482	Ap(O ₂)-Responsive MRI Contrast Agent Based on the Redox Switch of Manganese(II /III) - Porphyrin Complexes. <i>Angewandte Chemie - International Edition</i> , 2000, 39, 747-750.	7.2	150
483	Isolation and ¹³ C-NMR characterization of an insoluble proteinaceous fraction from substantia nigra of patients with parkinson's disease. <i>Movement Disorders</i> , 2000, 15, 977-981.	2.2	42
484	Non-covalent Conjugates between Cationic Polyamino Acids and GdIII Chelates: A Route for Seeking Accumulation of MRI-Contrast Agents at Tumor Targeting Sites. <i>Chemistry - A European Journal</i> , 2000, 6, 2609-2617.	1.7	69
485	Facile oxidative addition/reductive elimination of HX (X=Cl, Br, I) on (1/4-H)2Os3(CO)10. <i>Journal of Organometallic Chemistry</i> , 2000, 593-594, 135-141.	0.8	4
486	Nuclear magnetic relaxation dispersion profiles of substantia nigra pars compacta in Parkinson's disease patients are consistent with protein aggregation. <i>Neurochemistry International</i> , 2000, 37, 331-336.	1.9	31

#	ARTICLE	IF	CITATIONS
487	The Selectivity of Reversible Oxy-Anion Binding in Aqueous Solution at a Chiral Europium and Terbium Center: A Signaling of Carbonate Chelation by Changes in the Form and Circular Polarization of Luminescence Emission. <i>Journal of the American Chemical Society</i> , 2000, 122, 9674-9684.	6.6	292
488	Hetero-dinuclear sodium-lanthanide(III) complexes with an asymmetric compartmental macrocycle. <i>Chemical Communications</i> , 2000, , 145-146.	2.2	13
489	Ternary Gd(III)-L-HSA adducts: evidence for the replacement of inner-sphere water molecules by coordinating groups of the protein. Implications for the design of contrast agents for MRI. <i>Journal of Biological Inorganic Chemistry</i> , 2000, 5, 488-497.	1.1	140
490	Correlation of Water Exchange Rate with Isomeric Composition in Diastereoisomeric Gadolinium Complexes of Tetra(carboxyethyl)dota and Related Macrocyclic Ligands. <i>Journal of the American Chemical Society</i> , 2000, 122, 9781-9792.	6.6	189
491	Conjugates of cyclodextrins with charged and neutral macrocyclic europium, terbium and gadolinium complexes: sensitised luminescence and relaxometric investigations and an example of supramolecular relaxivity enhancement. <i>Perkin Transactions II RSC</i> , 2000, , 1329-1338.	1.1	41
492	Enhanced mobility of the cluster Ru ₃ (CO) ₁₂ in the solid state formed in situ by the reaction of CO and Ru ₃ (CO) ₆ (1/4-CO)(1/4-3-2-3-5-C12H8) on silica. <i>Chemical Communications</i> , 2000, , 1425-1426.	2.2	4
493	Multinuclear and multifrequency NMR study of gadolinium(III) complexes with bis-amide derivatives of ethylenedioxydiethylenedinitrilotetraacetic acid. <i>Dalton Transactions RSC</i> , 2000, , 3435-3440.	2.3	12
494	The role of molecular shape and axial symmetry in the solid state molecular dynamics of metal carbonyl compounds included in cyclodextrin cavities. <i>Dalton Transactions RSC</i> , 2000, , 4075-4077.	2.3	10
495	Syntheses and Relaxation Properties of Mixed Gadolinium Hydroxypyridinonate MRI Contrast Agents. <i>Inorganic Chemistry</i> , 2000, 39, 5747-5756.	1.9	95
496	An NMR Study of H(1/4-H)Os ₃ (CO) ₁₁ . <i>Inorganic Chemistry</i> , 2000, 39, 2422-2425.	1.9	9
497	First 17O NMR Observation of Coordinated Water on Both Isomers of [Eu(DOTAM)(H ₂ O)] ₃ : A Direct Access to Water Exchange and its Role in the Isomerization. <i>Journal of the American Chemical Society</i> , 2000, 122, 1506-1512.	6.6	163
498	A Tris-hydroxymethyl-Substituted Derivative of Gd-TREN-Me-3,2-HOPO: An MRI Relaxation Agent with Improved Efficiency. <i>Journal of the American Chemical Society</i> , 2000, 122, 11228-11229.	6.6	83
499	[GdPCP2A(H ₂ O) ₂]: A Paramagnetic Contrast Agent Designed for Improved Applications in Magnetic Resonance Imaging. <i>Journal of Medicinal Chemistry</i> , 2000, 43, 4017-4024.	2.9	86
500	A novel non-facial mode of azulene cluster co-ordination. Crystal and molecular structure of the dicarbenoid Os ₃ (1/4-H) ₂ (CO) ₉ (1/4-3-1-1-1-1-C10H6). <i>Dalton Transactions RSC</i> , 2000, , 2215-2217.	2.3	9
501	Coordination of Imines on Os ₃ Clusters: Effect of the Solvent in Addressing Isomer Formation. <i>Organometallics</i> , 2000, 19, 707-710.	1.1	11
502	Chemical Shift and Relaxation Reagents in NMR. , 1999, , 223-231.		0
503	NMR, Relaxometric, and Structural Studies of the Hydration and Exchange Dynamics of Cationic Lanthanide Complexes of Macrocyclic Tetraamide Ligands. <i>Journal of the American Chemical Society</i> , 1999, 121, 5762-5771.	6.6	267
504	A holmium complex of a macrocyclic ligand (DOTA) and its isostructural europium analogue. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1999, 55, 353-356.	0.4	37

#	ARTICLE	IF	CITATIONS
505	¹ H and ¹⁷ O-NMR relaxometric investigations of paramagnetic contrast agents for MRI. Clues for higher relaxivities. <i>Coordination Chemistry Reviews</i> , 1999, 185-186, 321-333.	9.5	48
506	Contrast agents for magnetic resonance angiographic applications: ¹ H and ¹⁷ O NMR relaxometric investigations on two gadolinium(III) DTPA-like chelates endowed with high binding affinity to human serum albumin. <i>Journal of Biological Inorganic Chemistry</i> , 1999, 4, 766-774.	1.1	99
507	A novel ¹⁹ F-NMR method for the investigation of the antioxidant capacity of biomolecules and biofluids. <i>Free Radical Biology and Medicine</i> , 1999, 27, 356-363.	1.3	4
508	Contrast Agents for Magnetic Resonance Imaging: A Novel Route to Enhanced Relaxivities Based on the Interaction of a Gd(III) Chelate with Poly- β -cyclodextrins. <i>Chemistry - A European Journal</i> , 1999, 5, 1253-1260.	1.7	45
509	Spectral Discrimination of Chiral Macrocyclic Paramagnetic Metal Complexes by NMR Techniques. <i>Chemistry - A European Journal</i> , 1999, 5, 1261-1266.	1.7	20
510	Novel Paramagnetic Macromolecular Complexes Derived from the Linkage of a Macrocyclic Gd(III) Complex to Polyamino Acids through a Squaric Acid Moiety. <i>Bioconjugate Chemistry</i> , 1999, 10, 192-199.	1.8	66
511	A macromolecular Gd(III) complex as pH-responsive relaxometric probe for MRI applications. <i>Chemical Communications</i> , 1999, , 1577-1578.	2.2	83
512	Structure and relaxivity of macrocyclic gadolinium complexes incorporating pyridyl and 4-morpholinopyridyl substituents. <i>New Journal of Chemistry</i> , 1999, 23, 669.	1.4	41
513	Fast carbonyl exchange in the solid state for β -cyclodextrin/(arene)Cr(CO) ₃ inclusion complexes. <i>Chemical Communications</i> , 1999, , 281-282.	2.2	9
514	Synthesis, X-ray Structure, and Solution NMR Studies of Ln(III) Complexes with a Macrocyclic Asymmetric Compartmental Schiff Base. Preference of the Ln(III) Ions for a Crown-Like Coordination Site. <i>Inorganic Chemistry</i> , 1999, 38, 2906-2916.	1.9	44
515	Chemical Shift and Relaxation Reagents in NMR*. , 1999, , 253-261.		0
516	Dependence of the relaxivity and luminescence of gadolinium and europium amino-acid complexes on hydrogencarbonate and pH. <i>Chemical Communications</i> , 1999, , 1047-1048.	2.2	71
517	Prototropic and Water-Exchange Processes in Aqueous Solutions of Gd(III) Chelates. <i>Accounts of Chemical Research</i> , 1999, 32, 941-949.	7.6	198
518	Metal Complexes as Allosteric Effectors of Human Hemoglobin: An NMR Study of the Interaction of the Gadolinium(III) Bis(m-boroxypheylamide)diethylenetriaminepentaacetic Acid Complex with Human Oxygenated and Deoxygenated Hemoglobin. <i>Biophysical Journal</i> , 1999, 76, 2735-2743.	0.2	18
519	Coordination of an Imine Ligand on an Os ₃ Cluster Stabilized by Intramolecular Os ^{IV} H--H ^{IV} N Hydrogen Bonding. <i>Organometallics</i> , 1999, 18, 2030-2032.	1.1	28
520	NOVEL PARAMAGNETIC MACROMOLECULAR COMPLEXES DERIVED FROM THE LINKAGE OF A MACROCYCLIC Gd(III) COMPLEX TO POLYAMINO ACIDS THROUGH A SQUARIC ACID MOIETY. <i>Bioconjugate Chemistry</i> , 1999, 10, 701-701.	1.8	28
521	A Novel Application of para H ₂ : The Reversible Addition/Elimination of H ₂ at a Ru ₃ Cluster Revealed by the Enhanced NMR Emission Resonance from Molecular Hydrogen. <i>Journal of Physical Chemistry A</i> , 1999, 103, 9702-9705.	1.1	40
522	Contrast Agents for Magnetic Resonance Imaging: A Novel Route to Enhanced Relaxivities Based on the Interaction of a Gd(III) Chelate with Poly- β -cyclodextrins. , 1999, 5, 1253.		1

#	ARTICLE	IF	CITATIONS
523	NMR characterization of four new isomers of $H(\frac{1}{4}H)Os_3(CO)_{10}$ (phosphine). <i>Inorganica Chimica Acta</i> , 1998, 275-276, 521-527.	1.2	14
524	NMR and luminescence studies on the formation of ternary adducts between HSA and $Ln(III)$ malonate complexes (Ln=Eu, Gd, Tb). <i>BBA - Proteins and Proteomics</i> , 1998, 1385, 7-16.	2.1	6
525	Determination of the Prototropic Exchange Rate at the Water Molecule Coordinated to an Anionic Paramagnetic Gd(III) Chelate. <i>European Journal of Inorganic Chemistry</i> , 1998, 1283-1289.	1.0	17
526	Direct NMR Spectroscopic Observation of a Lanthanide-Coordinated Water Molecule whose Exchange Rate Is Dependent on the Conformation of the Complexes. <i>Angewandte Chemie - International Edition</i> , 1998, 37, 2673-2675.	7.2	133
527	NMR relaxometric studies of Gd(III) complexes with heptadentate macrocyclic ligands. <i>Magnetic Resonance in Chemistry</i> , 1998, 36, S200-S208.	1.1	129
528	Stabilization of the T-state of ferrous human adult and fetal hemoglobin by $Ln(III)$ complexes: A thermodynamic study. <i>Journal of Inorganic Biochemistry</i> , 1998, 71, 37-43.	1.5	15
529	NMR relaxometric investigations of solid lipid nanoparticles (SLN) containing gadolinium(III) complexes. <i>European Journal of Pharmaceutics and Biopharmaceutics</i> , 1998, 45, 157-163.	2.0	88
530	Detection of a novel intermediate in the addition of thiols to osmium carbonyl clusters. <i>Chemical Communications</i> , 1998, , 2721-2722.	2.2	15
531	Relaxometric and luminescence behaviour of triaquahexaazamacrocyclic complexes, the gadolinium complex displaying a high relaxivity with a pronounced pH dependence. <i>New Journal of Chemistry</i> , 1998, 22, 627-631.	1.4	49
532	Synthesis, NMR, relaxometry and circularly polarised luminescence studies of macrocyclic monoamidetris(phosphinate) complexes bearing a remote chiral centre. <i>Journal of the Chemical Society Dalton Transactions</i> , 1998, , 881-892.	1.1	32
533	Synthesis, crystal structure and solid-state dynamics of the $La(hfa)_3 \cdot Me(OCH_2CH_2)_4OMe$ ($Hhfa = 1,1,1,5,5,5$ -hexafluoropentane-2,4-dione) precursor for MOCVD applications. <i>Journal of the Chemical Society Dalton Transactions</i> , 1998, , 1509-1512.	1.1	19
534	Addition of NH_3 to the Cluster Complexes $M_3H(\frac{1}{4}H)(CO)_{11}$ (M = Ru, Os). <i>Organometallics</i> , 1998, 17, 5353-5357.	1.1	5
535	Novel Synthesis, Solution Structure, Dynamics, and Protonation of $H(\frac{1}{4}H)Ru_3(CO)_{11}$. <i>Organometallics</i> , 1998, 17, 3182-3185.	1.1	16
536	Longitudinal Nuclear Relaxation in an A2Spin System Initially Polarized through Para-Hydrogen. <i>Journal of the American Chemical Society</i> , 1998, 120, 6770-6773.	6.6	61
537	Lanthanide(III) chelates for NMR biomedical applications. <i>Chemical Society Reviews</i> , 1998, 27, 19-29.	18.7	698
538	Role of $Os \sim H \sim N$ Interactions in Directing the Stereochemistry of Carbonyl Cluster Hydride Derivatives. <i>Organometallics</i> , 1997, 16, 5140-5141.	1.1	45
539	A versatile route to $[Ru_3(CO)_{11}(alkyne)]$ complexes. <i>Chemical Communications</i> , 1997, , 267-268.	2.2	6
540	Nuclear magnetic resonance, luminescence and structural studies of lanthanide complexes with octadentate macrocyclic ligands bearing benzylphosphinate groups. <i>Journal of the Chemical Society Dalton Transactions</i> , 1997, , 3623-3636.	1.1	82

#	ARTICLE	IF	CITATIONS
541	A Multinuclear NMR Study on the Structure and Dynamics of Lanthanide(III) Complexes of the Poly(amino carboxylate) EGTA4-in Aqueous Solution. <i>Inorganic Chemistry</i> , 1997, 36, 5104-5112.	1.9	74
542	A Novel Compound in the Lanthanide(III) DOTA Series. X-ray Crystal and Molecular Structure of the Complex Na[La(DOTA)La(HDOTA)]·10H ₂ O. <i>Inorganic Chemistry</i> , 1997, 36, 4287-4289.	1.9	87
543	Conformational and Coordination Equilibria on DOTA Complexes of Lanthanide Metal Ions in Aqueous Solution Studied by ¹ H-NMR Spectroscopy. <i>Inorganic Chemistry</i> , 1997, 36, 2059-2068.	1.9	333
544	PrototropicvsWhole Water Exchange Contributions to the Solvent Relaxation Enhancement in the Aqueous Solution of a Cationic Gd ³⁺ -Macrocyclic Complex. <i>Journal of the American Chemical Society</i> , 1997, 119, 4767-4768.	6.6	108
545	Synthesis and NMR Studies of Three Pyridine-Containing Triaza Macrocyclic Triacetate Ligands and Their Complexes with Lanthanide Ions. <i>Inorganic Chemistry</i> , 1997, 36, 2992-3000.	1.9	119
546	EPR investigations of the iron domain in neuromelanin. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 1997, 1361, 49-58.	1.8	52
547	Non-ionic Ln(III) chelates as MRI contrast agents: Synthesis, characterisation and ¹ H NMR relaxometric investigations of bis(benzylamide)diethylenetriaminepentaacetic acid Lu(III) and Gd(III) complexes. <i>Inorganica Chimica Acta</i> , 1997, 254, 63-70.	1.2	63
548	Towards MRI contrast agents of improved efficacy. NMR relaxometric investigations of the binding interaction to HSA of a novel heptadentate macrocyclic triphosphonate Gd(III)-complex. <i>Journal of Biological Inorganic Chemistry</i> , 1997, 2, 470-479.	1.1	77
549	Relaxometric Determination of the Exchange Rate of the Coordinated Water Protons in a Neutral Gd ^{III} Chelate. <i>Chemistry - A European Journal</i> , 1997, 3, 1499-1504.	1.7	29
550	Relaxometric, Structural, and Dynamic NMR Studies of DOTA-like Ln(III) Complexes (Ln = La, Gd, Ho, Yb) Containing ap-Nitrophenyl Substituent. <i>Inorganic Chemistry</i> , 1996, 35, 2726-2736.	1.9	77
551	Structural determinants of fluoride and formate binding to hemoglobin and myoglobin: crystallographic and ¹ H-NMR relaxometric study. <i>Biophysical Journal</i> , 1996, 70, 482-488.	0.2	49
552	H/D Isotopic Exchange on the Surface of a Triosmium Carbonyl Cluster. <i>Organometallics</i> , 1996, 15, 4967-4970.	1.1	10
553	Relationship between Solid State NMR Parameters and X-ray Structural Data in Tricadmium Phosphates. <i>Inorganic Chemistry</i> , 1996, 35, 149-154.	1.9	16
554	Extent of hydration of octadentate lanthanide complexes incorporating phosphinate donors: solution relaxometry and luminescence studies. <i>Journal of the Chemical Society Dalton Transactions</i> , 1996, , 17.	1.1	69
555	Paramagnetic complexes as novel NMR pH indicators. <i>Chemical Communications</i> , 1996, , 1265.	2.2	17
556	An NMR relaxometric indicator of the formation of OH [•] radicals in fenton-type reactions. <i>Chemical Communications</i> , 1996, , 1509.	2.2	4
557	¹ H-NMR relaxometric study of pancreatic serine (pro)enzyme inhibition by a Gd(III) chelate bearing boronic functionalities. <i>IUBMB Life</i> , 1996, 39, 741-746.	1.5	3
558	Chapter 8 considerations on high resolution solid state NMR in paramagnetic molecules. <i>Coordination Chemistry Reviews</i> , 1996, 150, 221-242.	9.5	24

#	ARTICLE	IF	CITATIONS
559	Gd(III) complexes as contrast agents for magnetic resonance imaging: a proton relaxation enhancement study of the interaction with human serum albumin. <i>Journal of Biological Inorganic Chemistry</i> , 1996, 1, 312-319.	1.1	167
560	A new ytterbium chelate as contrast agent in chemical shift imaging and temperature sensitive probe for MR spectroscopy. <i>Magnetic Resonance in Medicine</i> , 1996, 35, 648-651.	1.9	83
561	Crystal structure and solution dynamics of the lutetium(III) chelate of DOTA. <i>Inorganica Chimica Acta</i> , 1996, 246, 423-429.	1.2	141
562	Azide, cyanide, fluoride, imidazole and pyridine binding to ferric and ferrous native horse heart cytochrome c and to its carboxymethylated derivative: A comparative study. <i>Journal of Inorganic Biochemistry</i> , 1996, 62, 213-222.	1.5	42
563	Nuclear magnetic resonance spectroscopy characterization and iron content determination of human mesencephalic neuromelanin. <i>Advances in Neurology</i> , 1996, 69, 263-70.	0.8	5
564	NMR Relaxometric Investigation on Human Methemoglobin and Fluoromethemoglobin. An Improved Quantitative in Vitro Assay of Human Methemoglobin. <i>Magnetic Resonance in Medicine</i> , 1995, 33, 827-831.	1.9	14
565	Dipolar Oscillations in the ^{13}C { 1 H } Cross-Polarization/Magic-Angle-Spinning Spectrum of $\text{H}_2\text{Os}_3(\text{CO})_{10}$. <i>Journal of Magnetic Resonance Series A</i> , 1995, 116, 251-254.	1.6	6
566	Solution structures and dynamics of $\text{M}_3(\text{CO})_{12}\text{L}_x(\text{NCCH}_3)_x$ ($\text{M} = \text{Ru}$, $x = 1, 2, 3$; $\text{M} = \text{Os}$, $x = 1, 2$). <i>Inorganica Chimica Acta</i> , 1995, 235, 357-366.	1.2	18
567	Solid-Gas Reactions of the "Lightly Stabilized" $\text{Os}_3(\text{CO})_{11}\text{L}$ ($\text{L} = \text{NCCH}_3, \text{C}_2\text{H}_4$) Clusters with CO , NH_3 , and H_2 . <i>Organometallics</i> , 1995, 14, 3224-3228.	1.1	21
568	NMR Evidence for Interconversion between Two Enantiomeric Forms of Macrocyclic Schiff Base Lanthanide(III) Complexes Through Reversible Ring Contraction and Expansion. <i>Inorganic Chemistry</i> , 1995, 34, 5825-5831.	1.9	51
569	Evaluation of the energy barrier for carbonyl exchange in the highly fluxional $\text{Ru}_3(\text{CO})_{12}$ system. <i>Organometallics</i> , 1995, 14, 4435-4438.	1.1	14
570	^1H MAS NMR Spectra of Transition Metal Carbonyl Hydrides. <i>Inorganic Chemistry</i> , 1995, 34, 3557-3559.	1.9	4
571	Solution Structures and Dynamics of $\text{Ru}_4(\mu\text{-H})_4(\text{CO})_{12}\text{L}_x$ ($x = 1-4$, $\text{L} = \text{P}(\text{OEt})_3, \text{PPh}_3$; $x = 1-2$, $\text{L} = \text{Tj ETQq1}$) $\text{Ru}_4(\mu\text{-H})_4(\text{CO})_{10}(\text{P}(\text{OEt})_3)_2$. <i>Organometallics</i> , 1995, 14, 3693-3703.	1.1	22
572	Molecular Recognition of R- and T-States of Human Adult Hemoglobin by a Paramagnetic Gd(III) Complex by Means of the Measurement of Solvent Water Proton Relaxation Rate. <i>Journal of the American Chemical Society</i> , 1995, 117, 9365-9366.	6.6	39
573	NMR conformational study of the lanthanide(III) complexes of DOTA in aqueous solution. <i>Journal of Alloys and Compounds</i> , 1995, 225, 303-307.	2.8	56
574	High-resolution NMR and relaxometric studies of Ln(III) complexes of relevance to MRI. <i>Journal of Alloys and Compounds</i> , 1995, 225, 274-278.	2.8	5
575	Novel Contrast Agents for Magnetic Resonance Imaging. Synthesis and Characterization of the Ligand BOPTA and Its Ln(III) Complexes ($\text{Ln} = \text{Gd}, \text{La}, \text{Lu}$). X-ray Structure of Disodium (TPS-9-145337286-C-S)-[4-Carboxy-5,8,11-tris(carboxymethyl)-1-phenyl-2-oxa-5,8,11-triazatridecan-13-oato(5-)]gadolate(2-) in a Mixture with Its Enantiomer. <i>Inorganic Chemistry</i> , 1995, 34, 633-643.	1.9	180
576	Nuclear magnetic resonance studies of neutral lanthanide(III) complexes with tetraaza-macrocyclic ligands containing three phosphinate and one carboxamide co-ordinating arms. <i>Journal of the Chemical Society Dalton Transactions</i> , 1995, , 2259.	1.1	43

#	ARTICLE	IF	CITATIONS
577	MRI Contrast agents: macrocyclic lanthanide(III) complexes with improved relaxation efficiency. <i>Journal of the Chemical Society Chemical Communications</i> , 1995, , 1885.	2.0	38
578	Solution and solid-state NMR studies of MeHgII and RSHgII (R = Me, Et) complexes with pyrazolyl-containing ligands. <i>Polyhedron</i> , 1994, 13, 2695-2702.	1.0	9
579	Synthesis and VT-NMR studies of cationic hydride platinum clusters. <i>Journal of Cluster Science</i> , 1994, 5, 523-533.	1.7	2
580	Stereochemical Nonrigidity in $(\mu_2\text{-H})(\text{H})\text{Os}_3(\text{CO})_{11}$. Evidence for the Occurrence of a Racemization Process at Low Temperature. <i>Organometallics</i> , 1994, 13, 3737-3739.	1.1	12
581	Solution and Solid-State Characterization of Highly Rigid, Eight-Coordinate Lanthanide(III) Complexes of a Macrocyclic Tetrabenzylphosphinate. <i>Inorganic Chemistry</i> , 1994, 33, 4696-4706.	1.9	152
582	^1H and ^{13}C NMR Study of the Solid-State Dynamics of $\text{H}_4\text{Ru}_4(\text{CO})_{12}$. <i>Organometallics</i> , 1994, 13, 2375-2379.	1.1	19
583	Reactions of Solid $\text{H}_2\text{Os}_3(\text{CO})_{10}$ with Gaseous CO , NH_3 , and H_2S . <i>Organometallics</i> , 1994, 13, 4232-4236.	1.1	17
584	NMR Evidence of a Long Exchange Lifetime for the Coordinated Water in Ln(III)-Bis(methyl amide)-DTPA Complexes (Ln = Gd, Dy). <i>Inorganic Chemistry</i> , 1994, 33, 4707-4711.	1.9	95
585	Evidence for a Glycidic-Lipidic Matrix in Human Neuromelanin, Potentially Responsible for the Enhanced Iron Sequestering Ability of Substantia Nigra. <i>Journal of Neurochemistry</i> , 1994, 62, 369-371.	2.1	40
586	NMR relaxometric studies of water accessibility to haem cavity in horse heart and sperm whale myoglobin. <i>Magnetic Resonance in Chemistry</i> , 1993, 31, S85-S89.	1.1	9
587	Gd(DOTP) $_5$ - outer-sphere relaxation enhancement promoted by nitrogen bases. <i>Magnetic Resonance in Medicine</i> , 1993, 30, 583-591.	1.9	71
588	High-resolution NMR studies of (diphosphine)(diene) rhodium complexes in the solid state. <i>Journal of Organometallic Chemistry</i> , 1993, 463, 223-226.	0.8	4
589	Solid-state ^{31}P NMR vs. solution study of bis(tertiary phosphines). <i>Solid State Nuclear Magnetic Resonance</i> , 1993, 2, 245-251.	1.5	3
590	^1H and ^{13}C NMR study of cyclopentadienyl metal carbonyls in the solid state. <i>Spectrochimica Acta Part A: Molecular Spectroscopy</i> , 1993, 49, 1307-1314.	0.1	3
591	Solid state ^{13}C NMR spectra of metal carbonyl clusters. <i>Journal of Cluster Science</i> , 1993, 4, 1-8.	1.7	8
592	Paramagnetic GdIII–FeIII heterobimetallic complexes of DTPA-bis-salicylamide. <i>Spectrochimica Acta Part A: Molecular Spectroscopy</i> , 1993, 49, 1315-1322.	0.1	29
593	Crystal structure and NMR investigation of the serine proteinase inhibitor MR889, a cyclic thiolic compound. <i>Journal of the Chemical Society Perkin Transactions II</i> , 1993, , 2253.	0.9	5
594	Solid-gas reactions of molecular organometallic complexes. Detection of a metastable intermediate in the carbonylation of a nickel(0) complex. <i>Organometallics</i> , 1993, 12, 4757-4763.	1.1	11

#	ARTICLE	IF	CITATIONS
595	Synthesis and characterization of a novel DTPA-like gadolinium(III) complex: a potential reagent for the determination of glycosylated proteins by water proton NMR relaxation measurements. <i>Inorganic Chemistry</i> , 1993, 32, 2068-2071.	1.9	35
596	Synthesis, molecular structure, crystal packing, and dynamic behaviour in the solid state of $[\text{Fe}_2(\mu\text{-C}_5\text{H}_5)_2(\mu\text{-CO})(\text{CO})_2\{\mu\text{-CR}(\text{CN})\}](\text{R} = \text{H or CN})$. <i>Journal of the Chemical Society Dalton Transactions</i> , 1992, , 2961-2966.	1.1	10
597	Determination of metal-proton distances and electronic relaxation times in lanthanide complexes by nuclear magnetic resonance spectroscopy. <i>Journal of the Chemical Society Dalton Transactions</i> , 1992, , 225-228.	1.1	54
598	NMR study of solution structures and dynamics of lanthanide(III) complexes of DOTA. <i>Inorganic Chemistry</i> , 1992, 31, 4291-4299.	1.9	323
599	Magnetic resonance imaging of the gastrointestinal tract: Investigation of baby milk as a low cost contrast medium. <i>European Journal of Radiology</i> , 1992, 15, 171-174.	1.2	12
600	Synthesis and NMRD studies of gadolinium(3+) complexes of macrocyclic polyamino polycarboxylic ligands bearing .beta.-benzyloxy-.alpha.-propionic residues. <i>Inorganic Chemistry</i> , 1992, 31, 1100-1103.	1.9	62
601	Synthesis, characterization, and 1/T1 NMRD profiles of gadolinium(III) complexes of monoamide derivatives of DOTA-like ligands. X-ray structure of the 10-[2-[[2-hydroxy-1-(hydroxymethyl)ethyl]amino]-1-[(phenylmethoxy)methyl]-2-oxoethyl]-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid-gadolinium(III) complex. <i>Inorganic Chemistry</i> , 1992, 31, 2422-2428.	1.9	122
602	Reorientational motions of permethylated cyclopentadienyl rings in polycrystalline organometallic compounds. <i>Inorganic Chemistry</i> , 1992, 31, 3054-3059.	1.9	12
603	Quantitative Determination of Methemoglobin by Measuring the Solvent-Water Proton-Nuclear Magnetic Resonance Relaxation Rate. <i>Clinical Chemistry</i> , 1992, 38, 2401-2404.	1.5	6
604	Paramagnetic water proton relaxation enhancement: From contrast agents in MRI to reagents for quantitative <i>in vitro</i> assays. <i>Magnetic Resonance Imaging</i> , 1992, 10, 849-854.	1.0	16
605	Metal ion content in <i>Sepia officinalis</i> melanin. <i>Marine Chemistry</i> , 1992, 39, 243-250.	0.9	20
606	Use of solid-state carbon-13 NMR spectroscopy to quantify the degree of asymmetry of bonding for semibridging carbonyl groups in iron carbonyl complexes. <i>Inorganic Chemistry</i> , 1991, 30, 1489-1493.	1.9	23
607	NMR study of the chemical exchange of the hydrido ligands in the bis(bis(diphenylphosphino)ethane)trihydridodiplatinum(II) cation by T1, T2, and HD isotopic perturbation. <i>Inorganic Chemistry</i> , 1991, 30, 316-318.	1.9	12
608	Proton spin-lattice NMR relaxation studies of hydride carbonyl clusters: a method to evaluate distances involving hydrido ligands. <i>Inorganic Chemistry</i> , 1991, 30, 1614-1617.	1.9	15
609	Synthesis, reactivity, and ligand dynamics of $\text{Ru}_3(\text{CO})_9(\mu\text{-CO})(\mu\text{-}\eta\text{-2-alkyne})$ compounds. <i>Organometallics</i> , 1991, 10, 2854-2856.	1.1	41
610	Dynamic processes in the solid state. Proton relaxation studies and potential energy barrier calculations for (arene) $\text{M}(\text{CO})_3$ species. X-ray crystal structures of $(1,2,3\text{-C}_6\text{H}_3\text{Me}_3)\text{Cr}(\text{CO})_3$ and $(1,2,4,5\text{-C}_6\text{H}_2\text{Me}_4)\text{Cr}(\text{CO})_3$. <i>Inorganic Chemistry</i> , 1991, 30, 951-956.	1.9	24
611	Trends in NMR studies of paramagnetic Gd(III) complexes as potential contrast agents in MRI. <i>Magnetic Resonance Imaging</i> , 1991, 9, 843-847.	1.0	22
612	An NMR study of the interaction between Melanin Free Acid and Mn^{2+} ions as a model to mimic the enhanced proton relaxation rates in melanotic melanoma. <i>Magnetic Resonance Imaging</i> , 1991, 9, 963-968.	1.0	9

#	ARTICLE	IF	CITATIONS
613	An NMR relaxation study of aqueous solutions of Gd(III) chelates. <i>Journal of Magnetic Resonance</i> , 1991, 92, 572-580.	0.5	3
614	NMR Studies of Melanins: Characterization of a Soluble Melanin Free Acid From Sepia Ink. <i>Pigment Cell & Melanoma Research</i> , 1991, 4, 216-221.	4.0	28
615	Inclusion complexes between β -cyclodextrin and β -benzyloxy- β -propionic derivatives of paramagnetic DOTA- and DPTA-Gd(III) complexes. <i>Magnetic Resonance in Chemistry</i> , 1991, 29, 923-927.	1.1	58
616	Water signal suppression by T2-relaxation enhancement promoted by Dy(III) complexes. <i>Magnetic Resonance in Chemistry</i> , 1991, 29, S85-S88.	1.1	21
617	NMR studies of the reorientational motions of cyclopentadienyl ligands in solidcis- and trans- $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Fe}_2(\text{CO})_2(\eta^4\text{-CO})_2]$. <i>Magnetic Resonance in Chemistry</i> , 1990, 28, S52-S58.	1.1	13
618	Solution structure and dynamics of DTPA-Ln(III) complexes (DTPA=diethylene triamine penta acetate); Tj ETQq0 0 0 rgBT /Overlock 10 T	1.2	42
619	Complexes of diethylenetriaminepentaacetic acid as contrast agents in NMR imaging. Computer simulation of equilibria in human blood plasma. <i>Analytica Chimica Acta</i> , 1990, 233, 85-100.	2.6	7
620	Solid state ^{13}C CP-MAS NMR spectra of iron (alkyne) carbonyl compounds. <i>Journal of Organometallic Chemistry</i> , 1989, 368, 331-338.	0.8	6
621	NMR studies of L-Dopa melanin-manganese(II) complex in water solution. <i>Journal of Inorganic Biochemistry</i> , 1989, 36, 1-9.	1.5	6
622	Relationships between structure and ligand dynamics. <i>Journal of Organometallic Chemistry</i> , 1989, 365, 163-185.	0.8	46
623	High-field magic-angle-spinning carbon-13 NMR spectroscopy of tetracobalt dodecacarbonyl. <i>Inorganic Chemistry</i> , 1989, 28, 1196-1198.	1.9	22
624	Solution structure and dynamic behaviour of two isomers of $[\text{Fe}_3(\text{CO})_9\{\text{P}(\text{OR})_3\}_3]$ (R = Me, Et, or Ph) derivatives. <i>Journal of the Chemical Society Dalton Transactions</i> , 1989, , 1277.	1.1	4
625	^{13}C Solid-State Cross Polarization/Magic-Angle-Spinning Nuclear Magnetic Resonance Spectra of Natural and Synthetic Melanins. <i>Pigment Cell & Melanoma Research</i> , 1988, 1, 355-357.	4.0	10
626	^{13}C solid State MAS NMR spectra of $[\text{M}_3(\text{CO})_{12}]$ complexes (M = Ru and Os). Evidence for motional effects in $[\text{Ru}_3(\text{CO})_{12}]$. <i>Inorganica Chimica Acta</i> , 1988, 146, 151-152.	1.2	27
627	Organometal clusters as models for corrosion inhibitors. The reaction of $\text{Os}_3(\text{CO})_{10}(\text{NCCH}_3)_2$ with benzotriazole. <i>Journal of Organometallic Chemistry</i> , 1988, 353, 251-257.	0.8	6
628	Carbon-metal hydrogen interchange in organometal clusters of ruthenium and osmium. <i>Organometallics</i> , 1988, 7, 856-858.	1.1	9
629	Notes. Carbon-13 and oxygen-17 nuclear magnetic resonance relaxation studies of $[\text{M}_3(\text{CO})_{12}]$ derivatives (M = Fe, Ru, or Os). <i>Journal of the Chemical Society Dalton Transactions</i> , 1988, , 791.	1.1	11
630	A New Class of Contrast Agents for Magnetic Resonance Imaging Based on Selective Reduction of Water-T2 by Chemical Exchange. <i>Investigative Radiology</i> , 1988, 23, S267-S270.	3.5	20

#	ARTICLE	IF	CITATIONS
631	Factors Determining the Proton T1 Relaxivity in Solutions Containing Gd-DTPA. Investigative Radiology, 1988, 23, S264-S266.	3.5	9
632	NONACARBONYL- η^4 -HYDRIDO (η^3 -1,3-DIMETHYL 3-ALLENYL) Triangulo-TRIRUTHENIUM (3 Ru-Ru) and NONACARBONYL- η^4 -HYDRIDO (η^3 -1,3-DIMETHYL ALLYL)Triangulo TRIRUTHENIUM (3 Ru-Ru). , 1988, , 253-255.		2
633	NONACARBONYL- η^4 -DIHYDRIDO (η^3 -2-PENTYNE)Triangulo-TRIRUTHENIUM (3 Ru-Ru). , 1988, , 258-259.		0
634	NONACARBONYL- η^4 -HYDRIDO (η^3 -t-BUTYLETYNYL)Triangulo-TRIRUTHENIUM (3 Ru-Ru). , 1988, , 256-257.		0
635	Oximation of acetyl(hydroxyimino)acetone: nuclear magnetic resonance spectroscopic, chemical, and X-ray crystallographic studies of the reaction products. Journal of the Chemical Society Perkin Transactions II, 1987, , 523.	0.9	16
636	Thermal and photochemical behaviour of organotetraruthenium clusters: solution structures and dynamics of phosphine-substituted derivatives. Journal of the Chemical Society Dalton Transactions, 1987, , 349.	1.1	4
637	Synthesis of the alkoxo(hydrido)-clusters $[\text{Ru}_3(\mu\text{-H})(\text{CO})_{10}(\mu\text{-OR})]$ [R = Me, Et, Prn, or Bun] catalysed by dinuclear carbonyl iron complexes. Journal of the Chemical Society Dalton Transactions, 1987, , 253-254.	1.1	8
638	Synthesis, structure, and ligand dynamics of $(\mu\text{-H})(\mu\text{-}\eta^2\text{-CH}_3\text{C}=\text{NCH}_2\text{CH}_3)\text{Ru}_3(\text{CO})_9$. Organometallics, 1987, 6, 2074-2078.	1.1	37
639	Proton NMR spin-lattice relaxation studies of hydridometal clusters of ruthenium and osmium. Inorganic Chemistry, 1987, 26, 2551-2552.	1.9	10
640	Bis(diphenylphosphino)methane-substituted products from $(\eta^3\text{-CCH}_3)\text{Co}_3(\text{CO})_9$. Journal of Organometallic Chemistry, 1987, 320, 229-237.	0.8	15
641	^{13}C solid state CP/MAS NMR studies of EDTA complexes. Inorganica Chimica Acta, 1987, 129, L23-L25.	1.2	14
642	Nuclear magnetic resonance of binuclear complexes containing a metallacyclopentadiene system. Journal of the Chemical Society Dalton Transactions, 1986, , 1863.	1.1	11
643	Molecular structures and dynamic behaviour of two isomers of $[\text{Ru}_3(\mu\text{-H})(\mu^3\text{-Me}_2\text{NC}_4\text{H}_4)(\text{CO})_9]$ formed from 1-dimethylaminobut-2-yne and containing η^2 -alkene-1,3-diyl and η^2 -alkene-1,2-diyl ligands respectively. Journal of the Chemical Society Dalton Transactions, 1986, , 1459-1463.	1.1	23
644	The alkyne-cluster interaction: structural, theoretical, and spectroscopic study on the parallel $\mu\text{-}\eta^2$ bonding mode in trinuclear carbonyl clusters of ruthenium and osmium. Inorganic Chemistry, 1986, 25, 4004-4010.	1.9	36
645	Dynamic behavior of the flyover bridge complexes $\text{M}_2(\text{CO})_5\text{L}\{\mu\text{-}[\text{C}(\text{R})\text{:C}(\text{R}')]\text{CO}\}$ (M = Ru, Fe; L = CO.) Tj ETQq _{1,1} 0.7843 ₁₅ 14 rgBT		
646	An examination of the unsaturation in the clusters $\text{Fe}_4(\text{CO})_{11}(\mu\text{-}4\text{-PR})_2$. Organometallics, 1986, 5, 245-252.	1.1	39
647	Deuterium isotope effects on oxygen-17 chemical shifts. Magnetic Resonance in Chemistry, 1986, 24, 919-920.	1.1	6
648	A new method for the determination of ^{17}O electric quadrupole couplings. Terminal, bridging, and capping CO groups in metallocarbonyls. Journal of Magnetic Resonance, 1986, 68, 597-599.	0.5	8

#	ARTICLE	IF	CITATIONS
649	Activation of the M π -CO bond in transition metal complexes. CO substitution reactions in di-, tri- and tetra-nuclear metal carbonyl compounds catalyzed by [Fe(CO) ₂ (η -5-C ₅ H ₅) ₂]. <i>Inorganica Chimica Acta</i> , 1986, 115, 129-133.	1.2	17
650	Solution structures and dynamics of [HO ₃ (CO) ₁₀ (η -f, η -vinyl)] complexes. <i>Inorganica Chimica Acta</i> , 1986, 111, 95-98.	1.2	6
651	The reaction of Co ₄ (CO) ₁₂ with Ph ₂ PCH ₂ CH ₂ PPh ₂ . Spectroscopic identification of polymeric products. <i>Journal of Organometallic Chemistry</i> , 1986, 309, C51-C55.	0.8	15
652	Stoichiometric and catalytic hydrogenation of the [Co ₂ (CO) ₆ (η -1,2-alkyne)] complexes. <i>Inorganica Chimica Acta</i> , 1985, 100, 97-105.	1.2	10
653	Reinvestigation of the solution structure of Co ₄ (CO) ₁₂ by ¹³ C and ⁵⁹ Co NMR. 2.. <i>Journal of Magnetic Resonance</i> , 1985, 65, 308-315.	0.5	8
654	Two-dimensional nuclear magnetic resonance for the quantitative analysis of three-site exchange in the ruthenium cluster [Ru ₃ (η -H)(CO) ₉ (MeCCHCMe)]. <i>Journal of Magnetic Resonance</i> , 1985, 65, 173-177.	0.5	14
655	Note. Two-dimensional nuclear magnetic resonance for the analysis of the carbon-13 spectra of carbonyl groups in metallocarbonyls. <i>Journal of the Chemical Society Dalton Transactions</i> , 1985, , 225.	1.1	20
656	UV photoelectron and theoretical studies of organometal carbonyl clusters of ruthenium and osmium. μ -Hydrido- μ -3-allyl and μ -hydrido- μ -3-allenyl triangulo cluster compounds. <i>Inorganic Chemistry</i> , 1985, 24, 570-575.	1.9	13
657	Mechanism of localized site exchange of carbonyl groups in HM ₃ (CO) ₉ L clusters. <i>Inorganic Chemistry</i> , 1985, 24, 231-233.	1.9	14
658	Electronic structure of ferracyclopentadienyl derivatives. UV PES ab initio study of Fe ₂ (μ -CO)(CO) ₅ (C ₄ R ₄) and Fe ₃ (μ -CO) ₂ (CO) ₆ (C ₄ R ₄). <i>Inorganic Chemistry</i> , 1985, 24, 1241-1246.	1.9	21
659	An inspection into the class of bis(alkyne)-undecacarbonyl-quadro-tetraruthenium complexes. Crystal structure and dynamic behaviour of Ru ₄ (CO) ₁₁ (η -4, η -2-MeC ₂ Ph) ₂ . <i>Inorganica Chimica Acta</i> , 1984, 85, 161-168.	1.2	18
660	η -3-S bonding mode in H ₂ M ₃ (CO) ₉ S clusters of Ru and Os. An UV π -PES and theoretical study. <i>Inorganica Chimica Acta</i> , 1984, 84, 95-100.	1.2	4
661	Photochemistry of hydrido ruthenium clusters. <i>Inorganica Chimica Acta</i> , 1984, 81, L11-L13.	1.2	2
662	¹³ C π - ¹³ C coupling constants in η -2- and η -4-(η -2-acetylene) complexes of cobalt. <i>Journal of Organometallic Chemistry</i> , 1984, 262, c1-c4.	0.8	15
663	Stoichiometric hydrogenation of the heterobimetallic (η -alkynyl) complex [η -1(η -f, 2 η -3 η -3 η -H ₂ C π -C η -1/4 C π -CH ₃)] [(CO) ₃ Fe π -Co(CO) ₃]. <i>Polyhedron</i> , 1984, 3, 175-181.	1.0	5
664	Oxygen-17 and carbon-13 relaxation studies of metal carbonyls: improved determination of ¹⁷ O electric quadrupole couplings. <i>Journal of the Chemical Society Dalton Transactions</i> , 1984, , 1863.	1.1	16
665	Electrochemical investigation of a series of organotriruthenium clusters. <i>Organometallics</i> , 1984, 3, 1374-1378.	1.1	12
666	Kinetic deuterium isotope effects on μ -hydride and carbonyl ligand migrations. <i>Organometallics</i> , 1984, 3, 1790-1795.	1.1	19

#	ARTICLE	IF	CITATIONS
667	The alkyne-cluster interaction: structural, theoretical and mechanistic studies on the $M_2M'(CO)_9(\mu_3\text{-}\eta^2\text{-alkyne})$ complex ($M = Fe$; $M' = Fe$ and Ru). <i>Organometallics</i> , 1984, 3, 1510-1515.	1.1	42
668	Solution dynamics of the osmium cluster $[Os_3(\mu\text{-H})_2(CO)_{10}]$ by ^{13}C and ^{17}O nuclear magnetic resonance. Estimation of the ^{13}C chemical shift anisotropy and ^{17}O quadrupole coupling constant. <i>Journal of the Chemical Society Dalton Transactions</i> , 1984, , 279-284.	1.1	15
669	Synthesis, structure, and isomerisation of triruthenium and triosmium clusters derived from 3-dimethylaminoprop-1-yne; X-ray crystal structure of $[Ru_3\mu\text{-}H(CO)_9(Me_2N\text{-}\eta^2\text{-}C_3H_7)]$. <i>Journal of the Chemical Society Dalton Transactions</i> , 1984, , 1981-1986.	1.1	19
670	Carbon-nitrogen bond cleavage in 1-dimethylaminobut-2-yne and derived ligands in triruthenium and triosmium clusters. <i>Journal of the Chemical Society Dalton Transactions</i> , 1984, , 1987.	1.1	22
671	The 1H And 2H NMR Spectra Of $HFeCo_3(CO)_{12}$. <i>Polyhedron</i> , 1983, 2, 1235-1240.	1.0	4
672	Synthesis and spectroscopic characterization of heterobimetallic (μ -alkynyl)iron-cobalt hexacarbonyl complexes, $[\mu_4\text{-}(1\text{-}f, 2\text{-}3, 1\text{-}2\text{-}R_2C\text{-}C\text{-}\frac{1}{4}CR_2)][(CO)_3Fe\text{-}Co(CO)_3]$ and $[\mu_4\text{-}(1\text{-}f, 2\text{-}3, 1\text{-}2\text{-}H_2C\text{-}C\text{-}\frac{1}{4}CCH_2OH)][(CO)_3Fe\text{-}Co(CO)_2(PPh_3)]$. <i>Polyhedron</i> , 1983, 2, 77-81.	1.0	23
673	Carbon-halogen bond activation at $Ru_3(CO)_{12}$. <i>Journal of Organometallic Chemistry</i> , 1983, 244, C47-C49.	0.8	13
674	Reactions of alkynes with $HFeCo_3(CO)_{12}$. Crystal structure of $FeCO_3(CO)_9[PhC_2(H)Ph]$ (PhC_2Ph). <i>Inorganica Chimica Acta</i> , 1983, 71, 141-147.	1.2	27
675	Fluxional behaviour of the binary carbonyls. A comparison of the isoelectronic and isostructural tetranuclear clusters: $Co_4(CO)_{12}$, $HFeCo_3(CO)_{12}$ and $[FeCo_3(CO)_{12}]^+$. <i>Inorganica Chimica Acta</i> , 1983, 68, 141-145.	1.2	9
676	The alkyne-cluster interaction in a ruthenium μ -butterfly μ -acetylenic carbonyl complex. An UV-PES and theoretical study. <i>Journal of Organometallic Chemistry</i> , 1983, 244, 383-391.	0.8	12
677	Oxygen-17 relaxation times of metalcarbonyls: the use of the ^{17}O electric quadrupole coupling as a new structural probe. <i>Journal of the Chemical Society Chemical Communications</i> , 1983, , 794.	2.0	5
678	Intramolecular nucleophilic attack at the μ -carbon atom of a μ_3 -alkynyl ligand in a triosmium cluster. <i>Journal of the Chemical Society Dalton Transactions</i> , 1983, , 1807-1809.	1.1	24
679	Gas-phase ultraviolet photoelectron spectra of $[\{Ni(\eta\text{-}C_5H_5)\}_2(\mu\text{-}C_2R_2)]$ ($R = H$ or CF_3) complexes. <i>Journal of the Chemical Society Dalton Transactions</i> , 1983, , 869-872.	1.1	6
680	UV photoelectron and theoretical studies of organometal carbonyl clusters of ruthenium and osmium. μ -Hydrido- μ_3 -alkynyl triangulo cluster compounds. <i>Inorganic Chemistry</i> , 1983, 22, 744-748.	1.9	18
681	UV-PES, carbon-13 NMR and theoretical studies on the alkyne-cluster interaction in $Fe_3(CO)_9(\mu_3\text{-}\eta^2\text{-}EtC_2Et)$. <i>Organometallics</i> , 1983, 2, 430-434.	1.1	35
682	Gas-phase UV photoelectron spectra of some edge-bridged decacarbonyltriosmium cluster. <i>Inorganic Chemistry</i> , 1983, 22, 3899-3903.	1.9	5
683	Acid-induced isomerisations of osmium and ruthenium clusters involving hydroxide transfer from carbon to metal atoms. X-Ray crystal structure of nonacarbonyl- μ_3 -(3,3-diphenylallene)- Tj $ETQq_1$ 1 0.784314 rgBJ/Overlock_10 Tf 50 <i>Transactions</i> , 1982, , 1625-1629.	1.1	29
684	Relationships between structure and ligand dynamics in organometal clusters. <i>Organometallics</i> , 1982, 1, 640-644.	1.1	28

#	ARTICLE	IF	CITATIONS
685	Theoretical and spectroscopic studies of (μ -butatriene)hexacarbonyldiiron compounds. <i>Inorganic Chemistry</i> , 1982, 21, 4073-4076.	1.9	9
686	Gas-phase helium(He I) photoelectron spectra of methynyltricobalt enneacarbonyl clusters. <i>Inorganic Chemistry</i> , 1982, 21, 1081-1084.	1.9	13
687	Reactions of 3-hexyne with cobalt and iron carbonyls. Crystal structure and fluxionality of $\text{FeCo}_2(\text{CO})_9\text{C}_2\text{Et}_2$. <i>Inorganic Chemistry</i> , 1982, 21, 501-505.	1.9	42
688	Unsymmetrically substituted furoxans. VII. A ^{13}C NMR study of a series of isomeric pairs of furoxans and the structure of the two isomeric chloro-phenyl-furoxans. <i>Journal of Heterocyclic Chemistry</i> , 1982, 19, 427-430.	1.4	28
689	Cluster expansion reactions of $\text{HRu}_3(\text{CO})_9\text{MeC}\hat{\text{i}}-\text{CH}\hat{\text{i}}-\text{CMe}$, a nido-Pentagonal-bipyramidal complex. <i>Inorganica Chimica Acta</i> , 1982, 57, 207-210.	1.2	11
690	Multinuclear n.m.r. studies of transition metal carbonyl clusters. <i>Inorganica Chimica Acta</i> , 1982, 62, 51-56.	1.2	20
691	Reaction of $\text{Ru}_3(\text{CO})_{12}$ with styrene. <i>Journal of Organometallic Chemistry</i> , 1982, 233, 247-252.	0.8	16
692	UV photoelectron spectra of some organometallic carbonyl osmium clusters. <i>Journal of Organometallic Chemistry</i> , 1982, 229, C27-C30.	0.8	5
693	Trinuclear osmium and ruthenium clusters containing ligands formed by dehydration and rearrangement of 1,4-dihydroxybut-2-yne. <i>Inorganic Chemistry</i> , 1981, 20, 2027-2031.	1.9	36
694	Reactions of protic acids with a hydridoorganometal cluster: $\text{HRu}_3(\text{CO})_9(\text{C}_2\text{C}(\text{CH}_3)_3)$. <i>Inorganic Chemistry</i> , 1981, 20, 1592-1597.	1.9	17
695	High-field oxygen-17 NMR spectroscopy: solution structures and dynamics of oxygen-17 enriched dodecacarbonyltetracobalt and dodecacarbonylhydroiron dicobaltate. <i>Journal of the American Chemical Society</i> , 1981, 103, 5920-5922.	6.6	45
696	Synthesis and reactivity of trinuclear osmium and dinuclear and trinuclear ruthenium compounds derived from hydroxyalkynes. <i>Journal of the Chemical Society Dalton Transactions</i> , 1981, , 828.	1.1	32
697	The structures and fluxional behaviour of the binary carbonyls; a new approach. Part 3. The fluxional behaviour of $[\text{Fe}_3(\text{CO})_{11}\text{L}]$, $\text{L} = \text{PR}_3$ or $\text{P}(\text{OR})_3$. <i>Journal of the Chemical Society Dalton Transactions</i> , 1981, , 1535.	1.1	12
698	Reaction of 4-methylpent-2-yne with dodecacarbonyltriruthenium. The structure of a novel alkyne dimerization dehydrogenation product. <i>Journal of the Chemical Society Dalton Transactions</i> , 1981, , 2023.	1.1	24
699	^{13}C n.m.r. spectra of highly ^{13}C -enriched metal carbonyls: observation of $2J_{\text{CC}}$ cis coupling constants. <i>Journal of the Chemical Society Chemical Communications</i> , 1981, , 300.	2.0	9
700	Triruthenium clusters containing dinethylamino-substituted allyl and allenyl ligands: The transmission of electronic substituent effects through clusters. <i>Journal of Organometallic Chemistry</i> , 1981, 214, C15-C18.	0.8	9
701	Iron-57 satellites in ^{13}C NMR spectra: an aid to elucidation of "hidden-processes" in the dynamics of metal carbonyls. <i>Journal of Organometallic Chemistry</i> , 1981, 214, C27-C30.	0.8	7
702	Ligand Dynamics in $\text{H}_2\text{Os}_3(\text{CO})_{10}$ and $\text{H}_2\text{Os}_3(\text{CO})_{10}\text{L}$. <i>Journal of Organometallic Chemistry</i> , 1981, 213, 207-213.	0.8	25

#	ARTICLE	IF	CITATIONS
703	Chalcogen derivatives of iron carbonyls. X. Reaction Mechanism of the CO substitution on $\text{Fe}_2(\text{CO})_6(\text{X})_2$ ($\text{X} = \text{S}, \text{Se}$) with ligands. <i>Inorganica Chimica Acta</i> , 1980, 40, 131-140.	1.2	13
704	Mixed iron and cobalt acetylenic carbonyl derivatives. <i>Inorganica Chimica Acta</i> , 1980, 40, X118.	1.2	0
705	Carbon-carbon bond cleavage in η^5 -hydroxyalkynes on reaction with dodecacarbonyl-triosmium and -triruthenium. <i>Journal of the Chemical Society Chemical Communications</i> , 1980, , 1168-1169.	2.0	16
706	Dynamic properties and solution structures of the clusters $[\text{Fe}_3(\text{CO})_9\text{X}_2]$ ($\text{X} = \text{S}, \text{Se}, \text{Te}, \text{or NMe}$) and their derivatives. <i>Journal of the Chemical Society Dalton Transactions</i> , 1980, , 46.	1.1	15
707	Regiospecific substitution of trialkylphosphines for carbon monoxide in hydridoorganoruthenium clusters. <i>Inorganic Chemistry</i> , 1980, 19, 1571-1575.	1.9	38
708	Synthesis and crystal structure of heptacarbonyl-Hydrido-(<i>t</i> -butylethynyl)-(2,4-hexadiene)-triangulo-triruthenium, $\text{HRu}_3(\text{CO})_7(\text{C}_6\text{H}_9)(\text{C}_6\text{H}_{10})$. <i>Inorganica Chimica Acta</i> , 1979, 32, 163-169.	1.2	12
709	Products of the reaction of $\text{Ru}_3(\text{CO})_{12}$ with 1,3-cyclohexadiene. X-ray structure of (η^6 -benzene)nonacarbonyl-1-cyclohexen-1,2-ylene-tetraruthenium, $\text{Ru}_4(\text{CO})_9(\text{C}_6\text{H}_6)(\text{C}_6\text{H}_8)$. <i>Inorganica Chimica Acta</i> , 1979, 34, 49-55.	1.2	29
710	^{17}O NMR spectroscopy of transition metal carbonyls. <i>Journal of Organometallic Chemistry</i> , 1979, 178, 171-175.	0.8	25
711	Solution structures and dynamic behaviour of some iron chalcogen derivatives. <i>Transition Metal Chemistry</i> , 1979, 4, 322-325.	0.7	5
712	CO activation in the reactions of iron carbonyls and alkynes. Syntheses, crystal structures, and nuclear magnetic resonance studies on hydroxyferrole and maleoyliron derivatives. <i>Journal of the Chemical Society Dalton Transactions</i> , 1979, , 1664.	1.1	37
713	Activation of alkynes by transition metals. <i>Journal of the Chemical Society Chemical Communications</i> , 1979, , 704.	2.0	7
714	Synthesis, crystal structure, and stereochemical non-rigidity of μ -[1- η^5 -2-carboxylato-1-ethylbut-1-enyl-O(2Fe)]-bis(tricarbonyl-iron) (Fe_2), a new complex obtained from iron carbonyls under apolar conditions. <i>Journal of the Chemical Society Dalton Transactions</i> , 1979, , 1155-1159.	1.1	22
715	Reactivity of the ruthenium carbonyl hydride $\text{HRu}_3(\text{CO})_9\text{C}_2\text{C}(\text{CH}_3)_3$. <i>Inorganica Chimica Acta</i> , 1978, 26, 223-226.	1.2	37
716	Crystal and molecular structure of $\text{Ru}_3(\text{CO})_6(\text{C}_{12}\text{H}_{20}\text{CO})(\text{C}_{12}\text{H}_{20})$: an example of four alkynic groups bonded to a metal cluster. <i>Inorganica Chimica Acta</i> , 1978, 27, 145-151.	1.2	19
717	Carbon-hydrogen bond activation in novel cyclopentadienyl acetylenic carbonyl clusters of iron, ruthenium and nickel. <i>Inorganica Chimica Acta</i> , 1978, 30, 9-12.	1.2	28
718	Solution structures and dynamics of $\text{Co}_4(\text{CO})_{12}$ and $\text{HFeCo}_3(\text{CO})_{12} \cdot x\text{L}_x$ ($x = 0-3$; L = group V Ligand). <i>Inorganica Chimica Acta</i> , 1978, 30, 45-49.	1.2	20
719	Solution structure and dynamics of $\text{H}_2\text{Ru}_4(\text{CO})_{13}$. <i>Inorganica Chimica Acta</i> , 1978, 29, L211-L212.	1.2	20
720	Dynamic properties of the trinuclear carbonyl cluster $\text{FeCo}_2(\text{CO})_9\text{S}$ and its Derivatives. <i>Inorganica Chimica Acta</i> , 1978, 31, 291.	1.2	0

#	ARTICLE	IF	CITATIONS
721	Nitrogen derivatives of iron carbonyls. Part 5. New routes in the mechanism of reaction of dodecacarbonyl-triangulo-tri-iron with nitro-alkanes, and X-ray analysis of μ -[acetone oximate(η^5 -NO)]- μ -isopropyl-amido-bis(tricarbonyliron) ($\text{Fe} \leftarrow \text{Fe}$). Journal of the Chemical Society Dalton Transactions, 1978, , 534-540.	1.1	17
722	Crystal structure and stereochemical non-rigidity of octacarbonyl-(η^1 -1,3-dimethyl-2-vinylcyclopentadienyl)- μ_3 -propylidyne-triangulo-tri-iron. Journal of the Chemical Society Dalton Transactions, 1977, , 227-232.	1.1	26
723	^1H and ^{13}C NMR studies of acetylenic complexes of $\text{Co}_2(\text{CO})_8$. Inorganica Chimica Acta, 1977, 22, 135-139.	1.2	59
724	Dynamic properties of the trinuclear carbonyl cluster $\text{FeCo}_2(\text{CO})_9\text{S}$ and its derivatives. Inorganica Chimica Acta, 1977, 25, 103-108.	1.2	40
725	Dynamic ^{13}C NMR spectroscopy of metal carbonyls. Progress in Nuclear Magnetic Resonance Spectroscopy, 1977, 11, 183-210.	3.9	48
726	Carbon-13 nuclear magnetic resonance spectra of η^2, η^1 -binuclear complexes prepared from dodecacarbonyltri-iron and acetylenes. Journal of the Chemical Society Dalton Transactions, 1976, , 838-840.	1.1	20
727	Multinuclear magnetic resonance studies. Part 2. Diphosphanes and dithioxodi- η^5 -phosphanes. Journal of the Chemical Society Dalton Transactions, 1976, , 2144-2153.	1.1	36
728	X-Ray molecular structure of μ -(isopropylamino)- μ -(propanone) $\text{Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 467 Td}$ (oximate-O,N)-bis(tricarbonyliron) of the Chemical Society Chemical Communications, 1976, .	2.0	11
729	Stereochemically non rigid acetylenic carbonyl complexes of iron and cobalt. Inorganica Chimica Acta, 1976, 16, L7-L8.	1.2	14
730	A ^{13}C Nmr investigation of stereochemical non rigidity in hydrido olefinic carbonyl clusters of ruthenium. Inorganica Chimica Acta, 1976, 20, 217-220.	1.2	16
731	^{13}C NMR study of methynyltricobalt enneacarbonyls. Inorganica Chimica Acta, 1976, 18, 9-11.	1.2	34
732	Stereochemically nonrigid carbonyl complexes of group VIII B metal clusters. Inorganica Chimica Acta, 1975, 15, 53-56.	1.2	61
733	Observation of localised CO exchange mechanisms at the iron and ruthenium centres of di- μ -carbonyl ($\text{Fe} : \text{Ru}_2, \text{Ru}_3$)-di- μ -hydrido ($\text{Ru}_1 : \text{Ru}_2$), $\text{Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 262 Td}$ (Ru_3)-tetrahedro-tris(tricarbonylruthenium) of the Chemical Society Chemical Communications, 1975, , 452-454.	2.0	14
734	Butadienic derivatives and metal carbonyls II. Some rearrangements of butadienic ligands in reactions with dodecacarbonyltriruthenium. Inorganica Chimica Acta, 1974, 8, 71-75.	1.2	60
735	Multinuclear magnetic resonance studies. I. Tetramethyldiphosphine- ^{13}C . Journal of Magnetic Resonance, 1974, 13, 236-238.	0.5	2
736	Intramolecular motion in tetra- <i>t</i> -butyldiphosphine. Journal of the Chemical Society Chemical Communications, 1974, , 426.	2.0	16
737	New Radiotracers, Reporter Probes and Contrast Agents. , 0, , 191-221.		0