

Toshifumi Minamoto

List of Publications by Citations

Source: <https://exaly.com/author-pdf/4971129/toshifumi-minamoto-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

129
papers

4,332
citations

32
h-index

63
g-index

147
ext. papers

6,181
ext. citations

3.6
avg, IF

5.93
L-index

#	Paper	IF	Citations
129	Estimation of fish biomass using environmental DNA. <i>PLoS ONE</i> , 2012 , 7, e35868	3.7	441
128	MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: detection of more than 230 subtropical marine species. <i>Royal Society Open Science</i> , 2015 , 2, 150088	3.3	395
127	Using environmental DNA to estimate the distribution of an invasive fish species in ponds. <i>PLoS ONE</i> , 2013 , 8, e56584	3.7	253
126	Environmental DNA metabarcoding reveals local fish communities in a species-rich coastal sea. <i>Scientific Reports</i> , 2017 , 7, 40368	4.9	203
125	Environmental DNA analysis for estimating the abundance and biomass of stream fish. <i>Freshwater Biology</i> , 2017 , 62, 30-39	3.1	179
124	The release rate of environmental DNA from juvenile and adult fish. <i>PLoS ONE</i> , 2014 , 9, e114639	3.7	177
123	Use of droplet digital PCR for estimation of fish abundance and biomass in environmental DNA surveys. <i>PLoS ONE</i> , 2015 , 10, e0122763	3.7	165
122	Surveillance of fish species composition using environmental DNA. <i>Limnology</i> , 2012 , 13, 193-197	1.7	150
121	A basin-scale application of environmental DNA assessment for rare endemic species and closely related exotic species in rivers: a case study of giant salamanders in Japan. <i>Journal of Applied Ecology</i> , 2015 , 52, 358-365	5.8	128
120	Droplet digital polymerase chain reaction (PCR) outperforms real-time PCR in the detection of environmental DNA from an invasive fish species. <i>Environmental Science & Technology</i> , 2015 , 49, 5601-8	10.3	126
119	Environmental DNA as a Snapshot of Fish Distribution: A Case Study of Japanese Jack Mackerel in Maizuru Bay, Sea of Japan. <i>PLoS ONE</i> , 2016 , 11, e0149786	3.7	121
118	A novel environmental DNA approach to quantify the cryptic invasion of non-native genotypes. <i>Molecular Ecology Resources</i> , 2016 , 16, 415-22	8.4	88
117	Water temperature-dependent degradation of environmental DNA and its relation to bacterial abundance. <i>PLoS ONE</i> , 2017 , 12, e0176608	3.7	87
116	The use of environmental DNA of fishes as an efficient method of determining habitat connectivity. <i>Ecological Indicators</i> , 2016 , 62, 147-153	5.8	83
115	A simple method for preserving environmental DNA in water samples at ambient temperature by addition of cationic surfactant. <i>Limnology</i> , 2017 , 18, 233-241	1.7	78
114	Effect of water temperature and fish biomass on environmental DNA shedding, degradation, and size distribution. <i>Ecology and Evolution</i> , 2019 , 9, 1135-1146	2.8	76
113	Rapid degradation of longer DNA fragments enables the improved estimation of distribution and biomass using environmental DNA. <i>Molecular Ecology Resources</i> , 2017 , 17, e25-e33	8.4	73

112	Techniques for the practical collection of environmental DNA: filter selection, preservation, and extraction. <i>Limnology</i> , 2016 , 17, 23-32	1.7	64
111	Environmental DNA reflects spatial and temporal jellyfish distribution. <i>PLoS ONE</i> , 2017 , 12, e0173073	3.7	56
110	Effects of sample processing on the detection rate of environmental DNA from the Common Carp (<i>Cyprinus carpio</i>). <i>Biological Conservation</i> , 2015 , 183, 64-69	6.2	53
109	Dispersion and degradation of environmental DNA from caged fish in a marine environment. <i>Fisheries Science</i> , 2019 , 85, 327-337	1.9	46
108	Comparing local- and regional-scale estimations of the diversity of stream fish using eDNA metabarcoding and conventional observation methods. <i>Freshwater Biology</i> , 2018 , 63, 569-580	3.1	45
107	Detection of cyprinid herpesvirus 3 DNA in river water during and after an outbreak. <i>Veterinary Microbiology</i> , 2009 , 135, 261-6	3.3	44
106	Use of environmental DNA to survey the distribution of an invasive submerged plant in ponds. <i>Freshwater Science</i> , 2016 , 35, 748-754	2	40
105	Environmental DNA method for estimating salamander distribution in headwater streams, and a comparison of water sampling methods. <i>PLoS ONE</i> , 2017 , 12, e0176541	3.7	37
104	A novel isoform of vertebrate ancient opsin in a smelt fish, <i>Plecoglossus altivelis</i> . <i>Biochemical and Biophysical Research Communications</i> , 2002 , 290, 280-6	3.4	36
103	Quantification of cyprinid herpesvirus 3 in environmental water by using an external standard virus. <i>Applied and Environmental Microbiology</i> , 2010 , 76, 161-8	4.8	35
102	Detection of cyprinid herpesvirus-3 DNA in lake plankton. <i>Research in Veterinary Science</i> , 2011 , 90, 530-22.5		34
101	Transmission dynamics of an emerging infectious disease in wildlife through host reproductive cycles. <i>ISME Journal</i> , 2011 , 5, 244-51	11.9	34
100	Nuclear internal transcribed spacer-1 as a sensitive genetic marker for environmental DNA studies in common carp <i>Cyprinus carpio</i> . <i>Molecular Ecology Resources</i> , 2017 , 17, 324-333	8.4	32
99	Molecular cloning of lysozyme-encoding cDNAs expressed in the salivary gland of a wood-feeding termite, <i>Reticulitermes speratus</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2002 , 32, 1615-24	4.5	32
98	Environmental DNA analysis reveals the spatial distribution, abundance, and biomass of Japanese eels at the river-basin scale. <i>Aquatic Conservation: Marine and Freshwater Ecosystems</i> , 2019 , 29, 361-373	2.6	32
97	Molecular cloning of cone opsin genes and their expression in the retina of a smelt, Ayu (<i>Plecoglossus altivelis</i> , Teleostei). <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 2005 , 140, 197-205	2.3	29
96	On-site filtration of water samples for environmental DNA analysis to avoid DNA degradation during transportation. <i>Ecological Research</i> , 2016 , 31, 963-967	1.9	28
95	Sedimentary eDNA provides different information on timescale and fish species composition compared with aqueous eDNA. <i>Environmental DNA</i> , 2020 , 2, 505-518	7.6	27

94	Evaluating intraspecific genetic diversity using environmental DNA and denoising approach: A case study using tank water. <i>Environmental DNA</i> , 2020 , 2, 42-52	7.6	26
93	An illustrated manual for environmental DNA research: Water sampling guidelines and experimental protocols. <i>Environmental DNA</i> , 2021 , 3, 8-13	7.6	26
92	Effects of water pH and proteinase K treatment on the yield of environmental DNA from water samples. <i>Limnology</i> , 2017 , 18, 1-7	1.7	25
91	Use of a Filter Cartridge for Filtration of Water Samples and Extraction of Environmental DNA. <i>Journal of Visualized Experiments</i> , 2016 ,	1.6	25
90	Seasonal distribution of cyprinid herpesvirus 3 in Lake Biwa, Japan. <i>Applied and Environmental Microbiology</i> , 2009 , 75, 6900-4	4.8	25
89	Evaluation of the Environmental DNA Method for Estimating Distribution and Biomass of Submerged Aquatic Plants. <i>PLoS ONE</i> , 2016 , 11, e0156217	3.7	25
88	Application of environmental DNA analysis for the detection of <i>Opisthorchis viverrini</i> DNA in water samples. <i>Acta Tropica</i> , 2017 , 169, 1-7	3.2	22
87	Water sampling for environmental DNA surveys by using an unmanned aerial vehicle. <i>Limnology and Oceanography: Methods</i> , 2017 , 15, 939-944	2.6	21
86	Nationwide Cyprinid herpesvirus 3 contamination in natural rivers of Japan. <i>Research in Veterinary Science</i> , 2012 , 93, 508-14	2.5	21
85	Seasonal reactivation enables Cyprinid herpesvirus 3 to persist in a wild host population. <i>FEMS Microbiology Ecology</i> , 2014 , 87, 536-42	4.3	20
84	Usefulness of environmental DNA for detecting <i>Schistosoma mansoni</i> occurrence sites in Madagascar. <i>International Journal of Infectious Diseases</i> , 2018 , 76, 130-136	10.5	20
83	Reservoirs of Cyprinid herpesvirus 3 (CyHV-3) DNA in sediments of natural lakes and ponds. <i>Veterinary Microbiology</i> , 2012 , 155, 183-90	3.3	19
82	Quantitative monitoring of multispecies fish environmental DNA using high-throughput sequencing		19
81	Environmental DNA metabarcoding to detect pathogenic <i>Leptospira</i> and associated organisms in leptospirosis-endemic areas of Japan. <i>Scientific Reports</i> , 2019 , 9, 6575	4.9	18
80	Distinct seasonal migration patterns of Japanese native and non-native genotypes of common carp estimated by environmental DNA. <i>Ecology and Evolution</i> , 2017 , 7, 8515-8522	2.8	18
79	Real-time polymerase chain reaction assays for environmental DNA detection of three salmonid fish in Hokkaido, Japan: Application to winter surveys. <i>Ecological Research</i> , 2019 , 34, 237-242	1.9	18
78	Stress response to daily temperature fluctuations in common carp, <i>Cyprinus carpio</i> L.. <i>Hydrobiologia</i> , 2011 , 675, 65-73	2.4	17
77	Estimating shedding and decay rates of environmental nuclear DNA with relation to water temperature and biomass. <i>Environmental DNA</i> , 2020 , 2, 140-151	7.6	17

76	Identifying a breeding habitat of a critically endangered fish, <i>Acheilognathus typus</i> , in a natural river in Japan. <i>Die Naturwissenschaften</i> , 2017 , 104, 100	2	16
75	Particle Size Distribution of Environmental DNA from the Nuclei of Marine Fish. <i>Environmental Science & Technology</i> , 2019 , 53, 9947-9956	10.3	15
74	First use of oceanic environmental DNA to study the spawning ecology of the Japanese eel <i>Anguilla japonica</i> . <i>Marine Ecology - Progress Series</i> , 2019 , 609, 187-196	2.6	15
73	Circadian clock in <i>Ciona intestinalis</i> revealed by microarray analysis and oxygen consumption. <i>Journal of Biochemistry</i> , 2010 , 147, 175-84	3.1	14
72	Isopropanol precipitation method for collecting fish environmental DNA. <i>Limnology and Oceanography: Methods</i> , 2017 , 15, 212-218	2.6	13
71	Effects of sampling seasons and locations on fish environmental DNA metabarcoding in dam reservoirs. <i>Ecology and Evolution</i> , 2020 , 10, 5354-5367	2.8	13
70	Environmental DNA reveals nonmigratory individuals of <i>Palaemon paucidens</i> overwintering in Lake Biwa shallow waters. <i>Freshwater Science</i> , 2018 , 37, 307-314	2	13
69	A new method for random mutagenesis by error-prone polymerase chain reaction using heavy water. <i>Journal of Biotechnology</i> , 2012 , 157, 71-4	3.7	13
68	An emerging infectious pathogen endangers an ancient lineage of common carp by acting synergistically with conspecific exotic strains. <i>Animal Conservation</i> , 2013 , 16, 324-330	3.2	13
67	Comparison of inhibition resistance among PCR reagents for detection and quantification of environmental DNA. <i>Environmental DNA</i> , 2019 , 1, 359-367	7.6	12
66	Determining an effective sampling method for eDNA metabarcoding: a case study for fish biodiversity monitoring in a small, natural river. <i>Limnology</i> , 2021 , 22, 221-235	1.7	12
65	Messenger RNA typing of environmental RNA (eRNA): A case study on zebrafish tank water with perspectives for the future development of eRNA analysis on aquatic vertebrates. <i>Environmental DNA</i> , 2021 , 3, 14-21	7.6	12
64	Using environmental DNA to estimate the seasonal distribution and habitat preferences of a Japanese basket clam in Lake Shinji, Japan. <i>Estuarine, Coastal and Shelf Science</i> , 2019 , 221, 15-20	2.9	11
63	Detection of <i>Schistosoma japonicum</i> and <i>Oncomelania hupensis quadrasi</i> environmental DNA and its potential utility to schistosomiasis japonica surveillance in the Philippines. <i>PLoS ONE</i> , 2019 , 14, e0224617	2.7	11
62	Environmental DNA analysis shows high potential as a tool for estimating intraspecific genetic diversity in a wild fish population. <i>Molecular Ecology Resources</i> , 2020 , 20, 1248-1258	8.4	11
61	Comparing the efficiency of open and enclosed filtration systems in environmental DNA quantification for fish and jellyfish. <i>PLoS ONE</i> , 2020 , 15, e0231718	3.7	10
60	Biomass-dependent emission of environmental DNA in jack mackerel <i>Trachurus japonicus</i> juveniles. <i>Journal of Fish Biology</i> , 2019 , 95, 979-981	1.9	10
59	Effects of daily temperature fluctuation on the survival of carp infected with Cyprinid herpesvirus 3. <i>Aquaculture</i> , 2014 , 433, 208-213	4.4	10

58	Molecular cloning and characterization of rhodopsin in a teleost (<i>Plecoglossus altivelis</i> , Osmeridae). <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 2003 , 134, 559-70	2.3	10
57	Sedimentary DNA tracks decadal-centennial changes in fish abundance. <i>Communications Biology</i> , 2020 , 3, 558	6.7	10
56	Environmental DNA monitoring for short-term reproductive migration of endemic anadromous species, Shishamo smelt (<i>Spirinchus lanceolatus</i>). <i>Environmental DNA</i> , 2020 , 2, 130-139	7.6	9
55	Estimating fish population abundance by integrating quantitative data on environmental DNA and hydrodynamic modelling. <i>Molecular Ecology</i> , 2021 , 30, 3057-3067	5.7	8
54	Discovery of an unrecorded population of Yamato salamander (<i>Hynobius vandenburghi</i>) by GIS and eDNA analysis. <i>Environmental DNA</i> , 2019 , 1, 281-289	7.6	8
53	Effect of different personal histories on valuation for forest ecosystem services in urban areas: A case study of Mt. Rokko, Kobe, Japan. <i>Urban Forestry and Urban Greening</i> , 2017 , 28, 110-117	5.4	8
52	Multiplex real-time PCR enables the simultaneous detection of environmental DNA from freshwater fishes: a case study of three exotic and three threatened native fishes in Japan. <i>Biological Invasions</i> , 2020 , 22, 455-471	2.7	8
51	Spatial variation and temporal stability of littoral water temperature relative to lakeshore morphometry: environmental analysis from the view of fish thermal ecology. <i>Limnology</i> , 2010 , 11, 71-76 ^{1.7}	1.7	7
50	Environmental DNA detection and quantification of invasive red-eared sliders, <i>Trachemys scripta</i> , in ponds and the influence of water quality. <i>PeerJ</i> , 2019 , 7, e8155	3.1	7
49	Habitat selection and migration of the common shrimp, <i>Palaemon paucidens</i> in Lake Biwa, Japan. An eDNA-based study. <i>Environmental DNA</i> , 2019 , 1, 54-63	7.6	6
48	<i>Apis cerana japonica</i> discriminates between floral color phases of the oriental orchid, <i>Cymbidium floribundum</i> . <i>Zoological Science</i> , 2010 , 27, 901-6	0.8	6
47	Estimations of Riverine Distribution, Abundance, and Biomass of Anguillid Eels in Japan and Taiwan Using Environmental DNA Analysis. <i>Zoological Studies</i> , 2020 , 59, e17	0.6	6
46	Monitoring fish pathogenic viruses in natural lakes in Yunnan, China. <i>Limnology</i> , 2015 , 16, 69-77	1.7	5
45	Analysis of Environmental DNA and Edaphic Factors for the Detection of the Snail Intermediate Host. <i>Pathogens</i> , 2019 , 8,	4.5	5
44	Differences between domesticated Eurasian and Japanese indigenous strains of the common carp (<i>Cyprinus carpio</i>) in cortisol release following acute stress. <i>Ichthyological Research</i> , 2014 , 61, 165-168	0.8	5
43	Suppression of environmental DNA degradation in water samples associated with different storage temperature and period using benzalkonium chloride. <i>Limnology and Oceanography: Methods</i> , 2020 , 18, 437-445	2.6	5
42	Seasonal change in environmental DNA concentration of a submerged aquatic plant species. <i>Freshwater Science</i> , 2019 , 38, 654-660	2	4
41	Complex interactions between environmental DNA (eDNA) state and water chemistries on eDNA persistence suggested by meta-analyses. <i>Molecular Ecology Resources</i> , 2021 , 21, 1490-1503	8.4	4

40	Broad-scale detection of environmental DNA for an invasive macrophyte and the relationship between DNA concentration and coverage in rivers. <i>Biological Invasions</i> , 2021 , 23, 507-520	2.7	4
39	Simultaneous absolute quantification and sequencing of fish environmental DNA in a mesocosm by quantitative sequencing technique. <i>Scientific Reports</i> , 2021 , 11, 4372	4.9	4
38	The life history with seasonal migration of the lacustrine shrimp <i>Palaemon paucidens</i> in an ancient lake in Japan. <i>Ecosphere</i> , 2019 , 10, e02628	3.1	3
37	Selective collection of long fragments of environmental DNA using larger pore size filter. <i>Science of the Total Environment</i> , 2020 , 735, 139462	10.2	3
36	Cloning of genes encoding the visual pigments in the silkworm, <i>Bombyx mori</i> . <i>Applied Entomology and Zoology</i> , 1998 , 33, 199-204	1.5	3
35	Environmental DNA revealed the fish community of Hokkaido Island, Japan, after invasion by rainbow trout. <i>Biodiversity Data Journal</i> , 2020 , 8, e56876	1.8	3
34	A molecular survey based on eDNA to assess the presence of a clown featherback () in a confined environment. <i>PeerJ</i> , 2020 , 8, e10338	3.1	3
33	Evaluating intraspecific genetic diversity of a fish population using environmental DNA: An approach to distinguish true haplotypes from erroneous sequences		2
32	Estimating fish population abundance by integrating quantitative data on environmental DNA and hydrodynamic modelling		2
31	Compilation of real-time PCR conditions toward the standardization of environmental DNA methods. <i>Ecological Research</i> , 2021 , 36, 379-388	1.9	2
30	Molecular detection of giant snakeheads, <i>Channa micropeltes</i> (Cuvier, 1831), one of the most troublesome fish species. <i>Scientific Reports</i> , 2021 , 11, 9943	4.9	2
29	Utility of environmental DNA analysis for effective monitoring of invasive fish species in reservoirs. <i>Ecosphere</i> , 2021 , 12, e03643	3.1	2
28	Application of environmental DNA metabarcoding in a lake with extensive algal blooms. <i>Limnology</i> , 2021 , 22, 363-370	1.7	2
27	Characterizing the spatial and temporal occurrence patterns of the endangered botiid loach <i>Parabotia curtus</i> by environmental DNA analysis using a newly developed species-specific primer set. <i>Ichthyological Research</i> , 2021 , 68, 152-157	0.8	2
26	eDNA-based detection of a vulnerable crocodile newt (<i>Tylototriton uyenoi</i>) to influence government policy and raise public awareness. <i>Diversity and Distributions</i> , 2021 , 27, 1958-1965	5	2
25	Linking the state of environmental DNA to its application for biomonitoring and stock assessment: Targeting mitochondrial/nuclear genes, and different DNA fragment lengths and particle sizes. <i>Environmental DNA</i> ,	7.6	2
24	Detection of herbivory: eDNA detection from feeding marks on leaves. <i>Environmental DNA</i> , 2020 , 2, 627-634		1
23	Random Mutagenesis by Error-Prone Polymerase Chain Reaction Using a Heavy Water Solvent. <i>Methods in Molecular Biology</i> , 2017 , 1498, 491-495	1.4	1

22	Spatial-temporal analysis of water temperatures during spring in Lake Erhai, China: implications for fisheries. <i>Inland Waters</i> , 2012 , 2, 129-136	2.4	1
21	Expression of spliced variants of period mRNA in the Japanese honeybee <i>Apis cerana japonica</i> . <i>Biological Rhythm Research</i> , 2012 , 43, 125-135	0.8	1
20	Prevalence of Antimicrobial-Resistant <i>Escherichia coli</i> in Migratory Greater White-Fronted Geese (<i>Anser albifrons</i>) and their Habitat in Miyajimanuma, Japan. <i>Journal of Wildlife Diseases</i> , 2021 , 57, 954-958	1.3	1
19	STUDY ON SEASONAL, DAILY AND MORPHOLOGICAL CHANGE OF ENVIRONMENTAL DNA FOR SEAGRASS BED. <i>Journal of Japan Society of Civil Engineers Ser B2 (Coastal Engineering)</i> , 2019 , 75, I_1075-I_1080	0.1	1
18	Projection range of eDNA analysis in marshes: a suggestion from the Siberian salamander (<i>Pleurodeles waltl</i>) inhabiting the Kushiro marsh, Japan. <i>PeerJ</i> , 2020 , 8, e9764	3.1	1
17	Effects of species traits and ecosystem characteristics on species detection by eDNA metabarcoding in lake fish communities		1
16	Environmental DNA detection of an invasive ant species (<i>Linepithema humile</i>) from soil samples. <i>Scientific Reports</i> , 2021 , 11, 10712	4.9	1
15	Continuous prevalence of VEB-3 extended-spectrum β -lactamase-producing <i>Aeromonas hydrophila</i> in a local river in gifu city, Japan. <i>Microbiology and Immunology</i> , 2021 , 65, 99-100	2.7	1
14	Development of primer/probe sets for environmental DNA-based monitoring of pond smelt <i>Hypomesus nipponensis</i> and Japanese icefish <i>Salangichthys microdon</i> . <i>Landscape and Ecological Engineering</i> , 1	2	1
13	Environment-human linkage as a model for environmental diseases. <i>Ecological Research</i> , 2011 , 26, 1011-1016	1.9	0
12	Development of environmental DNA detection assays for snakes in paddy fields in Japan. <i>Landscape and Ecological Engineering</i> , 1	2	0
11	Environmental DNA emission by two carangid fishes in single and mixed-species tanks. <i>Fisheries Science</i> , 1	1.9	0
10	Spatiotemporal distribution of <i>Flavobacterium psychrophilum</i> and ayu <i>Plecoglossus altivelis</i> in rivers revealed by environmental DNA analysis. <i>Fisheries Science</i> , 2021 , 87, 321-330	1.9	0
9	Revealing an Invasion Risk of Fish Species in Qingdao Underwater World by Environmental DNA Metabarcoding. <i>Journal of Ocean University of China</i> , 2021 , 20, 124-136	1	0
8	Environmental DNA preserved in marine sediment for detecting jellyfish blooms after a tsunami. <i>Scientific Reports</i> , 2021 , 11, 16830	4.9	0
7	Pre-centrifugation before DNA extraction mitigates extraction efficiency reduction of environmental DNA caused by the preservative solution (benzalkonium chloride) remaining in the filters. <i>Limnology</i> , 1	1.7	0
6	Detection of multiple mycetozoa pathogens using fungal metabarcoding analysis of soil DNA in an endemic area of Sudan.. <i>PLoS Neglected Tropical Diseases</i> , 2022 , 16, e0010274	4.8	0
5	Detection of periodic patterns in microarray data reveals novel oscillating transcripts of biological rhythms in <i>Ciona intestinalis</i> . <i>Artificial Life and Robotics</i> , 2015 , 20, 347-352	0.6	

4	Seasonal changes in the distribution of black sea bream <i>Acanthopagrus schlegelii</i> estimated by environmental DNA. <i>Fisheries Science</i> , 2022 , 88, 91-107	1.9
3	Autumn dispersal and limited success of reproduction of the deepbody bitterling (<i>Acheilognathus longipinnis</i>) in terrestrialized floodplain. <i>Knowledge and Management of Aquatic Ecosystems</i> , 2022 , 4	1.4
2	SEASONAL CHANGES IN ENVIRONMENTAL DNA IN SEAGRASS BEDS AND EXAMINATION OF WATER SAMPLING PROCEDURE. <i>Journal of Japan Society of Civil Engineers Ser B2 (Coastal Engineering)</i> , 2020 , 76, I_943-I_948	0.1
1	STUDY ON AN ENVIRONMENTAL DNA ANALYSIS PROTOCOL FOR SEAGRASS BED MONITORING. <i>Journal of Japan Society of Civil Engineers Ser B2 (Coastal Engineering)</i> , 2021 , 77, I_895-I_900	0.1