

Tianhong Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4971378/publications.pdf>

Version: 2024-02-01

147
papers

2,298
citations

279798

23
h-index

315739

38
g-index

153
all docs

153
docs citations

153
times ranked

3077
citing authors

#	ARTICLE	IF	CITATIONS
1	Immediate psychological distress in quarantined patients with COVID-19 and its association with peripheral inflammation: A mixed-method study. <i>Brain, Behavior, and Immunity</i> , 2020, 88, 17-27.	4.1	303
2	Prodromal psychosis detection in a counseling center population in China: An epidemiological and clinical study. <i>Schizophrenia Research</i> , 2014, 152, 391-399.	2.0	104
3	Abnormal white matter microstructure in drug-naïve first episode schizophrenia patients before and after eight weeks of antipsychotic treatment. <i>Schizophrenia Research</i> , 2016, 172, 1-8.	2.0	75
4	Role of childhood traumatic experience in personality disorders in China. <i>Comprehensive Psychiatry</i> , 2012, 53, 829-836.	3.1	67
5	Prediction of psychosis in prodrome: development and validation of a simple, personalized risk calculator. <i>Psychological Medicine</i> , 2019, 49, 1990-1998.	4.5	59
6	Elevated serum levels of FGF-2, NGF and IGF-1 in patients with manic episode of bipolar disorder. <i>Psychiatry Research</i> , 2014, 218, 54-60.	3.3	58
7	Validating the Predictive Accuracy of the NAPLS-2 Psychosis Risk Calculator in a Clinical High-Risk Sample From the SHARP (Shanghai At Risk for Psychosis) Program. <i>American Journal of Psychiatry</i> , 2018, 175, 906-908.	7.2	54
8	Meta-analysis of cognitive function in Chinese first-episode schizophrenia: MATRICS Consensus Cognitive Battery (MCCB) profile of impairment. <i>Annals of General Psychiatry</i> , 2019, 32, e100043.	3.1	52
9	Functional connectome organization predicts conversion to psychosis in clinical high-risk youth from the SHARP program. <i>Molecular Psychiatry</i> , 2020, 25, 2431-2440.	7.9	49
10	Mobile health in China: Current status and future development. <i>Asian Journal of Psychiatry</i> , 2014, 10, 101-104.	2.0	43
11	Decreased serum fibroblast growth factor - 2 levels in pre- and post-treatment patients with major depressive disorder. <i>Neuroscience Letters</i> , 2014, 579, 168-172.	2.1	39
12	Altered neuronal spontaneous activity correlates with glutamate concentration in medial prefrontal cortex of major depressed females: An fMRI-MRS study. <i>Journal of Affective Disorders</i> , 2016, 201, 153-161.	4.1	39
13	The relationship between facial emotion recognition and executive functions in first-episode patients with schizophrenia and their siblings. <i>BMC Psychiatry</i> , 2015, 15, 241.	2.6	38
14	Prevalence of personality disorders using two diagnostic systems in psychiatric outpatients in Shanghai, China: a comparison of uni-axial and multi-axial formulation. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012, 47, 1409-1417.	3.1	36
15	Reduced γ -Aminobutyric Acid and Glutamate+Glutamine Levels in Drug-Naïve Patients with First-Episode Schizophrenia but Not in Those at Ultrahigh Risk. <i>Neural Plasticity</i> , 2016, 2016, 1-9.	2.2	34
16	Increased resting-state global functional connectivity density of default mode network in schizophrenia subjects treated with electroconvulsive therapy. <i>Schizophrenia Research</i> , 2018, 197, 192-199.	2.0	33
17	Neuropsychological Impairment in Prodromal, First-Episode, and Chronic Psychosis: Assessing RBANS Performance. <i>PLoS ONE</i> , 2015, 10, e0125784.	2.5	29
18	Add-on rTMS for the acute treatment of depressive symptoms is probably more effective in adolescents than in adults: Evidence from real-world clinical practice. <i>Brain Stimulation</i> , 2019, 12, 103-109.	1.6	29

#	ARTICLE	IF	CITATIONS
19	Altered Cellular White Matter But Not Extracellular Free Water on Diffusion MRI in Individuals at Clinical High Risk for Psychosis. <i>American Journal of Psychiatry</i> , 2019, 176, 820-828.	7.2	28
20	Aberrant resting-state functional connectivity of salience network in first-episode schizophrenia. <i>Brain Imaging and Behavior</i> , 2020, 14, 1350-1360.	2.1	28
21	P300 as an index of transition to psychosis and of remission: Data from a clinical high risk for psychosis study and review of literature. <i>Schizophrenia Research</i> , 2020, 226, 74-83.	2.0	26
22	Psychometric Properties of Prodromal Questionnaire-Brief Version among Chinese Help-Seeking Individuals. <i>PLoS ONE</i> , 2016, 11, e0148935.	2.5	26
23	Brain functional connectivity data enhance prediction of clinical outcome in youth at risk for psychosis. <i>NeuroImage: Clinical</i> , 2020, 26, 102108.	2.7	25
24	Baseline Cortical Thickness Reductions in Clinical High Risk for Psychosis: Brain Regions Associated with Conversion to Psychosis Versus Non-Conversion as Assessed at One-Year Follow-Up in the Shanghai-At-Risk-for-Psychosis (SHARP) Study. <i>Schizophrenia Bulletin</i> , 2021, 47, 562-574.	4.3	25
25	Reduced functional connectivity between bilateral precuneus and contralateral parahippocampus in schizotypal personality disorder. <i>BMC Psychiatry</i> , 2017, 17, 48.	2.6	24
26	Functional near-infrared spectroscopy (fNIRS) as a tool to assist the diagnosis of major psychiatric disorders in a Chinese population. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 745-757.	3.2	24
27	Calculating individualized risk components using a mobile app-based risk calculator for clinical high risk of psychosis: findings from ShangHai At Risk for Psychosis (SHARP) program. <i>Psychological Medicine</i> , 2021, 51, 653-660.	4.5	24
28	Real-world effectiveness of antipsychotic treatment in psychosis prevention in a 3-year cohort of 517 individuals at clinical high risk from the SHARP (ShangHai At Risk for Psychosis). <i>Australian and New Zealand Journal of Psychiatry</i> , 2020, 54, 696-706.	2.3	24
29	Screening schizotypal personality disorder for detection of clinical high risk of psychosis in Chinese mental health services. <i>Psychiatry Research</i> , 2015, 228, 664-670.	3.3	22
30	Salivary microbiome profiling reveals a dysbiotic schizophrenia-associated microbiota. <i>NPJ Schizophrenia</i> , 2021, 7, 51.	3.6	22
31	Clinical subtypes that predict conversion to psychosis: A canonical correlation analysis study from the ShangHai At Risk for Psychosis program. <i>Australian and New Zealand Journal of Psychiatry</i> , 2020, 54, 482-495.	2.3	21
32	Progressive decline of cognition during the conversion from prodrome to psychosis with a characteristic pattern of the theory of mind compensated by neurocognition. <i>Schizophrenia Research</i> , 2018, 195, 554-559.	2.0	20
33	Duration of untreated prodromal symptoms in a Chinese sample at a high risk for psychosis: demographic, clinical, and outcome. <i>Psychological Medicine</i> , 2018, 48, 1274-1281.	4.5	20
34	ECT-induced brain plasticity correlates with positive symptom improvement in schizophrenia by voxel-based morphometry analysis of grey matter. <i>Brain Stimulation</i> , 2019, 12, 319-328.	1.6	20
35	Tumor Necrosis Factor- α Variations in Patients With Major Depressive Disorder Before and After Antidepressant Treatment. <i>Frontiers in Psychiatry</i> , 2020, 11, 518837.	2.6	20
36	Both volumetry and functional connectivity of Heschl's gyrus are associated with auditory P300 in first episode schizophrenia. <i>Schizophrenia Research</i> , 2014, 160, 57-66.	2.0	18

#	ARTICLE	IF	CITATIONS
37	Dynamic Functional Connectivity Within the Fronto-Limbic Network Induced by Intermittent Theta-Burst Stimulation: A Pilot Study. <i>Frontiers in Neuroscience</i> , 2019, 13, 944.	2.8	18
38	Cognitive dysfunction in a psychotropic medication-naïve, clinical high-risk sample from the Shanghai-At-Risk-for-Psychosis (SHARP) study: Associations with clinical outcomes. <i>Schizophrenia Research</i> , 2020, 226, 138-146.	2.0	18
39	Interaction of social role functioning and coping in people with recent-onset attenuated psychotic symptoms: a case study of three Chinese women at clinical high risk for psychosis. <i>Neuropsychiatric Disease and Treatment</i> , 2015, 11, 1647.	2.2	17
40	Faux pas recognition performance in a help-seeking population at clinical high risk of psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2016, 266, 71-78.	3.2	17
41	Abnormal auditory-evoked gamma band oscillations in first-episode schizophrenia during both eye open and eye close states. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 86, 279-286.	4.8	17
42	Using “WeChat” online social networking in a real-world needs analysis of family members of youths at clinical high risk of psychosis. <i>Australian and New Zealand Journal of Psychiatry</i> , 2018, 52, 375-382.	2.3	17
43	Cognitive Behavioral Therapy for Prodromal Stage of Psychosis—Outcomes for Transition, Functioning, Distress, and Quality of Life: A Systematic Review and Meta-analysis. <i>Schizophrenia Bulletin</i> , 2022, 48, 8-19.	4.3	17
44	Age and Remission of Personality Pathology in the Psychotic Disorders Compared to Mood and/or Anxiety Disorders. <i>International Journal of Psychiatry in Medicine</i> , 2012, 44, 241-255.	1.8	16
45	Correlation of social cognition and neurocognition on psychotic outcome: a naturalistic follow-up study of subjects with attenuated psychosis syndrome. <i>Scientific Reports</i> , 2016, 6, 35017.	3.3	15
46	Co-morbidity of personality disorder in schizophrenia among psychiatric outpatients in China: data from epidemiologic survey in a clinical population. <i>BMC Psychiatry</i> , 2016, 16, 224.	2.6	15
47	Theory of Mind Impairments in Youth at Clinical High Risk of Psychosis. <i>Psychiatry (New York)</i> , 2016, 79, 40-55.	0.7	15
48	Association between catechol-O-methyltransferase genetic variation and functional connectivity in patients with first-episode schizophrenia. <i>Schizophrenia Research</i> , 2018, 199, 214-220.	2.0	15
49	Co-Morbidity of DSM-IV Personality Disorder in Major Depressive Disorder Among Psychiatric Outpatients in China: A Further Analysis of an Epidemiologic Survey in a Clinical Population. <i>Frontiers in Psychiatry</i> , 2019, 10, 833.	2.6	15
50	A comparison of conversion rates, clinical profiles and predictors of outcomes in two independent samples of individuals at clinical high risk for psychosis in China. <i>Schizophrenia Research</i> , 2018, 197, 509-515.	2.0	14
51	Conversion to psychosis in adolescents and adults: similar proportions, different predictors. <i>Psychological Medicine</i> , 2021, 51, 2003-2011.	4.5	14
52	Prolonged cortical silent period among drug-naive subjects at ultra-high risk of psychosis. <i>Schizophrenia Research</i> , 2014, 160, 124-130.	2.0	13
53	Eye Movement Indices in the Study of Depressive Disorder. <i>Shanghai Archives of Psychiatry</i> , 2016, 28, 326-334.	0.7	13
54	Changes in event-related potentials in patients with first-episode schizophrenia and their siblings. <i>BMC Psychiatry</i> , 2017, 17, 20.	2.6	12

#	ARTICLE	IF	CITATIONS
55	Altered functional connectivity of the thalamus induced by modified electroconvulsive therapy for schizophrenia. <i>Schizophrenia Research</i> , 2020, 218, 209-218.	2.0	12
56	Abnormal Function in Dentate Nuclei Precedes the Onset of Psychosis: A Resting-State fMRI Study in High-Risk Individuals. <i>Schizophrenia Bulletin</i> , 2021, 47, 1421-1430.	4.3	12
57	Temporal Dynamics in Degree Centrality of Brain Functional Connectome in First-Episode Schizophrenia with Different Short-Term Treatment Responses: A Longitudinal Study. <i>Neuropsychiatric Disease and Treatment</i> , 2021, Volume 17, 1505-1516.	2.2	12
58	Nerve growth factor variations in patients with mood disorders: no changes in eight weeks of clinical treatment. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 835.	2.2	11
59	Relationship between duration of untreated prodromal symptoms and symptomatic and functional recovery. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2019, 269, 871-877.	3.2	11
60	A potential objective marker in first-episode schizophrenia based on abnormal niacin response. <i>Schizophrenia Research</i> , 2022, 243, 405-412.	2.0	11
61	Effective schizophrenia recognition using discriminative eye movement features and model-metric based features. <i>Pattern Recognition Letters</i> , 2020, 138, 608-616.	4.2	11
62	Spatio-Temporal Self-Attention Network for Video Saliency Prediction. <i>IEEE Transactions on Multimedia</i> , 2023, 25, 1161-1174.	7.2	11
63	Transdiagnostic Dimensions towards Personality Pathology and Childhood Traumatic Experience in a Clinical Sample: Subtype Classification by a Cross-sectional Analysis. <i>Scientific Reports</i> , 2019, 9, 11248.	3.3	10
64	Comparative efficacy of add-on rTMS in treating the somatic and psychic anxiety symptoms of depression comorbid with anxiety in adolescents, adults, and elderly patientsâ€”A real-world clinical application. <i>Journal of Affective Disorders</i> , 2020, 276, 305-311.	4.1	10
65	Changes in correlation characteristics of time consumption and mind-reading performance in pre-onset and post-onset psychosis. <i>Psychiatry Research</i> , 2018, 262, 168-174.	3.3	9
66	Isolated hallucination is less predictive than thought disorder in psychosis: Insight from a longitudinal study in a clinical population at high risk for psychosis. <i>Scientific Reports</i> , 2018, 8, 13962.	3.3	9
67	Neurological soft signs and grey matter abnormalities in individuals with ultraâ€”high risk for psychosis. <i>PsyCh Journal</i> , 2019, 8, 252-260.	1.1	9
68	Poor functional recovery is better predicted than conversion in studies of outcomes of clinical high risk of psychosis: insight from SHARP. <i>Psychological Medicine</i> , 2020, 50, 1578-1584.	4.5	9
69	Predicting atypical visual saliency for autism spectrum disorder via scale-adaptive inception module and discriminative region enhancement loss. <i>Neurocomputing</i> , 2021, 453, 610-622.	5.9	9
70	Antipsychotics Effects on Network-Level Reconfiguration of Cortical Morphometry in First-Episode Schizophrenia. <i>Schizophrenia Bulletin</i> , 2022, 48, 231-240.	4.3	9
71	Further evidence that antipsychotic medication does not prevent long-term psychosis in higher-risk individuals. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2022, 272, 591-602.	3.2	9
72	Identification and prediction of clinical high risk of psychosis in Chinese outpatients using two-stage screening. <i>Schizophrenia Research</i> , 2018, 202, 284-290.	2.0	8

#	ARTICLE	IF	CITATIONS
73	Parietal memory network and default mode network in first-episode drug-naïve schizophrenia: Associations with auditory hallucination. <i>Human Brain Mapping</i> , 2020, 41, 1973-1984.	3.6	8
74	When to initiate antipsychotic treatment for psychotic symptoms: At the premorbid phase or first episode of psychosis?. <i>Australian and New Zealand Journal of Psychiatry</i> , 2021, 55, 314-323.	2.3	8
75	The role of circulating blood microRNA-374 and microRNA-10 levels in the pathogenesis and therapeutic mechanisms of major depressive disorder. <i>Neuroscience Letters</i> , 2021, 763, 136184.	2.1	8
76	An Open-label Trial of Adjuvant High-frequency Left Prefrontal Repetitive Transcranial Magnetic Stimulation for Treating Suicidal Ideation in Adolescents and Adults With Depression. <i>Journal of ECT</i> , 2021, 37, 140-146.	0.6	8
77	Different levels of facial expression recognition in patients with first-episode schizophrenia: A functional MRI study. <i>Annals of General Psychiatry</i> , 2018, 31, e000014.	3.1	7
78	Precise theta burst transcranial magnetic stimulation selectively reduced duration-related mismatch negativity. <i>Biological Psychology</i> , 2018, 137, 125-132.	2.2	7
79	25 Hz Magnetic Seizure Therapy Is Feasible but Not Optimal for Chinese Patients With Schizophrenia: A Case Series. <i>Frontiers in Psychiatry</i> , 2018, 9, 224.	2.6	7
80	Detection Study of Bipolar Depression Through the Application of a Model-Based Algorithm in Terms of Clinical Feature and Peripheral Biomarkers. <i>Frontiers in Psychiatry</i> , 2019, 10, 266.	2.6	7
81	A Strategy to Address High Comorbidity of Personality Disorders in a Chinese Population: A Principal and Subordinate Diagnostic Model. <i>Psychiatry (New York)</i> , 2019, 82, 1-11.	0.7	7
82	MK-Curve improves sensitivity to identify white matter alterations in clinical high risk for psychosis. <i>NeuroImage</i> , 2021, 226, 117564.	4.2	7
83	Imbalance Model of Heart Rate Variability and Pulse Wave Velocity in Psychotic and Nonpsychotic Disorders. <i>Schizophrenia Bulletin</i> , 2021, , .	4.3	7
84	Plasma metabolic alterations and potential biomarkers in individuals at clinical high risk for psychosis. <i>Schizophrenia Research</i> , 2022, 239, 19-28.	2.0	7
85	The D2R-DISC1 protein complex and associated proteins are altered in schizophrenia and normalized with antipsychotic treatment. <i>Journal of Psychiatry and Neuroscience</i> , 2022, 47, E134-E147.	2.4	7
86	Effective differentiation between depressed patients and controls using discriminative eye movement features. <i>Journal of Affective Disorders</i> , 2022, 307, 237-243.	4.1	7
87	Elevated hippocampal choline level is associated with altered functional connectivity in females with major depressive disorder: A pilot study. <i>Psychiatry Research - Neuroimaging</i> , 2018, 278, 48-55.	1.8	6
88	Effect of Adjunct Repetitive Transcranial Magnetic Stimulation in Elderly Patients with Acute Depressive Episode: Supporting Evidence from a Real-World Observation. <i>American Journal of Geriatric Psychiatry</i> , 2019, 27, 91-92.	1.2	6
89	Decreasing risk of psychosis by sulforaphane study protocol for a randomized, double-blind, placebo-controlled, clinical multi-centre trial. <i>Microbial Biotechnology</i> , 2021, 15, 585-594.	1.7	6
90	Neurocognitive Assessments Are More Important Among Adolescents Than Adults for Predicting Psychosis in Clinical High Risk. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2022, 7, 56-65.	1.5	6

#	ARTICLE	IF	CITATIONS
91	Temporal and timeâ€“frequency features of auditory oddball response in distinct subtypes of patients at clinical high risk for psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2022, 272, 449-459.	3.2	6
92	Biological mechanisms and clinical efficacy of sulforaphane for mental disorders. <i>Annals of General Psychiatry</i> , 2022, 35, e100700.	3.1	6
93	Frequency of narcissistic personality disorder in a counseling center population in China. <i>BMC Psychiatry</i> , 2019, 19, 212.	2.6	5
94	Screening of the college students at clinical high risk for psychosis in China: a multicenter epidemiological study. <i>BMC Psychiatry</i> , 2021, 21, 253.	2.6	5
95	Identify autism spectrum disorder via dynamic filter and deep spatiotemporal feature extraction. <i>Signal Processing: Image Communication</i> , 2021, 94, 116195.	3.2	5
96	Childhood Adversities in Narcissistic Personality Disorder in China. <i>Psychiatry (New York)</i> , 2021, 84, 81-83.	0.7	5
97	Magnetic Seizure Therapy Compared to Electroconvulsive Therapy for Schizophrenia: A Randomized Controlled Trial. <i>Frontiers in Psychiatry</i> , 2021, 12, 770647.	2.6	5
98	Antipsychotic prescription, assumption and conversion to psychosis: resolving missing clinical links to optimize prevention through precision. <i>NPJ Schizophrenia</i> , 2022, 8, .	3.6	5
99	Attenuated niacin-induced skin flush response in individuals with clinical high risk for psychosis. <i>Annals of General Psychiatry</i> , 2022, 35, e100748.	3.1	5
100	Comorbidity of personality disorder in obsessive-compulsive disorder: special emphases on the clinical significance. <i>CNS Spectrums</i> , 2015, 20, 466-468.	1.2	4
101	Time consumption in mind-reading: A potentially important factor for social cognition assessment in early psychosis. <i>Schizophrenia Research</i> , 2018, 192, 491-492.	2.0	4
102	<p>>Real-World Effectiveness and Safety of Antipsychotics in Individuals at Clinical High-Risk for Psychosis: Study Protocol for a Prospective Observational Study (ShangHai at Risk for) Tj ETQq0 0 0 rgBT /Overlock210 Tf 50 497 Td (Psy	2.1	4
103	Effect of DAOA genetic variation on white matter alteration in corpus callosum in patients with first-episode schizophrenia. <i>Brain Imaging and Behavior</i> , 2021, 15, 1748-1759.	2.1	4
104	Antidepressant Effect of Adjunct Repetitive Transcranial Magnetic Stimulation in Inpatients 60 Years and Older. <i>Journal of ECT</i> , 2020, 36, 216-221.	0.6	4
105	Subtypes of Clinical High Risk for Psychosis that Predict Antipsychotic Effectiveness in Long-Term Remission. <i>Pharmacopsychiatry</i> , 2021, 54, 23-30.	3.3	4
106	The effects of antipsychotics on interactions of dynamic functional connectivity in the triple-network in first episode schizophrenia. <i>Schizophrenia Research</i> , 2021, 236, 29-37.	2.0	4
107	Relationship Between Cognitive and Clinical Insight at Different Durations of Untreated Attenuated Psychotic Symptoms in High-Risk Individuals. <i>Frontiers in Psychiatry</i> , 2021, 12, 753130.	2.6	4
108	Effects of polygenic risk of schizophrenia on interhemispheric callosal white matter integrity and frontotemporal functional connectivity in first-episode schizophrenia. <i>Psychological Medicine</i> , 2022, , 1-10.	4.5	4

#	ARTICLE	IF	CITATIONS
109	Antipsychotic Exposure in Clinical High Risk of Psychosis. <i>Journal of Clinical Psychiatry</i> , 2022, 83, .	2.2	4
110	Comparison of electroconvulsive therapy and magnetic seizure therapy in schizophrenia: Structural changes/neuroplasticity. <i>Psychiatry Research</i> , 2022, 312, 114523.	3.3	4
111	Automatic auditory processing features in distinct subtypes of patients at clinical high risk for psychosis: Forecasting remission with mismatch negativity. <i>Human Brain Mapping</i> , 2022, 43, 5452-5464.	3.6	4
112	A compromise solution between overlapping and overlooking DSM personality disorders in Chinese psychiatric practice. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2018, 53, 99-106.	3.1	3
113	Gender differences in screening self-reported psychotic symptoms in a first help-seeking population. <i>Archives of Women's Mental Health</i> , 2022, 25, 291-299.	2.6	3
114	Psychosis risk syndrome is not prodromal psychosis. <i>Shanghai Archives of Psychiatry</i> , 2015, 27, 42-4.	0.7	3
115	Temporoparietal Connectivity Within Default Mode Network Associates With Clinical Improvements in Schizophrenia Following Modified Electroconvulsive Therapy. <i>Frontiers in Psychiatry</i> , 2021, 12, 768279.	2.6	3
116	Functional reconfiguration of cerebellum-cerebral neural loop in schizophrenia following electroconvulsive therapy. <i>Psychiatry Research - Neuroimaging</i> , 2022, 320, 111441.	1.8	3
117	Peripheral transcriptome of clinical high-risk psychosis reflects symptom alteration and helps prognosis prediction. <i>Psychiatry and Clinical Neurosciences</i> , 2022, 76, 268-270.	1.8	3
118	Reduced temporal activation during a verbal fluency test in clinical high risk of psychosis: a functional near-infrared spectroscopy-based study. <i>Annals of General Psychiatry</i> , 2022, 35, e100702.	3.1	3
119	Effect of cognitive insight on clinical insight from pre-morbid to early psychosis stages. <i>Psychiatry Research</i> , 2022, 313, 114613.	3.3	3
120	Relationships between self-reflectiveness and clinical symptoms in individuals during pre-morbid and early clinical stages of psychosis. <i>Annals of General Psychiatry</i> , 2022, 35, e100696.	3.1	3
121	Replication of the abnormal niacin response in first episode psychosis measured using laser Doppler flowmeter. <i>Asia-Pacific Psychiatry</i> , 2022, 14, .	2.2	3
122	A synopsis of recent influential papers published in mental health journals (2012â€“2013) in Mainland China. <i>Asian Journal of Psychiatry</i> , 2014, 10, 105-108.	2.0	2
123	Frequency of Self-reported Psychotic Symptoms among 2542 Outpatients at Their First Visit for Mental Health Services. <i>Psychiatry (New York)</i> , 2021, 84, 57-67.	0.7	2
124	A Preliminary Investigation on Plasma Cell Adhesion Molecules Levels by Protein Microarray Technology in Major Depressive Disorder. <i>Frontiers in Psychiatry</i> , 2021, 12, 627469.	2.6	2
125	Gender differences in attentive bias during social information processing in schizophrenia: An eye-tracking study. <i>Asian Journal of Psychiatry</i> , 2021, 66, 102871.	2.0	2
126	How should we intervene in psychosis risk syndromes?. <i>Shanghai Archives of Psychiatry</i> , 2013, 25, 6-9.	0.7	2

#	ARTICLE	IF	CITATIONS
127	Inefficient integration during multiple facial processing in pre-morbid and early phases of psychosis. World Journal of Biological Psychiatry, 2022, 23, 361-373.	2.6	2
128	Acute stress promotes brain oscillations and hippocampal-cortical dialog in emotional processing. Biochemical and Biophysical Research Communications, 2022, 598, 55-61.	2.1	2
129	Altered attentive bias towards interpersonal communication information across phases of schizophrenia: an eye-tracking study. Annals of General Psychiatry, 2022, 35, e100699.	3.1	2
130	Different patterns of association between white matter microstructure and plasma unsaturated fatty acids in those with high risk for psychosis and healthy participants. Annals of General Psychiatry, 2022, 35, e100703.	3.1	2
131	Artificial intelligence-assisted niacin skin flush screening in early psychosis identification and prediction. Annals of General Psychiatry, 2022, 35, e100753.	3.1	2
132	Abnormal neural oscillations in clinical high risk for psychosis: a magnetoencephalography method study. Annals of General Psychiatry, 2022, 35, e100712.	3.1	2
133	O10.5. ABNORMAL MODULAR ORGANIZATION OF THE FUNCTIONAL CONNECTOME PREDICTS CONVERSION TO PSYCHOSIS IN CLINICAL HIGH-RISK YOUTH. Schizophrenia Bulletin, 2018, 44, S104-S104.	4.3	1
134	Editorial: Identifying Individuals at Clinical High Risk of Psychosis in Different Cultures and Countries. Frontiers in Psychiatry, 2020, 11, 159.	2.6	1
135	A bibliometric analysis of international publications and citation trends of articles in mental health produced by Chinese institutions in mainland China (1990â€“2019). Global Mental Health (Cambridge,) Tj ETQq1 2020, 7, 1-14.	2.6	1
136	Individualized risk components guiding antipsychotic delivery in patients with a clinical high risk of psychosis: application of a risk calculator. Psychological Medicine, 2021, , 1-10.	4.5	1
137	Age-related changes in self-reported psychotic experiences in clinical help-seeking population: From 15 to 45 years. Microbial Biotechnology, 2022, 16, 1359-1367.	1.7	1
138	F14. REDUCED DURATION MISMATCH NEGATIVITY ASSOCIATED WITH DECREASED GLUTAMATE+GLUTAMINE LEVEL IN SUBJECTS AT CLINICAL HIGH-RISK FOR PSYCHOSIS. Schizophrenia Bulletin, 2018, 44, S223-S224.	4.3	0
139	O6.4. AUDITORY AND LANGUAGE AREAS DISTINGUISH CONVERTERS FROM NON-CONVERTERS AT BASELINE IN SHARP CLINICAL HIGH-RISK SUBJECTS FOR PSYCHOSIS STUDY. Schizophrenia Bulletin, 2018, 44, S90-S91.	4.3	0
140	S105. VALIDATING THE PREDICTIVE ACCURACY OF THE NAPLS-2 PSYCHOSIS RISK CALCULATOR IN A CLINICAL HIGH-RISK SAMPLE FROM THE SHARP (SHANGHAI AT RISK FOR PSYCHOSIS) PROGRAM. Schizophrenia Bulletin, 2018, 44, S366-S366.	4.3	0
141	21.4 BASELINE CLINICAL AND BIOLOGICAL VARIABLES PREDICTING 1 YEAR OUTCOME OF SUBJECTS AT CLINICAL HIGH RISK OF PSYCHOSIS: INSIGHT FROM SHANGHAI AT RISK FOR PSYCHOSIS (SHARP) PROGRAM. Schizophrenia Bulletin, 2018, 44, S36-S36.	4.3	0
142	S10. IMPAIRMENT OF REALITY TESTING IN INDIVIDUALS AT CLINICAL HIGH-RISK OF PSYCHOSIS. Schizophrenia Bulletin, 2019, 45, S309-S310.	4.3	0
143	S61. CLINICAL SUBTYPES THAT PREDICT CONVERSION TO PSYCHOSIS: A CANONICAL CORRELATION ANALYSIS STUDY FROM THE SHANGHAI AT RISK FOR PSYCHOSIS (SHARP) PROGRAM. Schizophrenia Bulletin, 2019, 45, S329-S330.	4.3	0
144	S69. NEURAL-SYNCHRONIZATION DURING COOPERATION TASK IN CLINICAL HIGH RISK OF PSYCHOSIS-A FNIRS-BASED HYPERSCANNING STUDY. Schizophrenia Bulletin, 2019, 45, S333-S333.	4.3	0

#	ARTICLE	IF	CITATIONS
145	S147. FUNCTIONAL BRAIN CONNECTIVITY DATA IMPROVE CLINICAL OUTCOME PREDICTION IN YOUTH AT RISK FOR PSYCHOSIS. Schizophrenia Bulletin, 2020, 46, S92-S92.	4.3	0
146	S171. EFFECT OF DAOA GENETIC VARIATION ON WHITE MATTER ALTERATION IN CORPUS CALLOSUM IN PATIENTS WITH FIRST-EPISODE SCHIZOPHRENIA. Schizophrenia Bulletin, 2020, 46, S102-S102.	4.3	0
147	Attenuated Psychosis Syndromes Among Youth and Young Adults in China: Early Identification and Intervention. , 2019, , 301-310.		0