

Lina Benson

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4977662/lina-benson-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

29
papers

587
citations

12
h-index

24
g-index

31
ext. papers

843
ext. citations

7.6
avg, IF

4.11
L-index

#	Paper	IF	Citations
29	Patient profile and outcomes associated with follow-up in specialty vs. primary care in heart failure.. <i>ESC Heart Failure</i> , 2022 ,	3.7	1
28	Use of evidence-based therapy in heart failure with reduced ejection fraction across age strata.. <i>European Journal of Heart Failure</i> , 2022 ,	12.3	4
27	Digoxin use in contemporary heart failure with reduced ejection fraction: an analysis from the Swedish Heart Failure Registry.. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2021 ,	6.4	2
26	A comprehensive characterization of acute heart failure with preserved vs. mildly reduced vs. reduced ejection fraction - insights from the ESC-HFA EORP Heart Failure Long-Term Registry.. <i>European Journal of Heart Failure</i> , 2021 ,	12.3	4
25	Eligibility of patients with heart failure with preserved ejection fraction for sacubitril/valsartan according to the PARAGON-HF trial. <i>ESC Heart Failure</i> , 2021 , 9, 164	3.7	1
24	Use of sodium-glucose co-transporter 2 inhibitors in patients with heart failure and type 2 diabetes mellitus: data from the Swedish Heart Failure Registry. <i>European Journal of Heart Failure</i> , 2021 , 23, 1012-1022	12.3	10
23	Association Between β Blockers and Outcomes in Heart Failure With Preserved Ejection Fraction: Current Insights From the SwedeHF Registry. <i>Journal of Cardiac Failure</i> , 2021 , 27, 1165-1174	3.3	1
22	Lower socioeconomic status predicts higher mortality and morbidity in patients with heart failure. <i>Heart</i> , 2021 , 107, 229-236	5.1	8
21	Association between renin-angiotensin-aldosterone system inhibitor use and COVID-19 hospitalization and death: a 1.4 million patient nationwide registry analysis. <i>European Journal of Heart Failure</i> , 2021 , 23, 476-485	12.3	24
20	Cardiac resynchronization therapy with or without defibrillator in patients with heart failure. <i>Europace</i> , 2021 ,	3.9	1
19	Non-cardiology vs. cardiology care of patients with heart failure and reduced ejection fraction is associated with lower use of guideline-based care and higher mortality: Observations from The Swedish Heart Failure Registry. <i>International Journal of Cardiology</i> , 2021 , 343, 63-72	3.2	1
18	A population-based study of post-endovascular aortic repair rupture during 15 years. <i>Journal of Vascular Surgery</i> , 2021 , 74, 701-710.e3	3.5	2
17	Cryoablation of atypical atrioventricular nodal reentry tachycardia. <i>Journal of Cardiovascular Electrophysiology</i> , 2021 , 32, 2971-2978	2.7	2
16	Phenotyping heart failure patients for iron deficiency and use of intravenous iron therapy: data from the Swedish Heart Failure Registry. <i>European Journal of Heart Failure</i> , 2021 , 23, 1844-1854	12.3	8
15	Association between potassium level and outcomes in heart failure with reduced ejection fraction: a cohort study from the Swedish Heart Failure Registry. <i>European Journal of Heart Failure</i> , 2020 , 22, 1390-1398	12.3	15
14	Association between beta-blocker use and mortality/morbidity in older patients with heart failure with reduced ejection fraction. A propensity score-matched analysis from the Swedish Heart Failure Registry. <i>European Journal of Heart Failure</i> , 2020 , 22, 103-112	12.3	15
13	Long-term outcome of patients with invasive electrophysiology procedure-related cardiac tamponade. <i>Europace</i> , 2020 , 22, 1547-1557	3.9	1

12	Heart Failure in Patients Undergoing Elective and Emergency Noncardiac Surgery: Still a Poorly Addressed Risk Factor. <i>Journal of Cardiac Failure</i> , 2020 , 26, 1034-1042	3.3	1
11	Association of Coronary Microvascular Dysfunction With Heart Failure Hospitalizations and Mortality in Heart Failure With Preserved Ejection Fraction: A Follow-up in the PROMIS-HFpEF Study. <i>Journal of Cardiac Failure</i> , 2020 , 26, 1016-1021	3.3	9
10	Association Between Use of Primary-Prevention Implantable Cardioverter-Defibrillators and Mortality in Patients With Heart Failure: A Prospective Propensity Score-Matched Analysis From the Swedish Heart Failure Registry. <i>Circulation</i> , 2019 , 140, 1530-1539	16.7	41
9	Alcohol-related hospitalization is associated with increased risk of ischaemic stroke among low-risk patients with atrial fibrillation. <i>Europace</i> , 2018 , 20, 19-24	3.9	4
8	New York Heart Association functional class, QRS duration, and survival in heart failure with reduced ejection fraction: implications for cardiac resynchronization therapy. <i>European Journal of Heart Failure</i> , 2017 , 19, 366-376	12.3	21
7	Association between demographic, organizational, clinical, and socio-economic characteristics and underutilization of cardiac resynchronization therapy: results from the Swedish Heart Failure Registry. <i>European Journal of Heart Failure</i> , 2017 , 19, 1270-1279	12.3	54
6	Significance of Ischemic Heart Disease in Patients With Heart Failure and Preserved, Midrange, and Reduced Ejection Fraction: A Nationwide Cohort Study. <i>Circulation: Heart Failure</i> , 2017 , 10,	7.6	104
5	Association Between Use of Long-Acting Nitrates and Outcomes in Heart Failure With Preserved Ejection Fraction. <i>Circulation: Heart Failure</i> , 2017 , 10,	7.6	13
4	Association Between Use of Statins and Mortality in Patients With Heart Failure and Ejection Fraction of ≥ 50 . <i>Circulation: Heart Failure</i> , 2015 , 8, 862-70	7.6	64
3	Gender, underutilization of cardiac resynchronization therapy, and prognostic impact of QRS prolongation and left bundle branch block in heart failure. <i>Europace</i> , 2015 , 17, 424-31	3.9	39
2	Association between use of statins and outcomes in heart failure with reduced ejection fraction: prospective propensity score matched cohort study of 21 864 patients in the Swedish Heart Failure Registry. <i>Circulation: Heart Failure</i> , 2015 , 8, 252-60	7.6	30
1	Association between use of β -blockers and outcomes in patients with heart failure and preserved ejection fraction. <i>JAMA - Journal of the American Medical Association</i> , 2014 , 312, 2008-18	27.4	106