

# Sami H Rabei

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/498879/publications.pdf>

Version: 2024-02-01

10  
papers

64  
citations

1937685

4  
h-index

1720034

7  
g-index

10  
all docs

10  
docs citations

10  
times ranked

98  
citing authors

#	ARTICLE	IF	CITATIONS
1	The complementary role of root and leaf PIP1 and PIP2 aquaporins drives the anisohydric behavior in <i>Helinathus annuus</i> L.. <i>Environmental and Experimental Botany</i> , 2021, 182, 104314.	4.2	2
2	Floristic study of Saint Katherine protectorate, Sinai: with one new record to flora of Egypt. <i>Taeckholmia</i> , 2021, 41, 32-55.	0.3	5
3	Phenotypic Diversity of <i>Eragrostis Wolf</i> in Egypt. <i>Taeckholmia</i> , 2021, 41, 18-31.	0.3	0
4	Effect of Genetic Diversity on the Distribution of Endemic Species of the Genus <i>Silene</i> (Caryophyllaceae) in Saint Katherine Protectorate, Sinai, Egypt. <i>Plant Molecular Biology Reporter</i> , 2020, 39, 364.	1.8	1
5	(2728) Proposal to conserve the name <i>Datura innoxia</i> (<i>Solanaceae</i>) with that spelling. <i>Taxon</i> , 2019, 68, 1370-1371.	0.7	1
6	<i>Clinopodium serpyllifolium</i> subsp. <i>barbatum</i> (Lamiaceae) and <i>Datura ferox</i> , (Solanaceae) new records for the flora of Egypt. <i>Egyptian Journal of Botany</i> , 2018, .	0.2	0
7	Water use efficiency in the drought-stressed sorghum and maize in relation to expression of aquaporin genes. <i>Biologia Plantarum</i> , 2017, 61, 127-137.	1.9	37
8	Keys for and morphological character variation in some Egyptian cultivars of Cucurbitaceae. <i>Genetic Resources and Crop Evolution</i> , 2013, 60, 1353-1364.	1.6	9
9	Conventional keys for Convolvulaceae in the flora of Egypt. <i>Flora Mediterranea</i> , 2012, 22, 45-62.	0.1	2
10	Taxonomic Assessment of Five Numerical Methods and its Implications on the Classification of Hyptis s.l. (Labiatae). <i>International Journal of Botany</i> , 2007, 4, 85-92.	0.2	7