Lori R Covey

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4993503/publications.pdf Version: 2024-02-01



LODI P COVEY

#	Article	IF	CITATIONS
1	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). Autophagy, 2016, 12, 1-222.	9.1	4,701
2	A Complex Containing Polypyrimidine Tract-Binding Protein Is Involved in Regulating the Stability of CD40 Ligand (CD154) mRNA. Journal of Immunology, 2003, 170, 979-988.	0.8	45
3	CD40:CD40L Interactions in X-linked and Non-X-linked Hyper-IgM Syndromes. Immunologic Research, 2001, 24, 311-324.	2.9	36
4	Identification of a Complex that Binds to the CD154 3′ Untranslated Region: Implications for a Role in Message Stability During T Cell Activation. Journal of Immunology, 2000, 165, 4478-4486.	0.8	30
5	Post-transcriptional regulation in Lymphocytes: The case of CD154. RNA Biology, 2009, 6, 259-265.	3.1	25
6	The RNA-Binding Protein, Polypyrimidine Tract-Binding Protein 1 (PTBP1) Is a Key Regulator of CD4 T Cell Activation. PLoS ONE, 2016, 11, e0158708.	2.5	25
7	Isolation of cDNAS encoding T-BAM, a surface glycoprotein on CD4+ T cells mediating contact-dependent helper function for B cells: Identity with the CD40-ligand. Molecular Immunology, 1994, 31, 471-484.	2.2	24
8	A Polypyrimidine Tract-Binding Protein-Dependent Pathway of mRNA Stability Initiates with CpG Activation of Primary B Cells. Journal of Immunology, 2008, 181, 3336-3345.	0.8	20
9	<i>In vivo</i> postâ€transcriptional regulation of CD154 in mouse CD4 ⁺ T cells. European Journal of Immunology, 2009, 39, 2224-2232.	2.9	18
10	Polypyrimidine Tract-Binding Protein Is Critical for the Turnover and Subcellular Distribution of CD40 Ligand mRNA in CD4+ T Cells. Journal of Immunology, 2011, 186, 2164-2171.	0.8	17
11	CREB / ATF proteins enhance the basal and CD154- and IL-4-induced transcriptional activity of the human lγ1 proximal promoter. European Journal of Immunology, 2001, 31, 653-664.	2.9	10
12	Functional analysis of a tripartite stability element within the CD40 ligand 3' untranslated region. Immunology, 2008, 124, 368-379.	4.4	8
13	c-Rel plays a key role in deficient activation of B cells from a non–X-linked hyper-IgM patient. Blood, 2006, 108, 3769-3776.	1.4	5
14	c-Rel Deficiency Increases Caspase-4 Expression and Leads to ER Stress and Necrosis in EBV-Transformed Cells. PLoS ONE, 2011, 6, e25467.	2.5	5
15	Maintenance of the CD40-related immunodeficient response in hyper-IgM B cells immortalized with a LMP1-regulated mini-EBV. Journal of Leukocyte Biology, 2005, 78, 620-629.	3.3	4
16	A Posttranscriptional Pathway of CD40 Ligand mRNA Stability Is Required for the Development of an Optimal Humoral Immune Response. Journal of Immunology, 2021, 206, 2552-2565.	0.8	2
17	Identification of a novel form of câ€Rel in a human B cell line undergoing spontaneous apoptosis. FASEB Journal, 2008, 22, 1066.17.	0.5	0