## Mauricio Salinas-Santander

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5000035/publications.pdf

Version: 2024-02-01

1039880 996849 30 265 9 15 citations g-index h-index papers 30 30 30 452 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Incidence of BRCA1 and BRCA2 mutations in 54 Chilean families with breast/ovarian cancer, genotype–phenotype correlations. Breast Cancer Research and Treatment, 2006, 95, 81-87.	1.1	58
2	Association between PTPN22 C1858T polymorphism and alopecia areata risk. Experimental and Therapeutic Medicine, 2015, 10, 1953-1958.	0.8	25
3	Protein tyrosine phosphatase PTPN22 +1858C/T polymorphism is associated with active vitiligo. Experimental and Therapeutic Medicine, 2014, 8, 1433-1437.	0.8	18
4	Updates and new medical treatments for vitiligo (Review). Experimental and Therapeutic Medicine, 2021, 22, 797.	0.8	18
5	Novel immunological and genetic factors associated with vitiligo: A review. Experimental and Therapeutic Medicine, 2021, 21, 312.	0.8	17
6	Tumor necrosis factor- $\hat{l}_{\pm}$ -308G/A polymorphism is associated with active vitiligo vulgaris in a northeastern Mexican population. Experimental and Therapeutic Medicine, 2012, 3, 893-897.	0.8	16
7	CAPN3, DCT, MLANA and TYRP1 are overexpressed in skin of vitiligo vulgaris Mexican patients. Experimental and Therapeutic Medicine, 2018, 15, 2804-2811.	0.8	14
8	Uridine 5'â€'diphosphoâ€'glucronosyltrasferase: Its role in pharmacogenomics and human disease (Review). Experimental and Therapeutic Medicine, 2018, 16, 3-11.	0.8	14
9	Evaluation of skin expression profiles of patients with vitiligo treated with narrow-band UVB therapy by targeted RNA-seq. Anais Brasileiros De Dermatologia, 2018, 93, 843-851.	0.5	11
10	Tumor necrosis factor alpha promoterâ€308G/A polymorphism in Mexican patients with patchy alopecia areata. International Journal of Dermatology, 2012, 51, 571-575.	0.5	10
11	Evaluation of the Expression of Genes Associated with Inflammation and Apoptosis in Androgenetic Alopecia by Targeted RNA-Seq. Skin Appendage Disorders, 2018, 4, 268-273.	0.5	10
12	High prevalence of diabetic retinopathy and lack of association with integrin $\hat{l}\pm 2$ gene polymorphisms in patients with type 2 diabetes from Northeastern Mexico. Experimental and Therapeutic Medicine, 2015, 10, 435-444.	0.8	8
13	Inhibition of miR31 and miR92a as Oncological Biomarkers in RKO Colon Cancer Cells Treated with Kaempferol-3- <i>O</i> -Glycoside Isolated from Black Bean. Journal of Medicinal Food, 2020, 23, 50-55.	0.8	8
14	CTLA4 +49AG (rs231775) and CT60 (rs3087243) gene variants are not associated with alopecia areata in a Mexican population from Monterrey Mexico. Anais Brasileiros De Dermatologia, 2020, 95, 283-288.	0.5	6
15	Association of the Vascular Endothelial Growth Factor Gene PolymorphismÂ+936ÂC/T with Diabetic Neuropathy in Patients with Type 2 Diabetes Mellitus. Archives of Medical Research, 2019, 50, 181-186.	1.5	5
16	Expression of melanocortin 1 receptor before and after narrowband UVB phototherapy treatment in patients with stable vitiligo: A prospective study. Experimental and Therapeutic Medicine, 2020, 19, 1649-1654.	0.8	5
17	ACE, APOA5, and MTP Gene Polymorphisms Analysis in Relation to Triglyceride and Insulin Levels in Pediatric Patients. Archives of Medical Research, 2018, 49, 94-100.	1.5	4
18	Depressive Symptoms are Associated with low Serotonin Levels in Plasma but are not 5–HTTLPR Genotype Dependent in Older Adults. Spanish Journal of Psychology, 2021, 24, e28.	1.1	4

#	Article	IF	CITATIONS
19	TheÂ+1858ÂC/T Polymorphism in the PTPN22 Gene Is Associated with Cystic Fibrosis Patients in Northeast Mexico. Archives of Medical Research, 2016, 47, 403-406.	1.5	3
20	Rapid Detection of the GSTM3 A/B Polymorphism Using Real-time PCR with TaqMan® Probes. Archives of Medical Research, 2016, 47, 142-145.	1.5	3
21	Purshia plicata Triggers and Regulates Proteins Related to Apoptosis in HeLa Cancer Cells. Plants, 2021, 10, 2559.	1.6	3
22	HPV-16 E6 and E7 oncogene expression is downregulated as a result of Mdm2 knockdown. International Journal of Oncology, 2012, 41, 141-6.	1.4	2
23	Assessment of biochemical parameters and characterization of TNFα â^308G/A and PTPN22 +1858C/T gene polymorphisms in the risk of obesity in adolescents. Biomedical Reports, 2016, 4, 107-111.	0.9	2
24	Thymidylate synthase polymorphism in Mexican patients with colon cancer treated with 5-fluorouracil. Journal of B U on, 2016, 21, 935-940.	0.4	1
25	Distribution of the ATP-binding cassette transporter ABCG8 IVS1-2A>G genotype and clinical characteristics of gallbladder patients in Northeastern Mexico: A pilot study. Biomedical Reports, 2018, 9, 266-270.	0.9	0
26	TSH levels, overweight, BMI, and skin expression levels of DCT and CCBL2 genes are related to vitiligo treatment response with narrow band UVB phototherapy. Dermatologic Therapy, 2019, 32, e12893.	0.8	0
27	Pharmacology Evaluation of Bioactive Compounds that Regulate Cervical Cancer Cells. , 2019, , .		0
28	Novel polygenetic variants evidenced in a patient with Jervell and Lange-Nielsen syndrome. Cardiology Journal, 2021, 28, 786-789.	0.5	0
29	Punicalin and ellagic acid from pomegranate peel extract facilitate apoptotic behaviour in the Hela cell line Pakistan Journal of Pharmaceutical Sciences, 2021, 34, 2181-2189.	0.2	0
30	Transcriptional Profile of HCV Replicon Cells after Treatment with Acetylsalicylic Acid Annals of Clinical and Laboratory Science, 2022, 52, 222-229.	0.2	0