

Yufen Xu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5001217/publications.pdf>

Version: 2024-02-01

28
papers

201
citations

1162367

8
h-index

1199166

12
g-index

30
all docs

30
docs citations

30
times ranked

188
citing authors

#	ARTICLE	IF	CITATIONS
1	The IL-1B Gene Polymorphisms rs16944 and rs1143627 Contribute to an Increased Risk of Coronary Artery Lesions in Southern Chinese Children with Kawasaki Disease. <i>Journal of Immunology Research</i> , 2019, 2019, 1-7.	0.9	24
2	The lncRNA MALAT1 rs619586 G Variant Confers Decreased Susceptibility to Recurrent Miscarriage. <i>Frontiers in Physiology</i> , 2019, 10, 385.	1.3	24
3	The lncRNA <i>CCAT2</i> rs6983267 G allele is associated with decreased susceptibility to recurrent miscarriage. <i>Journal of Cellular Physiology</i> , 2019, 234, 20577-20583.	2.0	16
4	A <i>PEAR1</i> polymorphism (<i>rs12041331</i>) is associated with risk of coronary artery aneurysm in Kawasaki disease. <i>Annals of Human Genetics</i> , 2019, 83, 54-62.	0.3	12
5	The rs1625579 T>G polymorphism in the <i>miRNA-137</i> gene confers a risk of early-onset Kawasaki disease in a southern Chinese population. <i>Infection and Drug Resistance</i> , 2018, Volume 11, 1055-1060.	1.1	11
6	LncRNA HULC Polymorphism Is Associated With Recurrent Spontaneous Abortion Susceptibility in the Southern Chinese Population. <i>Frontiers in Genetics</i> , 2019, 10, 918.	1.1	10
7	P2RY12:rs7637803 TT variant genotype increases coronary artery aneurysm risk in Kawasaki disease in a southern Chinese population. <i>Journal of Gene Medicine</i> , 2019, 21, e3066.	1.4	10
8	ABCC4 Variants Modify Susceptibility to Kawasaki Disease in a Southern Chinese Population. <i>Disease Markers</i> , 2018, 2018, 1-7.	0.6	9
9	TBXA2R rs4523 G allele is associated with decreased susceptibility to Kawasaki disease. <i>Cytokine</i> , 2018, 111, 216-221.	1.4	9
10	The association between the <i>miR-146a</i> rs2910164 C>G polymorphism and Kawasaki disease in a southern Chinese population. <i>Bioscience Reports</i> , 2018, 38, .	1.1	9
11	An Angiotensinogen Gene Polymorphism (rs5050) Is Associated with the Risk of Coronary Artery Aneurysm in Southern Chinese Children with Kawasaki Disease. <i>Disease Markers</i> , 2019, 2019, 1-7.	0.6	9
12	Association between P2RY12 Gene Polymorphisms and IVIG Resistance in Kawasaki Patients. <i>Cardiovascular Therapeutics</i> , 2020, 2020, 1-6.	1.1	9
13	Association between the rs2288947 polymorphism of the lncRNA <i>TINCR</i> gene and the risk of recurrent miscarriage in a Southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22919.	0.9	7
14	Single-Nucleotide Polymorphism LncRNA AC008392.1/rs7248320 in CARD8 is Associated with Kawasaki Disease Susceptibility in the Han Chinese Population. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 4809-4816.	1.6	6
15	Lack of association between <i>miR-218</i> rs11134527 A>G and Kawasaki disease susceptibility. <i>Bioscience Reports</i> , 2018, 38, .	1.1	5
16	FNDC1 Polymorphism (rs3003174 C > T) Increased the Incidence of Coronary Artery Aneurysm in Patients with Kawasaki Disease in a Southern Chinese Population. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 2633-2640.	1.6	5
17	Homozygous of MRP4 Gene rs1751034 C Allele Is Related to Increased Risk of Intravenous Immunoglobulin Resistance in Kawasaki Disease. <i>Frontiers in Genetics</i> , 2021, 12, 510350.	1.1	4
18	The <i>miRNA-608</i> rs4919510 G>C polymorphism confers reduce coronary injury of Kawasaki disease in a Southern Chinese population. <i>Bioscience Reports</i> , 2019, 39, .	1.1	3

#	ARTICLE	IF	CITATIONS
19	The rs1051931 G>A Polymorphism in the PLA2G7 Gene Confers Resistance to Immunoglobulin Therapy in Kawasaki Disease in a Southern Chinese Population. <i>Frontiers in Pediatrics</i> , 2020, 8, 338.	0.9	3
20	The lncRNA ANRIL Gene rs2151280 GG Genotype is Associated with Increased Susceptibility to Recurrent Miscarriage in a Southern Chinese Population. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 2865-2872.	1.6	3
21	Association study of miR-149, miR-196a2, and miR-499a polymorphisms with coronary artery aneurysm of Kawasaki disease in southern Chinese population. <i>Journal of Gene Medicine</i> , 2022, 24, e3405.	1.4	3
22	Association between the TOX3 rs3803662 C>T polymorphism and recurrent miscarriage in a southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22992.	0.9	2
23	Integrin β 2 gene polymorphism is a risk factor of coronary artery lesions in Chinese children with Kawasaki disease. <i>Pediatric Rheumatology</i> , 2021, 19, 12.	0.9	2
24	The lncRNA CCAT2 Rs6983267 G Variant Contributes to Increased Sepsis Susceptibility in a Southern Chinese Population. <i>Infection and Drug Resistance</i> , 2021, Volume 14, 2969-2976.	1.1	2
25	The SERPINA4 rs2070777 AA Genotype is Associated with an Increased Risk of Recurrent Miscarriage in a Southern Chinese Population. <i>International Journal of Women's Health</i> , 2021, Volume 13, 111-117.	1.1	2
26	Association between the rs3802201 polymorphism of the lncRNA MIR2052HG gene and the risk of recurrent miscarriage in a Southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2022, 36, e24167.	0.9	2
27	Protective Effect of TNFRSF11A rs7239667 G > C Gene Polymorphism on Coronary Outcome of Kawasaki Disease in Southern Chinese Population. <i>Frontiers in Genetics</i> , 2021, 12, 691282.	1.1	0
28	The rs7404339 AA Genotype in CDH5 Contributes to Increased Risks of Kawasaki Disease and Coronary Artery Lesions in a Southern Chinese Child Population. <i>Frontiers in Cardiovascular Medicine</i> , 2022, 9, 760982.	1.1	0