

Yufen Xu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5001217/publications.pdf>

Version: 2024-02-01

28
papers

201
citations

1163117
8
h-index

1199594
12
g-index

30
all docs

30
docs citations

30
times ranked

188
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Association study of miR-149, miR-196a2, and miR-499a polymorphisms with coronary artery aneurysm of Kawasaki disease in southern Chinese population. <i>Journal of Gene Medicine</i> , 2022, 24, e3405. | 2.8 | 3 |
| 2 | Association between the rs3802201 polymorphism of the lncRNA MIR2052HG gene and the risk of recurrent miscarriage in a Southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2022, 36, e24167. | 2.1 | 2 |
| 3 | The rs7404339 AA Genotype in CDH5 Contributes to Increased Risks of Kawasaki Disease and Coronary Artery Lesions in a Southern Chinese Child Population. <i>Frontiers in Cardiovascular Medicine</i> , 2022, 9, 760982. | 2.4 | 0 |
| 4 | Integrin $\beta 2$ gene polymorphism is a risk factor of coronary artery lesions in Chinese children with Kawasaki disease. <i>Pediatric Rheumatology</i> , 2021, 19, 12. | 2.1 | 2 |
| 5 | Homozygous of MRP4 Gene rs1751034 C Allele Is Related to Increased Risk of Intravenous Immunoglobulin Resistance in Kawasaki Disease. <i>Frontiers in Genetics</i> , 2021, 12, 510350. | 2.3 | 4 |
| 6 | The lncRNA ANRIL Gene rs2151280 GG Genotype is Associated with Increased Susceptibility to Recurrent Miscarriage in a Southern Chinese Population. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 2865-2872. | 3.5 | 3 |
| 7 | FND1C1 Polymorphism (rs3003174 C > T) Increased the Incidence of Coronary Artery Aneurysm in Patients with Kawasaki Disease in a Southern Chinese Population. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 2633-2640. | 3.5 | 5 |
| 8 | The lncRNA CCAT2 Rs6983267 G Variant Contributes to Increased Sepsis Susceptibility in a Southern Chinese Population. <i>Infection and Drug Resistance</i> , 2021, Volume 14, 2969-2976. | 2.7 | 2 |
| 9 | Protective Effect of TNFRSF11A rs7239667 G > C Gene Polymorphism on Coronary Outcome of Kawasaki Disease in Southern Chinese Population. <i>Frontiers in Genetics</i> , 2021, 12, 691282. | 2.3 | 0 |
| 10 | Single-Nucleotide Polymorphism lncRNA AC008392.1/rs7248320 in CARD8 is Associated with Kawasaki Disease Susceptibility in the Han Chinese Population. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 4809-4816. | 3.5 | 6 |
| 11 | The SERPINA4 rs2070777 AA Genotype is Associated with an Increased Risk of Recurrent Miscarriage in a Southern Chinese Population. <i>International Journal of Women's Health</i> , 2021, Volume 13, 111-117. | 2.6 | 2 |
| 12 | Association between P2RY12 Gene Polymorphisms and IVIG Resistance in Kawasaki Patients. <i>Cardiovascular Therapeutics</i> , 2020, 2020, 1-6. | 2.5 | 9 |
| 13 | The rs1051931 G>A Polymorphism in the PLA2G7 Gene Confers Resistance to Immunoglobulin Therapy in Kawasaki Disease in a Southern Chinese Population. <i>Frontiers in Pediatrics</i> , 2020, 8, 338. | 1.9 | 3 |
| 14 | lncRNA HULC Polymorphism Is Associated With Recurrent Spontaneous Abortion Susceptibility in the Southern Chinese Population. <i>Frontiers in Genetics</i> , 2019, 10, 918. | 2.3 | 10 |
| 15 | Association between the TOX3 rs3803662 C>T polymorphism and recurrent miscarriage in a southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22992. | 2.1 | 2 |
| 16 | Association between the rs2288947 polymorphism of the lncRNA <i>TINCR</i> gene and the risk of recurrent miscarriage in a Southern Chinese population. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22919. | 2.1 | 7 |
| 17 | The IL-1B Gene Polymorphisms rs16944 and rs1143627 Contribute to an Increased Risk of Coronary Artery Lesions in Southern Chinese Children with Kawasaki Disease. <i>Journal of Immunology Research</i> , 2019, 2019, 1-7. | 2.2 | 24 |
| 18 | The <i>miRNA-608</i> rs4919510 G>C polymorphism confers reduce coronary injury of Kawasaki disease in a Southern Chinese population. <i>Bioscience Reports</i> , 2019, 39, . | 2.4 | 3 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | The lncRNA MALAT1 rs619586 G Variant Confers Decreased Susceptibility to Recurrent Miscarriage. <i>Frontiers in Physiology</i> , 2019, 10, 385. | 2.8 | 24 |
| 20 | The lncRNA <i>CCAT2</i> rs6983267 G allele is associated with decreased susceptibility to recurrent miscarriage. <i>Journal of Cellular Physiology</i> , 2019, 234, 20577-20583. | 4.1 | 16 |
| 21 | An Angiotensinogen Gene Polymorphism (rs5050) Is Associated with the Risk of Coronary Artery Aneurysm in Southern Chinese Children with Kawasaki Disease. <i>Disease Markers</i> , 2019, 2019, 1-7. | 1.3 | 9 |
| 22 | A <i>PEAR1</i> polymorphism (<i>rs12041331</i>) is associated with risk of coronary artery aneurysm in Kawasaki disease. <i>Annals of Human Genetics</i> , 2019, 83, 54-62. | 0.8 | 12 |
| 23 | P2RY12:rs7637803 TT variant genotype increases coronary artery aneurysm risk in Kawasaki disease in a southern Chinese population. <i>Journal of Gene Medicine</i> , 2019, 21, e3066. | 2.8 | 10 |
| 24 | Lack of association between <i>miR-218</i> rs11134527 A>G and Kawasaki disease susceptibility. <i>Bioscience Reports</i> , 2018, 38, . | 2.4 | 5 |
| 25 | ABCC4 Variants Modify Susceptibility to Kawasaki Disease in a Southern Chinese Population. <i>Disease Markers</i> , 2018, 2018, 1-7. | 1.3 | 9 |
| 26 | TBXA2R rs4523 G allele is associated with decreased susceptibility to Kawasaki disease. <i>Cytokine</i> , 2018, 111, 216-221. | 3.2 | 9 |
| 27 | The rs1625579 T>G polymorphism in the <i>miR-137</i> gene confers a risk of early-onset Kawasaki disease in a southern Chinese population. <i>Infection and Drug Resistance</i> , 2018, Volume 11, 1055-1060. | 2.7 | 11 |
| 28 | The association between the <i>miR-146a</i> rs2910164 C>G polymorphism and Kawasaki disease in a southern Chinese population. <i>Bioscience Reports</i> , 2018, 38, . | 2.4 | 9 |