
Kyung-Chae Park

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5006942/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	Prediction of prevalent but not incident nonâ€alcoholic fatty liver disease by levels of serum testosterone. Journal of Gastroenterology and Hepatology (Australia), 2015, 30, 1211-1216.	2.8	33
2	Coronary artery calcification is associated with high serum concentration of undercarboxylated osteocalcin in asymptomatic <scp>K</scp> orean men. Clinical Endocrinology, 2015, 83, 320-326.	2.4	25
3	Intermuscular adipose tissue is associated with monocyte chemoattractant protein-1, independent of visceral adipose tissue. Clinical Biochemistry, 2016, 49, 439-443.	1.9	11
4	Association of serum 25-hydroxyvitamin D and serum total cholesterol with depressive symptoms in Korean adults: the Fifth Korean National Health and Nutrition Examination Survey (KNHANES V,) Tj ETQq0 0 0 rgB	T ⊉Q verloc	k1110 Tf 50 6
5	Low hair copper concentration is related to a high risk of nonalcoholic fatty liver disease in adults. Journal of Trace Elements in Medicine and Biology, 2018, 50, 28-33.	3.0	11
6	Low Level of Osteocalcin Is Related With Arterial Stiffness in Korean Adults: An Inverse J-Shaped Relationship. Journal of Clinical Endocrinology and Metabolism, 2016, 101, 96-102.	3.6	8
7	A cross-sectional study of the association between adipokine levels and bone mineral density according to obesity and menopausal status in Korean women. Journal of Bone and Mineral Metabolism, 2017, 35, 642-648.	2.7	7
8	Sex Difference in the Association between Serum Homocysteine Level and Non-Alcoholic Fatty Liver Disease. Korean Journal of Family Medicine, 2016, 37, 242.	1.2	7

9	Association between Nonalcoholic Fatty Liver Disease and Concentration of Copper and Zinc in Hair Tissue in Korean Adults. Korean Journal of Family Practice, 2018, 8, 423-428.	c