

Giancarlo Palumbo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5012143/publications.pdf>

Version: 2024-02-01

32
papers

773
citations

623188

14
h-index

500791

28
g-index

32
all docs

32
docs citations

32
times ranked

733
citing authors

#	ARTICLE	IF	CITATIONS
1	Melphalan hypoxic perfusion with hemofiltration for melanoma locoregional metastases in the pelvis. <i>Journal of Surgical Research</i> , 2017, 215, 114-124.	0.8	7
2	MGMT methylation correlates with melphalan pelvic perfusion survival in stage III melanoma patients: a pilot study. <i>Melanoma Research</i> , 2017, 27, 439-447.	0.6	24
3	Does Locoregional Chemotherapy Still Matter in the Treatment of Advanced Pelvic Melanoma?. <i>International Journal of Molecular Sciences</i> , 2017, 18, 2382.	1.8	8
4	Surgical versus percutaneous isolated pelvic perfusion (IPP) for advanced melanoma: comparison in terms of melphalan pharmacokinetic pelvic bio-availability. <i>BMC Research Notes</i> , 2017, 10, 411.	0.6	10
5	Mesalazine vanishing time from rectal mucosa following its topical administration. <i>Journal of Crohn's and Colitis</i> , 2010, 4, 102-105.	0.6	21
6	Development of a method for the determination of vardenafil in human plasma by high performance liquid chromatography with UV detection. <i>Biomedical Chromatography</i> , 2009, 23, 759-763.	0.8	18
7	Thoracic stop-flow perfusion for refractory lymphoma: a phase I-II evaluation trial. <i>In Vivo</i> , 2009, 23, 447-57.	0.6	1
8	A validated HPLC method with electrochemical detection for simultaneous assay of 5-aminosalicylic acid and its metabolite in human plasma. <i>Biomedical Chromatography</i> , 2005, 19, 350-354.	0.8	47
9	Long-term oral plus topical mesalazine in frequently relapsing ulcerative colitis. <i>Digestive and Liver Disease</i> , 2005, 37, 92-96.	0.4	55
10	Development of a Liquid Chromatographic Method for the Determination of Sildenafil in Seminal Plasma. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2004, 27, 3039-3050.	0.5	2
11	Deliberate hypoxic pelvic and limb chemoperfusion in the treatment of recurrent melanoma. <i>American Journal of Surgery</i> , 2002, 183, 28-36.	0.9	25
12	N-Butyrate rectal transport in cirrhotic patients. <i>Digestive Diseases and Sciences</i> , 2001, 46, 2084-2088.	1.1	4
13	Neurochemical and Behavioural Effects of Chronic Aldicarb Administration in Rats. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2001, 89, 249-254.	0.0	4
14	Simultaneous determination of losartan and hydrochlorothiazide in tablets by high-performance liquid chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2000, 23, 185-189.	1.4	84
15	Mucosal 5-aminosalicylic acid concentration inversely correlates with severity of colonic inflammation in patients with ulcerative colitis. <i>Gut</i> , 2000, 47, 410-414.	6.1	121
16	Prevention of post-operative recurrence of Crohn's disease requires adequate mucosal concentration of mesalazine. <i>Alimentary Pharmacology and Therapeutics</i> , 1999, 13, 577-582.	1.9	35
17	Rectal and colonic mesalazine concentration in ulcerative colitis: oral vs. oral plus topical treatment. <i>Alimentary Pharmacology and Therapeutics</i> , 1999, 13, 1413-1417.	1.9	87
18	Mitomycin C in biological fluids: Comparison of disk and cartridge solid-phase extraction and quantitation by HPLC in human plasma and ultrafiltrate. <i>Chromatographia</i> , 1999, 50, 561-564.	0.7	4

#	ARTICLE	IF	CITATIONS
19	Simultaneous determination of rufloxacin, fenbufen and felbinac in human plasma using high-performance liquid chromatography. <i>Biomedical Applications</i> , 1996, 682, 315-319.	1.7	12
20	Simultaneous determination of 5-aminosalicylic acid, acetyl-5-aminosalicylic acid and 2,5-dihydroxybenzoic acid in endoscopic intestinal biopsy samples in humans by high-performance liquid chromatography with electrochemical detection. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1995, 14, 175-180.	1.4	43
21	Simultaneous determination of rufloxacin and theophylline by high-performance liquid chromatography in human plasma. <i>Analyst, The</i> , 1995, 120, 2493.	1.7	6
22	A high performance liquid chromatographic procedure for the simultaneous determination of norfloxacin and furprofen in rat plasma. <i>Biomedical Chromatography</i> , 1993, 7, 126-128.	0.8	13
23	Determination of Tenoxicam in Human Plasma using Solid-Phase Extraction and High-Performance Liquid Chromatography with Ultraviolet Detection. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1992, 15, 683-695.	0.9	18
24	Pharmacokinetics of intra-arterial nitonycin C with extracorporeal detoxification in humans. <i>Pharmacological Research</i> , 1992, 26, 98.	3.1	0
25	Indirect stereoselective determination of enantiomers of furprofen in human plasma by high-performance liquid chromatography. <i>Chromatographia</i> , 1992, 34, 618-620.	0.7	2
26	Determination of Rufloxacin, a New Tricyclic Fluoroquinolone in Biological Fluids Using High-Performance Liquid Chromatography with Ultraviolet Detection. <i>Therapeutic Drug Monitoring</i> , 1991, 13, 448-451.	1.0	5
27	Analytical procedure for the determination of rufloxacin, a new pyridobenzothiazine, in human serum and urine by high-performance liquid chromatography. <i>Biomedical Applications</i> , 1991, 564, 346-351.	1.7	13
28	A high-performance liquid chromatographic method for the evaluation of aldrin epoxidation by cytochrome P-450 dependent monooxygenase in small liver samples. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1989, 7, 783-788.	1.4	2
29	Chronic atrophic gastritis and risk of n-nitroso compounds carcinogenesis. <i>Langenbecks Archiv Fur Chirurgie</i> , 1988, 373, 82-90.	0.2	10
30	Determination of furprofen in human plasma by solid-phase extraction and reversed-phase high-performance liquid chromatography. <i>Biomedical Applications</i> , 1988, 424, 201-204.	1.7	3
31	Effects of Maternal Exposure to Polychlorobiphenyls (PCBs) on F1 Generation Behavior in the Rat. <i>Toxicological Sciences</i> , 1988, 11, 440-449.	1.4	70
32	Determination of Ofloxacin, A New Oxazine Derivative in Human Serum, Urine, and Bile by High-Performance Liquid Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1986, 9, 2539-2547.	0.9	19