

Kim HÃrslev-Petersen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5017359/publications.pdf>

Version: 2024-02-01

158
papers

5,431
citations

71102

41
h-index

98798

67
g-index

160
all docs

160
docs citations

160
times ranked

5515
citing authors

#	ARTICLE	IF	CITATIONS
1	Cardiovascular disease in patients with rheumatoid arthritis: results from the QUEST-RA study. <i>Arthritis Research and Therapy</i> , 2008, 10, R30.	3.5	387
2	MRI bone oedema is the strongest predictor of subsequent radiographic progression in early rheumatoid arthritis. Results from a 2-year randomised controlled trial (CIMESTRA). <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 384-390.	0.9	335
3	Radiographic progression and remission rates in early rheumatoid arthritis - MRI bone oedema and anti-CCP predicted radiographic progression in the 5-year extension of the double-blind randomised CIMESTRA trial. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1789-1795.	0.9	168
4	Combination treatment with methotrexate, cyclosporine, and intraarticular betamethasone compared with methotrexate and intraarticular betamethasone in early active rheumatoid arthritis: An investigator-initiated, multicenter, randomized, double-blind, parallel-group, placebo-controlled study. <i>Arthritis and Rheumatism</i> , 2006, 54, 1401-1409.	6.7	160
5	Serum YKL-40 concentrations in patients with rheumatoid arthritis: relation to disease activity. <i>British Journal of Rheumatology</i> , 1999, 38, 618-626.	2.3	156
6	Remission and rheumatoid arthritis: Data on patients receiving usual care in twenty-four countries. <i>Arthritis and Rheumatism</i> , 2008, 58, 2642-2651.	6.7	156
7	Adalimumab added to a treat-to-target strategy with methotrexate and intra-articular triamcinolone in early rheumatoid arthritis increased remission rates, function and quality of life. The OPERA Study: an investigator-initiated, randomised, double-blind, parallel-group, placebo-controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 654-661.	0.9	108
8	Plasma Adiponectin in Patients with Active, Early, and Chronic Rheumatoid Arthritis Who Are Steroid- and Disease-Modifying Antirheumatic Drug-Naive Compared with Patients with Osteoarthritis and Controls. <i>Journal of Rheumatology</i> , 2009, 36, 1885-1891.	2.0	106
9	Synovitis and Osteitis Are Very Frequent in Rheumatoid Arthritis Clinical Remission: Results from an MRI Study of 294 Patients in Clinical Remission or Low Disease Activity State. <i>Journal of Rheumatology</i> , 2011, 38, 2039-2044.	2.0	101
10	Increased Interleukin 21 (IL-21) and IL-23 Are Associated with Increased Disease Activity and with Radiographic Status in Patients with Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2010, 37, 2014-2020.	2.0	94
11	Aggressive combination therapy with intra-articular glucocorticoid injections and conventional disease-modifying anti-rheumatic drugs in early rheumatoid arthritis: second-year clinical and radiographic results from the CIMESTRA study. <i>Annals of the Rheumatic Diseases</i> , 2008, 67, 815-822.	0.9	93
12	Bone edema on magnetic resonance imaging is an independent predictor of rheumatoid arthritis development in patients with early undifferentiated arthritis. <i>Arthritis and Rheumatism</i> , 2011, 63, 2192-2202.	6.7	90
13	Low field dedicated magnetic resonance imaging in untreated rheumatoid arthritis of recent onset. <i>Annals of the Rheumatic Diseases</i> , 2001, 60, 770-776.	0.9	82
14	Pre-analytical and biological variability in circulating interleukin 6 in healthy subjects and patients with rheumatoid arthritis. <i>Biomarkers</i> , 2008, 13, 59-78.	1.9	80
15	Prevalence of degenerative and spondyloarthritis-related magnetic resonance imaging findings in the spine and sacroiliac joints in patients with persistent low back pain. <i>European Radiology</i> , 2016, 26, 1191-1203.	4.5	80
16	Increased soluble programmed death-1 (sPD-1) is associated with disease activity and radiographic progression in early rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2014, 43, 101-108.	1.1	79
17	Shared care or nursing consultations as an alternative to rheumatologist follow-up for rheumatoid arthritis outpatients with low disease activity—patient outcomes from a 2-year, randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 357-364.	0.9	78
18	Everyday Life with Rheumatoid Arthritis and Implications for Patient Education and Clinical Practice: A Focus Group Study. <i>Musculoskeletal Care</i> , 2012, 10, 29-38.	1.4	76

#	ARTICLE	IF	CITATIONS
19	Intra-articular hyaluronan is without clinical effect in knee osteoarthritis: a multicentre, randomised, placebo-controlled, double-blind study of 337 patients followed for 1 year. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1097.2-1102.	0.9	71
20	Determining a Magnetic Resonance Imaging Inflammatory Activity Acceptable State Without Subsequent Radiographic Progression in Rheumatoid Arthritis: Results from a Followup MRI Study of 254 Patients in Clinical Remission or Low Disease Activity. <i>Journal of Rheumatology</i> , 2014, 41, 398-406.	2.0	71
21	Associations Between Spondyloarthritis Features and Magnetic Resonance Imaging Findings: A Cross-sectional Analysis of 1,020 Patients With Persistent Low Back Pain. <i>Arthritis and Rheumatology</i> , 2016, 68, 892-900.	5.6	71
22	Short- and long-term efficacy of intra-articular injections with betamethasone as part of a treat-to-target strategy in early rheumatoid arthritis: impact of joint area, repeated injections, MRI findings, anti-CCP, IgM-RF and CRP. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 851-856.	0.9	70
23	Effect of Magnetic Resonance Imaging vs Conventional Treat-to-Target Strategies on Disease Activity Remission and Radiographic Progression in Rheumatoid Arthritis. <i>JAMA - Journal of the American Medical Association</i> , 2019, 321, 461.	7.4	70
24	Serum amino-terminal type III procollagen peptide in rheumatoid arthritis. Relationship to disease activity, treatment, and development of joint erosions. <i>Arthritis and Rheumatism</i> , 1986, 29, 592-599.	6.7	66
25	Procollagen type I N-terminal propeptide (PINP) as an indicator of type I collagen metabolism: ELISA development, reference interval, and hypovitaminosis D induced hyperparathyroidism. <i>Bone</i> , 1996, 19, 157-163.	2.9	63
26	Dynamic gadolinium-enhanced magnetic resonance imaging allows accurate assessment of the synovial inflammatory activity in rheumatoid arthritis knee joints: a comparison with synovial histology. <i>Scandinavian Journal of Rheumatology</i> , 2012, 41, 89-94.	1.1	61
27	CXCL13 predicts disease activity in early rheumatoid arthritis and could be an indicator of the therapeutic 'window of opportunity'. <i>Arthritis Research and Therapy</i> , 2014, 16, 434.	3.5	61
28	Incidence of rheumatoid arthritis from 1995 to 2001: impact of ascertainment from multiple sources. <i>Rheumatology International</i> , 2009, 29, 411-415.	3.0	59
29	Serum amino terminal type III procollagen peptide and serum hyaluronan in rheumatoid arthritis: relation to clinical and serological parameters of inflammation during 8 and 24 months' treatment with levamisole, penicillamine, or azathioprine.. <i>Annals of the Rheumatic Diseases</i> , 1988, 47, 116-126.	0.9	58
30	Low-cost, low-field dedicated extremity magnetic resonance imaging in early rheumatoid arthritis: a 1-year follow-up study. <i>Annals of the Rheumatic Diseases</i> , 2006, 65, 1208-1212.	0.9	58
31	Results from systematic screening for cardiovascular risk in outpatients with rheumatoid arthritis in accordance with the EULAR recommendations. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, 1771-1776.	0.9	58
32	Prediction of treatment response to adalimumab: a double-blind placebo-controlled study of circulating microRNA in patients with early rheumatoid arthritis. <i>Pharmacogenomics Journal</i> , 2016, 16, 141-146.	2.0	58
33	Serum aminoterminal type III procollagen peptide and the 7S domain of type IV collagen in patients with alcohol abuse Relation to ultrastructural fibrosis in the acinar zone 3 and to serum hyaluronan. <i>Liver</i> , 1987, 7, 339-346.	0.1	54
34	Magnetic resonance imaging and bone scintigraphy in the differential diagnosis of unclassified arthritis. <i>Annals of the Rheumatic Diseases</i> , 2008, 67, 48-51.	0.9	52
35	Active conventional treatment and three different biological treatments in early rheumatoid arthritis: phase IV investigator initiated, randomised, observer blinded clinical trial. <i>BMJ</i> , The, 2020, 371, m4328.	6.0	50
36	A treat-to-target strategy with methotrexate and intra-articular triamcinolone with or without adalimumab effectively reduces MRI synovitis, osteitis and tenosynovitis and halts structural damage progression in early rheumatoid arthritis: results from the OPERA randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 867-875.	0.9	49

#	ARTICLE	IF	CITATIONS
37	Limited Reliability of Radiographic Assessment of Sacroiliac Joints in Patients with Suspected Early Spondyloarthritis. <i>Journal of Rheumatology</i> , 2017, 44, 70-77.	2.0	48
38	Correlation between computer-aided dynamic gadolinium-enhanced MRI assessment of inflammation and semi-quantitative synovitis and bone marrow oedema scores of the wrist in patients with rheumatoid arthritis—a cohort study. <i>Rheumatology</i> , 2012, 51, 134-143.	1.9	47
39	Incidence of Rheumatoid Arthritis in the Southern part of Denmark from 1995 to 2001. <i>Open Rheumatology Journal</i> , 2007, 1, 18-23.	0.2	46
40	The impact on self-efficacy of different types of follow-up care and disease status in patients with rheumatoid arthritis—A randomized trial. <i>Patient Education and Counseling</i> , 2012, 88, 121-128.	2.2	43
41	Does Clinical Remission Lead to Normalization of EQ-5D in Patients with Rheumatoid Arthritis and Is Selection of Remission Criteria Important?. <i>Journal of Rheumatology</i> , 2010, 37, 285-290.	2.0	42
42	Bone Metabolism in Patients with Systemic Lupus Erythematosus: Effect of disease activity and glucocorticoid treatment. <i>Scandinavian Journal of Rheumatology</i> , 1998, 27, 197-206.	1.1	41
43	Clinical and radiographic outcome of a treat-to-target strategy using methotrexate and intra-articular glucocorticoids with or without adalimumab induction: a 2-year investigator-initiated, double-blinded, randomised, controlled trial (OPERA). <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 1645-1653.	0.9	40
44	What Factors Influence the Health Status of Patients with Rheumatoid Arthritis Measured by the SF-12v2 Health Survey and the Health Assessment Questionnaire?. <i>Journal of Rheumatology</i> , 2009, 36, 2183-2189.	2.0	39
45	Hepatic and renal extraction of circulating type III procollagen amino-terminal propeptide and hyaluronan in pig. <i>Journal of Hepatology</i> , 1989, 9, 177-183.	3.7	38
46	Serum Levels of Aminoterminal Type III Procollagen Peptide and Hyaluronan Predict Mortality in Systemic Sclerosis. <i>Scandinavian Journal of Rheumatology</i> , 1992, 21, 5-9.	1.1	37
47	Reliability and responsiveness of dynamic contrast-enhanced magnetic resonance imaging in rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2013, 42, 115-122.	1.1	37
48	Are human endogenous retroviruses triggers of autoimmune diseases? Unveiling associations of three diseases and viral loci. <i>Immunologic Research</i> , 2016, 64, 55-63.	2.9	37
49	Serum aminoterminal type III procollagen peptide. <i>Apmis</i> , 1988, 96, 793-804.	2.0	36
50	Being an outpatient with rheumatoid arthritis — a focus group study on patients' self-efficacy and experiences from participation in a short course and one of three different outpatient settings. <i>Scandinavian Journal of Caring Sciences</i> , 2011, 25, 394-403.	2.1	35
51	Plasma MicroRNA Profiles in Patients with Early Rheumatoid Arthritis Responding to Adalimumab plus Methotrexate vs Methotrexate Alone: A Placebo-controlled Clinical Trial. <i>Journal of Rheumatology</i> , 2018, 45, 53-61.	2.0	35
52	Shared care or nurse consultations as an alternative to rheumatologist follow-up for rheumatoid arthritis (RA) outpatients with stable low disease-activity RA: cost-effectiveness based on a 2-year randomized trial. <i>Scandinavian Journal of Rheumatology</i> , 2015, 44, 13-21.	1.1	34
53	Procollagen type III aminoterminal peptide in serum in idiopathic myelofibrosis and allied conditions: Relation to disease activity and effect of chemotherapy. <i>American Journal of Hematology</i> , 1990, 33, 18-26.	4.1	29
54	It Means Everything: Continuing Normality of Everyday Life for People with Rheumatoid Arthritis in Early Remission. <i>Musculoskeletal Care</i> , 2012, 10, 162-170.	1.4	28

#	ARTICLE	IF	CITATIONS
55	Soluble OX40L is associated with presence of autoantibodies in early rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2014, 16, 474.	3.5	28
56	Is the aminoterminal propeptide of type III procollagen degraded in the liver?. <i>Journal of Hepatology</i> , 1988, 6, 144-150.	3.7	27
57	Collagen Types I and III Propeptides as Markers of Healing in Chronic Leg Ulcers A Noninvasive Method for the Determination of Procollagen Propeptides in Wound Fluidâ€”Influence of Growth Hormone. <i>Annals of Surgery</i> , 1992, 216, 684-691.	4.2	26
58	Periarticular and generalised bone loss in patients with early rheumatoid arthritis: influence of alendronate and intra-articular glucocorticoid treatment. Post hoc analyses from the CIMESTRA trial. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 1123-1129.	0.9	26
59	The interleukin-20 receptor axis in early rheumatoid arthritis: novel links between disease-associated autoantibodies and radiographic progression. <i>Arthritis Research and Therapy</i> , 2016, 18, 61.	3.5	26
60	Magnetic resonance imaging assessed inflammation in the wrist is associated with patient-reported physical impairment, global assessment of disease activity and pain in early rheumatoid arthritis: longitudinal results from two randomised controlled trials. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 1707-1715.	0.9	26
61	Splanchnic and renal extraction of circulating type III procollagen aminoterminal propeptide in patients with normal liver function and in patients with alcoholic cirrhosis. <i>Hepatology</i> , 1990, 11, 957-963.	7.3	25
62	Serum and urinary aminoterminal type III procollagen peptide in progressive systemic sclerosis: relationship to scleroderma involvement, serum hyaluronan and urinary collagen metabolites. <i>Journal of Rheumatology</i> , 1988, 15, 460-7.	2.0	25
63	The aminoterminal-type-III procollagen peptide and proteoglycans in serum and synovial fluid of patients with rheumatoid arthritis or reactive arthritis. <i>Rheumatology International</i> , 1988, 8, 1-9.	3.0	24
64	Collagen type IV and procollagen type III during granulation tissue formation: a serological, biochemical, immunohistochemical and morphometrical study on the viscose cellulose sponge rat model. <i>European Journal of Clinical Investigation</i> , 1988, 18, 352-359.	3.4	23
65	Serum aminoterminal procollagen type III peptide in acute viral hepatitis. A longâ€”term followâ€”up study. <i>Liver</i> , 1987, 7, 96-105.	0.1	22
66	Selfâ€”Efficacy as an Outcome Measure and its Association with Physical Diseaseâ€”Related Variables in Persons with Rheumatoid Arthritis: A Literature Review. <i>Musculoskeletal Care</i> , 2011, 9, 125-140.	1.4	22
67	Brief Report: Mâ€”Ficolin Levels Reflect Disease Activity and Predict Remission in Early Rheumatoid Arthritis. <i>Arthritis and Rheumatism</i> , 2013, 65, 3045-3050.	6.7	22
68	Interactions between smoking, increased serum levels of anti-CCP antibodies, rheumatoid factors, and erosive joint disease in patients with early, untreated rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2015, 44, 8-12.	1.1	22
69	Expression of soluble CD83 in plasma from early-stage rheumatoid arthritis patients is not modified by anti-TNF-Î± therapy. <i>Cytokine</i> , 2017, 96, 1-7.	3.2	22
70	Goal-setting in multidisciplinary team care for patients with rheumatoid arthritis: An international multi-center evaluation of the contents using the International Classification of Functioning, Disability and Health as a reference. <i>Journal of Rehabilitation Medicine</i> , 2013, 45, 888-899.	1.1	21
71	Development of a framework identifying domains and elements of importance for arthritis rehabilitation. <i>Journal of Rehabilitation Medicine</i> , 2012, 44, 406-413.	1.1	20
72	Mannose-Binding Lectin Gene Polymorphisms Are Associated with Disease Activity and Physical Disability in Untreated, Anti-Cyclic Citrullinated Peptide-Positive Patients with Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2009, 36, 731-735.	2.0	19

#	ARTICLE	IF	CITATIONS
73	Self-efficacy in rheumatoid arthritis: translation and test of validity, reliability and sensitivity of the Danish version of the Rheumatoid Arthritis Self-Efficacy Questionnaire (RASE). <i>Musculoskeletal Care</i> , 2010, 8, 123-135.	1.4	19
74	Circulating surfactant protein -D is low and correlates negatively with systemic inflammation in early, untreated rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2010, 12, R39.	3.5	19
75	Upregulated baseline plasma CCL19 and CCR7 cell-surface expression on monocytes in early rheumatoid arthritis normalized during treatment and CCL19 correlated with radiographic progression. <i>Scandinavian Journal of Rheumatology</i> , 2014, 43, 91-100.	1.1	19
76	Diagnostic Delay in Rheumatoid Arthritis: A Qualitative Study of Symptom Interpretation Before the First Visit to the Doctor. <i>Musculoskeletal Care</i> , 2016, 14, 26-36.	1.4	19
77	Head-to-head comparison of aggressive conventional therapy and three biological treatments and comparison of two de-escalation strategies in patients who respond to treatment: study protocol for a multicenter, randomized, open-label, blinded-assessor, phase 4 study. <i>Trials</i> , 2017, 18, 161.	1.6	19
78	Anti-cyclic citrullinated peptide antibodies, 28-joint Disease Activity Score, and magnetic resonance imaging bone oedema at baseline predict 11 years' functional and radiographic outcome in early rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2019, 48, 1-8.	1.1	19
79	Differentiation between early rheumatoid arthritis patients and healthy persons by conventional and dynamic contrast-enhanced magnetic resonance imaging. <i>Scandinavian Journal of Rheumatology</i> , 2014, 43, 109-118.	1.1	18
80	Predictive value of a multi-biomarker disease activity score for clinical remission and radiographic progression in patients with early rheumatoid arthritis: a post-hoc study of the OPERA trial. <i>Scandinavian Journal of Rheumatology</i> , 2019, 48, 9-16.	1.1	18
81	Flares in rheumatoid arthritis: do patient-reported swollen and tender joints match clinical and ultrasonography findings?. <i>Rheumatology</i> , 2020, 59, 129-136.	1.9	18
82	A disintegrin and metalloprotease-17 and galectin-9 are important regulators of local 4-1BB activity and disease outcome in rheumatoid arthritis. <i>Rheumatology</i> , 2016, 55, 1871-1879.	1.9	17
83	The discriminative value of inflammatory back pain in patients with persistent low back pain. <i>Scandinavian Journal of Rheumatology</i> , 2016, 45, 321-328.	1.1	17
84	Different Effects of Thyroid Disease on Serum Levels of Procollagen III-N-Pep tide and Hyaluronic Acid. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1990, 71, 1016-1021.	3.6	16
85	Fibroblastic rheumatism: a Scandinavian case report. <i>Annals of the Rheumatic Diseases</i> , 2005, 64, 156-157.	0.9	16
86	Increased galectin-3 may serve as a serologic signature of pre-rheumatoid arthritis while markers of synovitis and cartilage do not differ between early undifferentiated arthritis subsets. <i>Arthritis Research and Therapy</i> , 2017, 19, 80.	3.5	16
87	Prevalence of Rheumatoid Arthritis in the Southern Part of Denmark. <i>Open Rheumatology Journal</i> , 2011, 5, 91-97.	0.2	16
88	Patterns of Magnetic Resonance Imaging Bone Erosion in Rheumatoid Arthritis – Which Bones Are Most Frequently Involved and Show the Most Change?: Figure 1.. <i>Journal of Rheumatology</i> , 2011, 38, 2014-2017.	2.0	15
89	Hand bone loss in early rheumatoid arthritis during a methotrexate-based treat-to-target strategy with or without adalimumab – a substudy of the optimized treatment algorithm in early RA (OPERA) trial. <i>Clinical Rheumatology</i> , 2017, 36, 781-789.	2.2	15
90	Juggling identities of rheumatoid arthritis, motherhood and paid work – a grounded theory study. <i>Disability and Rehabilitation</i> , 2019, 41, 1536-1544.	1.8	15

#	ARTICLE	IF	CITATIONS
91	Macrophage activity assessed by soluble CD163 in early rheumatoid arthritis: association with disease activity but different response patterns to synthetic and biologic DMARDs. <i>Clinical and Experimental Rheumatology</i> , 2015, 33, 498-502.	0.8	15
92	Cartilage Oligomeric Matrix Protein Associates Differentially with Erosions and Synovitis and Has a Different Temporal Course in Cyclic Citrullinated Peptide Antibody (Anti-CCP)-positive versus Anti-CCP-negative Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2011, 38, 1563-1568.	2.0	14
93	Within-day variation and influence of physical exercise on circulating Galectin-3 in patients with rheumatoid arthritis and healthy individuals. <i>Scandinavian Journal of Immunology</i> , 2015, 82, 70-75.	2.7	14
94	Cardiovascular risk profiles in a hospital-based population of patients with psoriatic arthritis and ankylosing spondylitis: a cross-sectional study. <i>Rheumatology International</i> , 2017, 37, 113-120.	3.0	14
95	The vitamin metabolite 1,25(OH) ₂ D in serum is associated with disease activity and Anti-Citrullinated Protein Antibodies in active and treatment naïve, early Rheumatoid Arthritis Patients. <i>Scandinavian Journal of Immunology</i> , 2018, 88, e12704.	2.7	14
96	Collagen metabolism during wound healing in rats. The aminoterminal propeptide of type III procollagen in serum and wound fluid in relation to formation of granulation tissue. <i>Apmis</i> , 1993, 101, 557-564.	2.0	13
97	Uncoupling of Collagen II Metabolism in Newly Diagnosed, Untreated Rheumatoid Arthritis Is Linked to Inflammation and Antibodies Against Cyclic Citrullinated Peptides. <i>Journal of Rheumatology</i> , 2010, 37, 1113-1120.	2.0	13
98	What's in the black box of arthritis rehabilitation? A comparison of rehabilitation practice for patients with inflammatory arthritis in Northern Europe. <i>Journal of Rehabilitation Medicine</i> , 2013, 45, 458-466.	1.1	13
99	Mortality and its predictors in patients with rheumatoid arthritis: a Danish population-based inception cohort study. <i>Scandinavian Journal of Rheumatology</i> , 2018, 47, 371-377.	1.1	13
100	CRP genotype and haplotype associations with serum C-reactive protein level and DAS28 in untreated early rheumatoid arthritis patients. <i>Arthritis Research and Therapy</i> , 2014, 16, 475.	3.5	12
101	Interleukin-23 in early disease development in rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2015, 44, 438-442.	1.1	12
102	Changes in Soluble CD18 in Murine Autoimmune Arthritis and Rheumatoid Arthritis Reflect Disease Establishment and Treatment Response. <i>PLoS ONE</i> , 2016, 11, e0148486.	2.5	12
103	A parametric study of effect of experimental tibialis posterior muscle pain on joint loading and muscle forces—Implications for patients with rheumatoid arthritis?. <i>Gait and Posture</i> , 2019, 72, 102-108.	1.4	12
104	Pain and Self-reported Swollen Joints Are Main Drivers of Patient-reported Flares in Rheumatoid Arthritis: Results from a 12-month Observational Study. <i>Journal of Rheumatology</i> , 2020, 47, 1305-1313.	2.0	12
105	Foot and Ankle Problems in Patients With Rheumatoid Arthritis in 2019: Still an Important Issue. <i>ACR Open Rheumatology</i> , 2021, 3, 396-402.	2.1	12
106	Serum aminoterminal type III procollagen peptide in inflammatory and degenerative rheumatic disorders. <i>Clinical Rheumatology</i> , 1988, 7, 61-68.	2.2	11
107	Impact of a magnetic resonance imaging-guided treat-to-target strategy on disease activity and progression in patients with rheumatoid arthritis (the IMAGINE-RA trial): study protocol for a randomized controlled trial. <i>Trials</i> , 2015, 16, 178.	1.6	11
108	Multifactorial intervention to prevent cardiovascular disease in patients with early rheumatoid arthritis: protocol for a multicentre randomised controlled trial. <i>BMJ Open</i> , 2016, 6, e009134.	1.9	11

#	ARTICLE	IF	CITATIONS
109	Soluble CD206 plasma levels in rheumatoid arthritis reflect decrease in disease activity. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2017, 77, 385-389.	1.2	11
110	Construction of meaningful identities in the context of rheumatoid arthritis, motherhood and paid work: A meta-ethnography. <i>Journal of Clinical Nursing</i> , 2017, 26, 4117-4128.	3.0	11
111	Serum aminoterminal type III procollagen peptide. Relation to biosynthesis of collagen type III in experimentally induced granulation tissue in rats. <i>Apmis</i> , 1988, 96, 793-804.	2.0	11
112	Circadian rhythm and the influence of physical activity on circulating surfactant protein D in early and long-standing rheumatoid arthritis. <i>Rheumatology International</i> , 2011, 31, 1617-1623.	3.0	10
113	Which patients improve the most after arthritis rehabilitation? A study of predictors in patients with inflammatory arthritis in Northern Europe, the STAR-ETIC collaboration. <i>Journal of Rehabilitation Medicine</i> , 2014, 46, 250-257.	1.1	10
114	Circulating serum interleukin-6, serum chitinase-3-like protein-1, and plasma vascular endothelial growth factor are not predictive for remission and radiographic progression in patients with early rheumatoid arthritis: post-hoc explorative and validation studies based on the CIMESTRA and OPERA trials. <i>Scandinavian Journal of Rheumatology</i> , 2018, 47, 259-269.	1.1	10
115	Impact of patient-reported flares on radiographic progression and functional impairment in patients with rheumatoid arthritis: a cohort study based on the AMBRA trial. <i>Scandinavian Journal of Rheumatology</i> , 2018, 47, 87-94.	1.1	10
116	Tibialis posterior muscle pain effects on hip, knee and ankle gait mechanics. <i>Human Movement Science</i> , 2019, 66, 98-108.	1.4	10
117	Programmed death ligand 2 – A link between inflammation and bone loss in rheumatoid arthritis. <i>Journal of Translational Autoimmunity</i> , 2020, 3, 100028.	4.0	10
118	Serum aminoterminal propeptide of type III procollagen after cardiac transplantation and the effect of rejection. <i>American Journal of Cardiology</i> , 1996, 78, 1406-1410.	1.6	9
119	Connective tissue metabolites in serum as markers of disease activity in patients with rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 1988, 6, 129-34.	0.8	9
120	Can magnetic resonance imaging differentiate undifferentiated arthritis?. <i>Arthritis Research and Therapy</i> , 2005, 7, 243.	3.5	8
121	Insulin-like growth factor I receptor density on CD4+T-lymphocytes from active early steroid- and DMARD-naïve rheumatoid arthritis patients is up-regulated and not influenced by 1 year of clinically effective treatment. <i>Rheumatology International</i> , 2012, 32, 501-504.	3.0	8
122	No diagnostic utility of antibody patterns against <i>Klebsiella pneumoniae</i> capsular serotypes in patients with axial spondyloarthritis vs. patients with non-specific low back pain: a cross-sectional study. <i>Scandinavian Journal of Rheumatology</i> , 2017, 46, 296-302.	1.1	8
123	Validity and Responsiveness of Combined Inflammation and Combined Joint Damage Scores Based on the OMERACT Rheumatoid Arthritis MRI Scoring System (RAMRIS). <i>Journal of Rheumatology</i> , 2019, 46, 1222-1227.	2.0	8
124	Serial magnetic resonance imaging and ultrasound examinations demonstrate differential inflammatory lesion patterns in soft tissue and bone upon patient-reported flares in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2020, 22, 19.	3.5	8
125	Impact of season on the association between vitamin D levels at diagnosis and one-year remission in early Rheumatoid Arthritis. <i>Scientific Reports</i> , 2020, 10, 7371.	3.3	8
126	Using an electronic platform interactively to improve treatment outcome in patients with rheumatoid arthritis: new developments from the DANBIO registry. <i>Clinical and Experimental Rheumatology</i> , 2016, 34, S75-S78.	0.8	8

#	ARTICLE	IF	CITATIONS
127	Effect of a treat-to-target strategy based on methotrexate and intra-articular betamethasone with or without additional cyclosporin on MRI-assessed synovitis, osteitis, tenosynovitis, bone erosion, and joint space narrowing in early rheumatoid arthritis: results from a 2-year randomized double-blind placebo-controlled trial (CIMESTRA). <i>Scandinavian Journal of Rheumatology</i> , 2017, 46, 335-345.	1.1	7
128	Different types of foot orthoses effect on gait mechanics in patients with rheumatoid arthritis. <i>Journal of Biomechanics</i> , 2021, , 110496.	2.1	7
129	Increased synovial galectin-3 induce inflammatory fibroblast activation and osteoclastogenesis in patients with rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2022, , 1-9.	1.1	7
130	Diurnal variation of connective tissue metabolites in early and long-standing rheumatoid arthritis and in healthy individuals. <i>Scandinavian Journal of Rheumatology</i> , 2011, 40, 88-94.	1.1	6
131	Hepcidin plasma levels are not associated with changes in haemoglobin in early rheumatoid arthritis patients. <i>Scandinavian Journal of Rheumatology</i> , 2017, 46, 441-445.	1.1	6
132	Predictors of joint damage progression and stringent remission in patients with established rheumatoid arthritis in clinical remission. <i>Rheumatology</i> , 2021, 60, 380-391.	1.9	6
133	Low-field magnetic resonance imaging or combined ultrasonography and anti-cyclic citrullinated peptide antibody improve correct classification of individuals as established rheumatoid arthritis: results of a population-based, cross-sectional study. <i>BMC Musculoskeletal Disorders</i> , 2014, 15, 268.	1.9	5
134	CXCL13 predicts long-term radiographic status in early rheumatoid arthritis. <i>Rheumatology</i> , 2022, 61, 2590-2595.	1.9	5
135	Circadian pattern and the effect of standardized physical exercise on procollagen IIA N-peptide (PIIANP) in rheumatoid arthritis at different stages and in healthy individuals. <i>Biomarkers</i> , 2010, 15, 80-85.	1.9	4
136	Association between baseline vitamin D metabolite levels and long-term cardiovascular events in patients with rheumatoid arthritis from the CIMESTRA trial: protocol for a cohort study with patient-record evaluated outcomes. <i>BMJ Open</i> , 2017, 7, e014816.	1.9	4
137	Similar lipid level changes in early rheumatoid arthritis patients following 1-year treat-to-target strategy with adalimumab plus methotrexate versus placebo plus methotrexate: secondary analyses from the randomised controlled OPERA trial. <i>Rheumatology International</i> , 2021, 41, 543-549.	3.0	4
138	Blood chemokine levels are markers of disease activity but not predictors of remission in early rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.8	4
139	Hepatic extraction of the aminoterminal propeptide of type III procollagen before and after bile duct ligation in pigs. <i>Liver</i> , 1991, 11, 310-315.	0.1	2
140	Does socioeconomic status make a difference? A register-based study on the extent to which cardiovascular screening in patients with inflammatory arthritis leads to recommended follow-up in general practice. <i>RMD Open</i> , 2020, 6, e000940.	3.8	2
141	The effect of foot orthoses on gait biomechanics and pain among people with rheumatoid arthritis: A quasi-experimental study. <i>Gait and Posture</i> , 2022, 95, 121-128.	1.4	2
142	Dose-Response Study of Human Growth Hormone in Venous Ulcers: Influence on Healing and Synthesis of Collagen Types I and III. <i>Phlebology</i> , 1994, 9, 92-98.	1.2	1
143	FR10017–Ox40 and ox40I are highly associated with autoantibody formation in early rheumatoid arthritis, and predict flare after anti-tnf discontinuation.. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A374.1-A374.	0.9	1
144	OP0223–M-ficolin, an activator of the complement system, predicts DAS28 remission in early DMARD na–ve rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013, 71, 131.1-131.	0.9	1

#	ARTICLE	IF	CITATIONS
145	Patient-specific foot orthotics improves postural control of rheumatoid arthritis patients: a pilot study. <i>Footwear Science</i> , 2017, 9, S17-S18.	2.1	1
146	Interactions between women with rheumatoid arthritis and nurses during outpatient consultations: A qualitative study. <i>Musculoskeletal Care</i> , 2019, 17, 363-371.	1.4	1
147	AB0316...Patient's global assessment of general health by vas at baseline predicts ACR/EULAR remission after 3, 6 and 12 months of efficient treatment in DMARD- and steroid naïve early rheumatoid arthritis patients. The danish cimestra-study. <i>Annals of the Rheumatic Diseases</i> , 2013, 71, 655.12-655.	0.9	0
148	SAT0391...Does dynamic contrast-enhanced mri provide better separation of active early rheumatoid arthritis patients and healthy controls than conventional MRI?.. <i>Annals of the Rheumatic Diseases</i> , 2013, 71, 604.2-605.	0.9	0
149	SAT0597...Course and Outcome of Rehabilitation Care in Different Rheumatological Diagnosis Groups. A Descriptive Study using the Star-Etic Registry.. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A784.3-A785.	0.9	0
150	THU0436...Screening for Cardiovascular Risk According to Euler 2010 Recommendations in Outpatients with Psoriasis Arthritis or Ankylosing Spondylitis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 333.2-333.	0.9	0
151	OP0073...Polymorphisms in the FCN1 Gene Coding for M-Ficolin are Associated with Disease Activity, Radiographic Damage and are the Strongest Predictors of DAS28 Remission in 180 DMARD Naive Early Rheumatoid Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 96.1-96.	0.9	0
152	FRI0535...Hand Bone Loss in Early Rheumatoid Arthritis Is Independent of Adalimumab Treatment. A Substudy of The Optimized Treatment Algorithm in Early RA (Opera) Trial. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 633.1-633.	0.9	0
153	AB0036...Soluble CD83 Plasma Levels Are Associated with Disease Activity and Course of Disease in Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 909.1-909.	0.9	0
154	THU0601...Cardiovascular screening among patients with inflammatory arthritis: to what extent do patients follow recommendations?., 2017, , .		0
155	...Isn't biological treatment something healthy? Lay people's perceptions of medical terms. <i>Scandinavian Journal of Rheumatology</i> , 2019, 48, 253-255.	1.1	0
156	Intra-articular and circulating levels of type I and III collagen markers in inflammatory and degenerative joint disease. <i>Acta Orthopaedica</i> , 1995, 66, 147-148.	1.4	0
157	AB0194...VITAMIN D TRAJECTORIES IN EARLY DIAGNOSED, AGGRESSIVELY TREATED RHEUMATOID ARTHRITIS PATIENTS: A 10 YEAR LONGITUDINAL COHORT STUDY BASED ON THE DANISH CIMESTRA TRIAL. <i>Annals of the Rheumatic Diseases</i> , 2020, 79, 1397-1397.	0.9	0
158	Blood chemokine levels are markers of disease activity but not predictors of remission in early rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2021, , .	0.8	0