

Ulf Bergström

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5018469/publications.pdf>

Version: 2024-02-01

61
papers

2,442
citations

172457

29
h-index

214800

47
g-index

62
all docs

62
docs citations

62
times ranked

2714
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Which factors can affect the productivity and dynamics of cod stocks in the Baltic Sea, Kattegat and Skagerrak?. <i>Ocean and Coastal Management</i> , 2022, 223, 106154. | 4.4 | 7 |
| 2 | Long-term decline in northern pike (<i>Esox lucius</i> L.) populations in the Baltic Sea revealed by recreational angling data. <i>Fisheries Research</i> , 2022, 251, 106307. | 1.7 | 22 |
| 3 | Ecological connectivity of the marine protected area network in the Baltic Sea, Kattegat and Skagerrak: Current knowledge and management needs. <i>Ambio</i> , 2022, 51, 1485-1503. | 5.5 | 14 |
| 4 | Increases of opportunistic species in response to ecosystem change: the case of the Baltic Sea three-spined stickleback. <i>ICES Journal of Marine Science</i> , 2022, 79, 1419-1434. | 2.5 | 16 |
| 5 | Habitat segregation of plate phenotypes in a rapidly expanding population of three-spined stickleback. <i>Ecosphere</i> , 2021, 12, e03561. | 2.2 | 7 |
| 6 | Environmental compensation for biodiversity and ecosystem services: A flexible framework that addresses human wellbeing. <i>Ecosystem Services</i> , 2021, 50, 101319. | 5.4 | 7 |
| 7 | Predator biomass and vegetation influence the coastal distribution of threespine stickleback morphotypes. <i>Ecology and Evolution</i> , 2021, 11, 12485-12496. | 1.9 | 3 |
| 8 | Predicting the effects of eutrophication mitigation on predatory fish biomass and the value of recreational fisheries. <i>Ambio</i> , 2020, 49, 1090-1099. | 5.5 | 4 |
| 9 | Feeding and growth of Atlantic cod (<i>Gadus morhua</i> L.) in the eastern Baltic Sea under environmental change. <i>ICES Journal of Marine Science</i> , 2020, 77, 624-632. | 2.5 | 55 |
| 10 | Perch and pike recruitment in coastal bays limited by stickleback predation and environmental forcing. <i>Estuarine, Coastal and Shelf Science</i> , 2020, 246, 107052. | 2.1 | 20 |
| 11 | A spatial regime shift from predator to prey dominance in a large coastal ecosystem. <i>Communications Biology</i> , 2020, 3, 459. | 4.4 | 56 |
| 12 | Long-term changes in spatial overlap between interacting cod and flounder in the Baltic Sea. <i>Hydrobiologia</i> , 2020, 847, 2541-2553. | 2.0 | 9 |
| 13 | Cumulative impact assessment for ecosystem-based marine spatial planning. <i>Science of the Total Environment</i> , 2020, 734, 139024. | 8.0 | 40 |
| 14 | A Model for Disentangling Dependencies and Impacts among Human Activities and Marine Ecosystem Services. <i>Environmental Management</i> , 2020, 65, 575-586. | 2.7 | 26 |
| 15 | Benefits and costs of two temporary no-take zones. <i>Marine Policy</i> , 2020, 117, 103883. | 3.2 | 3 |
| 16 | Spatial contraction of demersal fish populations in a large marine ecosystem. <i>Journal of Biogeography</i> , 2019, 46, 633-645. | 3.0 | 30 |
| 17 | The first large-scale assessment of three-spined stickleback (<i>Gasterosteus aculeatus</i>) biomass and spatial distribution in the Baltic Sea. <i>ICES Journal of Marine Science</i> , 2019, 76, 1653-1665. | 2.5 | 23 |
| 18 | Relative impacts of fishing and eutrophication on coastal fish assessed by comparing a no-take area with an environmental gradient. <i>Ambio</i> , 2019, 48, 565-579. | 5.5 | 24 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Recreational boating degrades vegetation important for fish recruitment. <i>Ambio</i> , 2019, 48, 539-551. | 5.5 | 33 |
| 20 | Evaluating complex relationships between ecological indicators and environmental factors in the Baltic Sea: A machine learning approach. <i>Ecological Indicators</i> , 2019, 101, 117-125. | 6.3 | 21 |
| 21 | Conflicts in the coastal zone: human impacts on commercially important fish species utilizing coastal habitat. <i>ICES Journal of Marine Science</i> , 2018, 75, 1203-1213. | 2.5 | 37 |
| 22 | Essential coastal habitats for fish in the Baltic Sea. <i>Estuarine, Coastal and Shelf Science</i> , 2018, 204, 14-30. | 2.1 | 48 |
| 23 | Competition for the fish " fish extraction from the Baltic Sea by humans, aquatic mammals, and birds. <i>ICES Journal of Marine Science</i> , 2018, 75, 999-1008. | 2.5 | 94 |
| 24 | Response to comments by Heikinheimo et al. (in press) on Hansson et al. (2018): competition for the fish" fish extraction from the Baltic Sea by humans, aquatic mammals, and birds. <i>ICES Journal of Marine Science</i> , 2018, 75, 1837-1839. | 2.5 | 3 |
| 25 | External nutrient loading from land, sea and atmosphere to all 656 Swedish coastal water bodies. <i>Marine Pollution Bulletin</i> , 2017, 114, 664-670. | 5.0 | 16 |
| 26 | Spatio-temporal dynamics of cod nursery areas in the Baltic Sea. <i>Progress in Oceanography</i> , 2017, 155, 28-40. | 3.2 | 14 |
| 27 | Modelling indices of abundance and size-based indicators of cod and flounder stocks in the Baltic Sea using newly standardized trawl survey data. <i>ICES Journal of Marine Science</i> , 2017, 74, 1322-1333. | 2.5 | 26 |
| 28 | Characterizing and predicting the distribution of Baltic Sea flounder (<i>Platichthys flesus</i>) during the spawning season. <i>Journal of Sea Research</i> , 2017, 126, 46-55. | 1.6 | 11 |
| 29 | DNA metabarcoding reveals diverse diet of the three-spined stickleback in a coastal ecosystem. <i>PLoS ONE</i> , 2017, 12, e0186929. | 2.5 | 59 |
| 30 | Size matters: relationships between body size and body mass of common coastal, aquatic invertebrates in the Baltic Sea. <i>PeerJ</i> , 2017, 5, e2906. | 2.0 | 35 |
| 31 | Spatio-temporal dynamics of a fish predator: Density-dependent and hydrographic effects on Baltic Sea cod population. <i>PLoS ONE</i> , 2017, 12, e0172004. | 2.5 | 22 |
| 32 | Distribution of mesopredatory fish determined by habitat variables in a predator-depleted coastal system. <i>Marine Biology</i> , 2016, 163, 201. | 1.5 | 18 |
| 33 | Coastal fish indicators response to natural and anthropogenic drivers" variability at temporal and different spatial scales. <i>Estuarine, Coastal and Shelf Science</i> , 2016, 183, 62-72. | 2.1 | 23 |
| 34 | Spawning areas of eastern Baltic cod revisited: Using hydrodynamic modelling to reveal spawning habitat suitability, egg survival probability, and connectivity patterns. <i>Progress in Oceanography</i> , 2016, 143, 13-25. | 3.2 | 23 |
| 35 | Top"down control as important as nutrient enrichment for eutrophication effects in North Atlantic coastal ecosystems. <i>Journal of Applied Ecology</i> , 2016, 53, 1138-1147. | 4.0 | 107 |
| 36 | Cod pots in a Baltic fishery: are they efficient and what affects their efficiency?. <i>ICES Journal of Marine Science</i> , 2015, 72, 1545-1554. | 2.5 | 32 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Declining coastal piscivore populations in the Baltic Sea: Where and when do sticklebacks matter?. <i>Ambio</i> , 2015, 44, 462-471. | 5.5 | 51 |
| 38 | Stickleback increase in the Baltic Sea – A thorny issue for coastal predatory fish. <i>Estuarine, Coastal and Shelf Science</i> , 2015, 163, 134-142. | 2.1 | 78 |
| 39 | Nursery habitat availability limits adult stock sizes of predatory coastal fish. <i>ICES Journal of Marine Science</i> , 2014, 71, 672-680. | 2.5 | 87 |
| 40 | Ecological value of coastal habitats for commercially and ecologically important species. <i>ICES Journal of Marine Science</i> , 2014, 71, 648-665. | 2.5 | 233 |
| 41 | Testing the Potential for Predictive Modeling and Mapping and Extending Its Use as a Tool for Evaluating Management Scenarios and Economic Valuation in the Baltic Sea (PREHAB). <i>Ambio</i> , 2014, 43, 82-93. | 5.5 | 11 |
| 42 | Pikeperch (<i>Sander lucioperca</i> (L.)) in Decline: High Mortality of Three Populations in the Northern Baltic Sea. <i>Ambio</i> , 2014, 43, 325-336. | 5.5 | 37 |
| 43 | Shoreline development and degradation of coastal fish reproduction habitats. <i>Ambio</i> , 2014, 43, 1020-1028. | 5.5 | 65 |
| 44 | Egg production of turbot, <i>Scophthalmus maximus</i> , in the Baltic Sea. <i>Journal of Sea Research</i> , 2013, 84, 77-86. | 1.6 | 9 |
| 45 | Evaluating eutrophication management scenarios in the Baltic Sea using species distribution modelling. <i>Journal of Applied Ecology</i> , 2013, 50, 680-690. | 4.0 | 43 |
| 46 | The influence of environmental conditions on early life stages of flounder (<i>Platichthys flesus</i>) in the central Baltic Sea. <i>Journal of Sea Research</i> , 2013, 75, 77-84. | 1.6 | 30 |
| 47 | Empirical modelling of benthic species distribution, abundance, and diversity in the Baltic Sea: evaluating the scope for predictive mapping using different modelling approaches. <i>ICES Journal of Marine Science</i> , 2013, 70, 1233-1243. | 2.5 | 45 |
| 48 | Estimating Competition between Wildlife and Humans – A Case of Cormorants and Coastal Fisheries in the Baltic Sea. <i>PLoS ONE</i> , 2013, 8, e83763. | 2.5 | 46 |
| 49 | Effects of an offshore wind farm on temporal and spatial patterns in the demersal fish community. <i>Marine Ecology - Progress Series</i> , 2013, 485, 199-210. | 1.9 | 94 |
| 50 | Ecological coherence of marine protected area networks: a spatial assessment using species distribution models. <i>Journal of Applied Ecology</i> , 2011, 48, 112-120. | 4.0 | 72 |
| 51 | A meso-predator release of stickleback promotes recruitment of macroalgae in the Baltic Sea. <i>Journal of Experimental Marine Biology and Ecology</i> , 2011, 397, 79-84. | 1.5 | 50 |
| 52 | Effects of Altered Offshore Food Webs on Coastal Ecosystems Emphasize the Need for Cross-Ecosystem Management. <i>Ambio</i> , 2011, 40, 786-797. | 5.5 | 100 |
| 53 | Recruitment failure of coastal predatory fish in the Baltic Sea coincident with an offshore ecosystem regime shift. <i>ICES Journal of Marine Science</i> , 2010, 67, 1587-1595. | 2.5 | 125 |
| 54 | Characterisation of juvenile flatfish habitats in the Baltic Sea. <i>Estuarine, Coastal and Shelf Science</i> , 2009, 82, 294-300. | 2.1 | 42 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Transferability of predictive fish distribution models in two coastal systems. <i>Estuarine, Coastal and Shelf Science</i> , 2009, 83, 90-96. | 2.1 | 55 |
| 56 | Spatial predictions of Baltic phytobenthic communities: Measuring robustness of generalized additive models based on transect data. <i>Journal of Marine Systems</i> , 2008, 74, S86-S96. | 2.1 | 22 |
| 57 | Plugging Space into Predator-Prey Models: An Empirical Approach. <i>American Naturalist</i> , 2006, 167, 246-259. | 2.1 | 35 |
| 58 | Spatial scale, heterogeneity and functional responses. <i>Journal of Animal Ecology</i> , 2004, 73, 487-493. | 2.8 | 29 |
| 59 | Small-scale spatial structure of Baltic Sea zoobenthos – inferring processes from patterns. <i>Journal of Experimental Marine Biology and Ecology</i> , 2002, 281, 123-136. | 1.5 | 39 |
| 60 | Estimating predation rates in experimental systems: scale-dependent effects of aggregative behaviour. <i>Oikos</i> , 2002, 97, 251-259. | 2.7 | 30 |
| 61 | Species diversity and distribution of aquatic macrophytes in the Northern Quark, Baltic Sea. <i>Nordic Journal of Botany</i> , 1999, 19, 375-383. | 0.5 | 26 |