

# Masanori Tamaoki

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/5036997/masanori-tamaoki-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

83

papers

3,424

citations

33

h-index

58

g-index

84

ext. papers

3,797

ext. citations

4.3

avg, IF

4.59

L-index

#	Paper	IF	Citations
83	Radiocesium Contamination in Wild Rodents Inhabiting Forested Areas Inside the Evacuated Area in Fukushima, Japan <b>2022</b> , 463-472		0
82	Monitoring of radioactive cesium in wild boars captured inside the difficult-to-return zone in Fukushima Prefecture over a 5-year period.. <i>Scientific Reports</i> , <b>2022</b> , 12, 5667	4.9	
81	Temporal Changes in the Parasite Fauna of the Large Japanese Field Mouse <i>Apodemus speciosus</i> in the Radioactive Contaminated Zone of Fukushima. <i>Japanese Journal of Zoo and Wildlife Medicine</i> , <b>2021</b> , 26, 1-5	0.1	
80	Untangling radiocesium dynamics of forest-stream ecosystems: A review of Fukushima studies in the decade after the accident. <i>Environmental Pollution</i> , <b>2021</b> , 288, 117744	9.3	1
79	Photosynthesis and growth of <i>Ulva ohnoi</i> and <i>Ulva pertusa</i> (Ulvophyceae) under high light and high temperature conditions, and implications for green tide in Japan. <i>Phycological Research</i> , <b>2020</b> , 68, 152-160	1.3	5
78	Occurrence of spilled genetically modified oilseed rape growing along a Japanese roadside over 10 years. <i>Weed Biology and Management</i> , <b>2020</b> , 20, 139-146	1.4	1
77	Estimation of Dose Rate for the Large Japanese Field Mouse ( <i>Apodemus speciosus</i> ) Distributed in the Difficult-to-Return Zone in Fukushima Prefecture <b>2020</b> , 17-30		3
76	Genetic Diversity of Invasive Loisel. (Poaceae) Introduced Unintentionally Into Japan and Its Invasion Pathway. <i>Frontiers in Plant Science</i> , <b>2020</b> , 11, 556039	6.2	9
75	Concentration of radioactive materials in small mammals collected from a restricted area in Fukushima, Japan since 2012. <i>Ecological Research</i> , <b>2019</b> , 34, 7-7	1.9	4
74	Comparison of 0.1 M Stable CsCl and 1 M NHNO as an Extraction Reagent to Evaluate Cs-137 Mobility in Soils. <i>Analytical Sciences</i> , <b>2019</b> , 35, 153-158	1.7	0
73	Impact of sea spider parasitism on host clams: susceptibility and intensity-dependent mortality. <i>Journal of the Marine Biological Association of the United Kingdom</i> , <b>2018</b> , 98, 735-742	1.1	3
72	Ozone-Sensitive Arabidopsis Mutants with Deficiencies in Photorespiratory Enzymes. <i>Plant and Cell Physiology</i> , <b>2017</b> , 58, 914-924	4.9	7
71	Molecular Mechanisms of Selenium Responses and Resistance in Plants. <i>Plant Ecophysiology</i> , <b>2017</b> , 35-51		11
70	Elevated Ozone Deteriorates Grain Quality of Japonica Rice cv. Koshihikari, Even if it Does Not Cause Yield Reduction. <i>Rice</i> , <b>2016</b> , 9, 7	5.8	11
69	Fixed-route monitoring and a comparative study of the occurrence of herbicide-resistant oilseed rape ( <i>Brassica napus</i> L.) along a Japanese roadside. <i>GM Crops and Food</i> , <b>2016</b> , 7, 20-37	2.7	13
68	Impact of sea spider parasitism on host clams: relationships between burial patterns and parasite loads, somatic condition and survival of host. <i>Hydrobiologia</i> , <b>2016</b> , 770, 15-26	2.4	7
67	Comparison of Potentials of Higher Plants for Phytoremediation of Radioactive Cesium from Contaminated Soil. <i>Environmental Control in Biology</i> , <b>2016</b> , 54, 65-69	0.9	5

66	Effects of environmental radiation on testes and spermatogenesis in wild large Japanese field mice ( <i>Apodemus speciosus</i> ) from Fukushima. <i>Scientific Reports</i> , <b>2016</b> , 6, 23601	4.9	35
65	Alteration of <i>Arabidopsis</i> SLAC1 promoter and its association with natural variation in drought tolerance. <i>Plant Signaling and Behavior</i> , <b>2015</b> , 10, e989761	2.5	5
64	$^{137}\text{Cs}$ concentrations in foliose lichens within Tsukuba-city as a reflection of radioactive fallout from the Fukushima Dai-ichi Nuclear Power Plant accident. <i>Journal of Environmental Radioactivity</i> , <b>2015</b> , 141, 38-43	2.4	9
63	Classification of the spermatogenic cycle, seasonal changes of seminiferous tubule morphology and estimation of the breeding season of the large Japanese field mouse ( <i>Apodemus speciosus</i> ) in Toyama and Aomori prefectures, Japan. <i>Journal of Veterinary Medical Science</i> , <b>2015</b> , 77, 799-807	1.1	6
62	18S analysis of the taxonomic position of an endoparasitic pycnogonid, <i>Nymphonella tapetis</i> (Arthropoda: Pycnogonida:Ascorhynchidae). <i>Journal of Crustacean Biology</i> , <b>2015</b> , 35, 491-494	0.8	4
61	Ozone-Induced Rice Grain Yield Loss Is Triggered via a Change in Panicle Morphology That Is Controlled by ABERRANT PANICLE ORGANIZATION 1 Gene. <i>PLoS ONE</i> , <b>2015</b> , 10, e0123308	3.7	36
60	Isolation and characterization of 25 polymorphic microsatellites of the large Japanese wood mouse ( <i>Apodemus speciosus</i> ). <i>Conservation Genetics Resources</i> , <b>2013</b> , 5, 1001-1003	0.8	
59	Quantitative trait locus analyses of ozone-induced grain yield reduction in rice. <i>Environmental and Experimental Botany</i> , <b>2013</b> , 88, 100-106	5.9	14
58	Characterization of hybrids between wild and genetically modified glyphosate-tolerant soybeans. <i>Plant Biotechnology</i> , <b>2013</b> , 30, 335-345	1.3	5
57	High-throughput capture of nucleotide sequence polymorphisms in three Brassica species (Brassicaceae) using DNA microarrays. <i>American Journal of Botany</i> , <b>2012</b> , 99, e94-6	2.7	1
56	The integral membrane protein SEN1 is required for symbiotic nitrogen fixation in <i>Lotus japonicus</i> nodules. <i>Plant and Cell Physiology</i> , <b>2012</b> , 53, 225-36	4.9	76
55	Expression and functions of myo-inositol monophosphatase family genes in seed development of <i>Arabidopsis</i> . <i>Journal of Plant Research</i> , <b>2011</b> , 124, 385-94	2.6	13
54	Seeds of a possible natural hybrid between herbicide-resistant <i>Brassica napus</i> and <i>Brassica rapa</i> detected on a riverbank in Japan. <i>GM Crops</i> , <b>2011</b> , 2, 201-10		38
53	Molecular mechanisms of selenium tolerance and hyperaccumulation in <i>Stanleya pinnata</i> . <i>Plant Physiology</i> , <b>2010</b> , 153, 1630-52	6.6	173
52	Rapeseed species and environmental concerns related to loss of seeds of genetically modified oilseed rape in Japan. <i>GM Crops</i> , <b>2010</b> , 1, 143-56		21
51	DEAR1, a transcriptional repressor of DREB protein that mediates plant defense and freezing stress responses in <i>Arabidopsis</i> . <i>Journal of Plant Research</i> , <b>2009</b> , 122, 633-43	2.6	115
50	Ethylene and salicylic acid control glutathione biosynthesis in ozone-exposed <i>Arabidopsis thaliana</i> . <i>Physiologia Plantarum</i> , <b>2009</b> , 136, 284-98	4.6	74
49	Monitoring the occurrence of genetically modified oilseed rape growing along a Japanese roadside: 3-year observations. <i>Environmental Biosafety Research</i> , <b>2009</b> , 8, 33-44		37

48	The Arabidopsis sweetie mutant is affected in carbohydrate metabolism and defective in the control of growth, development and senescence. <i>Plant Journal</i> , <b>2008</b> , 55, 665-86	6.9	53
47	Arabidopsis CAD1 negatively controls plant immunity mediated by both salicylic acid-dependent and -independent signaling pathways. <i>Plant Science</i> , <b>2008</b> , 175, 604-611	5.3	8
46	The role of phytohormone signaling in ozone-induced cell death in plants. <i>Plant Signaling and Behavior</i> , <b>2008</b> , 3, 166-74	2.5	58
45	New insights into the roles of ethylene and jasmonic acid in the acquisition of selenium resistance in plants. <i>Plant Signaling and Behavior</i> , <b>2008</b> , 3, 865-7	2.5	28
44	Cooperative ethylene and jasmonic acid signaling regulates selenite resistance in Arabidopsis. <i>Plant Physiology</i> , <b>2008</b> , 146, 1219-30	6.6	133
43	Disruption of a gene encoding C4-dicarboxylate transporter-like protein increases ozone sensitivity through deregulation of the stomatal response in Arabidopsis thaliana. <i>Plant and Cell Physiology</i> , <b>2008</b> , 49, 2-10	4.9	40
42	Transcriptome analyses give insights into selenium-stress responses and selenium tolerance mechanisms in Arabidopsis. <i>Physiologia Plantarum</i> , <b>2008</b> , 132, 236-53	4.6	119
41	An unidentified ultraviolet-B-specific photoreceptor mediates transcriptional activation of the cyclobutane pyrimidine dimer photolyase gene in plants. <i>Planta</i> , <b>2008</b> , 229, 25-36	4.7	14
40	SAZ, a new SUPERMAN-like protein, negatively regulates a subset of ABA-responsive genes in Arabidopsis. <i>Molecular Genetics and Genomics</i> , <b>2008</b> , 279, 183-92	3.1	22
39	Isolation of O <sub>3</sub> -response genes from Arabidopsis thaliana using cDNA macroarray. <i>Methods in Molecular Biology</i> , <b>2008</b> , 410, 29-42	1.4	
38	The isochlorismate pathway is negatively regulated by salicylic acid signaling in O <sub>3</sub> -exposed Arabidopsis. <i>Planta</i> , <b>2007</b> , 226, 1277-85	4.7	33
37	Complete genomic structure of the bloom-forming toxic cyanobacterium <i>Microcystis aeruginosa</i> NIES-843. <i>DNA Research</i> , <b>2007</b> , 14, 247-56	4.5	202
36	Glycosylation of bisphenol A by freshwater microalgae. <i>Chemosphere</i> , <b>2007</b> , 69, 934-41	8.4	78
35	Cytokinin and auxin inhibit abscisic acid-induced stomatal closure by enhancing ethylene production in Arabidopsis. <i>Journal of Experimental Botany</i> , <b>2006</b> , 57, 2259-66	7	160
34	Cytosolic dehydroascorbate reductase is important for ozone tolerance in Arabidopsis thaliana. <i>Plant and Cell Physiology</i> , <b>2006</b> , 47, 304-8	4.9	101
33	O <sub>2</sub> - activates leaf injury, ethylene and salicylic acid synthesis, and the expression of O <sub>3</sub> -induced genes in O <sub>3</sub> -exposed tobacco. <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , <b>2006</b> , 61, 856-64	1.7	2
32	Development of visible markers for transgenic plants and their availability for environmental risk assessment. <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , <b>2006</b> , 61, 377-86	1.7	4
31	Detection of feral transgenic oilseed rape with multiple-herbicide resistance in Japan. <i>Environmental Biosafety Research</i> , <b>2006</b> , 5, 77-87		54

30	Salicylic acid accumulation under O <sub>3</sub> exposure is regulated by ethylene in tobacco plants. <i>Plant and Cell Physiology</i> , <b>2005</b> , 46, 1062-72	4.9	79
29	The Arabidopsis gene CAD1 controls programmed cell death in the plant immune system and encodes a protein containing a MACPF domain. <i>Plant and Cell Physiology</i> , <b>2005</b> , 46, 902-12	4.9	78
28	Monitoring the escape of transgenic oilseed rape around Japanese ports and roadsides. <i>Environmental Biosafety Research</i> , <b>2005</b> , 4, 217-22		73
27	Ethylene inhibits abscisic acid-induced stomatal closure in Arabidopsis. <i>Plant Physiology</i> , <b>2005</b> , 138, 2337-43	4.3	295
26	Spermidine synthase genes are essential for survival of Arabidopsis. <i>Plant Physiology</i> , <b>2004</b> , 135, 1565-73	3.6	181
25	A method for diagnosis of plant environmental stresses by gene expression profiling using a cDNA microarray. <i>Environmental Pollution</i> , <b>2004</b> , 131, 137-45	9.3	30
24	Compensation for lack of a cytosolic ascorbate peroxidase in an Arabidopsis mutant by activation of multiple antioxidative systems. <i>Plant Science</i> , <b>2004</b> , 166, 1547-1554	5.3	12
23	Genomic structure of the cucumber CPD photolyase gene. <i>OMICS A Journal of Integrative Biology</i> , <b>2003</b> , 7, 203-9	3.8	1
22	Novel Marker Gene for Assessment of Behavior of Transgenic Plants in the Field. <i>Plant Biotechnology</i> , <b>2003</b> , 20, 225-227	1.3	
21	Transcriptome analysis of O <sub>3</sub> -exposed Arabidopsis reveals that multiple signal pathways act mutually antagonistically to induce gene expression. <i>Plant Molecular Biology</i> , <b>2003</b> , 53, 443-56	4.6	85
20	Differential ozone sensitivity among Arabidopsis accessions and its relevance to ethylene synthesis. <i>Planta</i> , <b>2003</b> , 216, 552-60	4.7	66
19	Light-controlled expression of a gene encoding l-galactono- $\gamma$ -lactone dehydrogenase which affects ascorbate pool size in Arabidopsis thaliana. <i>Plant Science</i> , <b>2003</b> , 164, 1111-1117	5.3	100
18	Isolation of an ozone-sensitive and jasmonate-semi-insensitive Arabidopsis mutant (oji1). <i>Plant and Cell Physiology</i> , <b>2003</b> , 44, 1301-10	4.9	53
17	Improvement in ozone tolerance of tobacco plants with an antisense DNA for 1-aminocyclopropane-1-carboxylate synthase. <i>Plant, Cell and Environment</i> , <b>2002</b> , 25, 727-735	8.4	32
16	Processing of bisphenol A by plant tissues: glucosylation by cultured BY-2 cells and glucosylation/translocation by plants of Nicotiana tabacum. <i>Plant and Cell Physiology</i> , <b>2002</b> , 43, 1036-42	4.9	55
15	Expression of genes encoding late nodulins characterized by a putative signal peptide and conserved cysteine residues is reduced in ineffective pea nodules. <i>Molecular Plant-Microbe Interactions</i> , <b>2002</b> , 15, 129-37	3.6	26
14	CDNA microarray assessment for ozone-stressed Arabidopsis thaliana. <i>Environmental Pollution</i> , <b>2002</b> , 117, 191-4	9.3	29
13	Effects of ozone exposure on the gene expression of ethylene biosynthetic enzymes in tomato leaves. <i>Plant Physiology and Biochemistry</i> , <b>2001</b> , 39, 993-998	5.4	18

12	Two types of pea leghemoglobin genes showing different O <sub>2</sub> -binding affinities and distinct patterns of spatial expression in nodules. <i>Plant Physiology</i> , <b>2001</b> , 125, 641-51	6.6	25
11	The Conserved KNOX Domain Mediates Specificity of Tobacco KNOTTED1-Type Homeodomain Proteins. <i>Plant Cell</i> , <b>1999</b> , 11, 1419	11.6	
10	The conserved KNOX domain mediates specificity of tobacco KNOTTED1-type homeodomain proteins. <i>Plant Cell</i> , <b>1999</b> , 11, 1419-32	11.6	42
9	The expression of tobacco knotted1-type class 1 homeobox genes correspond to regions predicted by the cytohistological zonation model. <i>Plant Journal</i> , <b>1999</b> , 18, 337-47	6.9	65
8	The homeobox gene NTH23 of tobacco is expressed in the basal region of leaf primordia. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , <b>1998</b> , 1399, 203-8		10
7	Alteration of hormone levels in transgenic tobacco plants overexpressing the rice homeobox gene OSH1. <i>Plant Physiology</i> , <b>1998</b> , 116, 471-6	6.6	112
6	Dorsoventral pattern formation of tobacco leaf involves spatial expression of a tobacco homeobox gene, NTH15.. <i>Genes and Genetic Systems</i> , <b>1997</b> , 72, 1-8	1.4	4
5	Two transcripts with different sizes derived from a rice homeobox gene, OSH1. <i>Biochemical and Biophysical Research Communications</i> , <b>1996</b> , 221, 408-13	3.4	3
4	Abnormal cell divisions in leaf primordia caused by the expression of the rice homeobox gene. <i>Molecular Genetics and Genomics</i> , <b>1996</b> , 251, 13		3
3	Alternative RNA products from a rice homeobox gene. <i>Plant Journal</i> , <b>1995</b> , 7, 927-38	6.9	66
2	Expression of nodulin genes in plant-determined ineffective nodules of pea. <i>Plant Molecular Biology</i> , <b>1995</b> , 28, 1027-38	4.6	17
1	Expression of rice OSH1 gene is localized in developing vascular strands and its ectopic expression in transgenic rice causes altered morphology of leaf. <i>Plant Cell Reports</i> , <b>1995</b> , 14, 555-9	5.1	13