

Hansruedi Glatt

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/503866/publications.pdf>

Version: 2024-02-01

197
papers

8,911
citations

38660

50
h-index

54797

84
g-index

203
all docs

203
docs citations

203
times ranked

6811
citing authors

#	ARTICLE	IF	CITATIONS
1	Feeding Brassica vegetables to rats leads to the formation of characteristic DNA adducts (from) Tj ETQq1 1 0.784314 rgBT /Qverlock 10	1.9	1
2	The comparison of cytotoxic and genotoxic activities of glucosinolates, isothiocyanates, and indoles. <i>Scientific Reports</i> , 2022, 12, 4875.	1.6	3
3	Potent aneugenicity of 1-methylpyrene in human cells dependent on metabolic activation by endogenous enzymes. <i>Archives of Toxicology</i> , 2021, 95, 703-713.	1.9	5
4	Influence of Bisphenol Compounds at Nanomolar Concentrations on Chromosome Damage Induced by Metabolically Activated Carcinogens in HepG2 Cells. <i>Environmental Science & Technology</i> , 2021, 55, 10001-10011.	4.6	20
5	Human CYP1B1-dependent genotoxicity of dioxin-like polychlorinated biphenyls in mammalian cells. <i>Toxicology</i> , 2020, 429, 152329.	2.0	13
6	Human CYP2E1-dependent mutagenicity of benzene and its hydroxylated metabolites in V79 derived cells: Suppression and enhancement by ethanol pretreatment. <i>Environmental and Molecular Mutagenesis</i> , 2020, 61, 622-634.	0.9	6
7	Interleukin-22 protects intestinal stem cells against genotoxic stress. <i>Nature</i> , 2019, 566, 249-253.	13.7	251
8	Human CYP3A4-mediated toxification of the pyrrolizidine alkaloid lasiocarpine. <i>Food and Chemical Toxicology</i> , 2019, 130, 79-88.	1.8	35
9	1-Methoxy-3-indolylmethyl DNA adducts in six tissues, and blood protein adducts, in mice under pak choi diet: time course and persistence. <i>Archives of Toxicology</i> , 2019, 93, 1515-1527.	1.9	5
10	Mutagenic Activity of Nitrosodiethylamine in Cell Lines Expressing Human CYP2E1 Adequacy of Dimethylsulfoxide as Solvent. <i>Environmental and Molecular Mutagenesis</i> , 2019, 60, 214-226.	0.9	5
11	Strong impact of sulfotransferases on DNA adduct formation by 4-aminobiphenyl in bladder and liver in mice. <i>Cancer Medicine</i> , 2018, 7, 5604-5610.	1.3	6
12	Hemoglobin adducts of furfuryl alcohol in genetically modified mouse models: Role of endogenous sulfotransferases 1a1 and 1d1 and transgenic human sulfotransferases 1A1/1A2. <i>Toxicology Letters</i> , 2018, 295, 173-178.	0.4	10
13	Role of human sulfotransferase 1A1 and N-acetyltransferase 2 in the metabolic activation of 16 heterocyclic amines and related heterocyclics to genotoxicants in recombinant V79 cells. <i>Archives of Toxicology</i> , 2017, 91, 3175-3184.	1.9	30
14	Potent mutagenicity of some non-planar tri- and tetrachlorinated biphenyls in mammalian cells, human CYP2E1 being a major activating enzyme. <i>Archives of Toxicology</i> , 2017, 91, 2663-2676.	1.9	27
15	Methyleugenol DNA adducts in human liver are associated with SULT1A1 copy number variations and expression levels. <i>Archives of Toxicology</i> , 2017, 91, 3329-3339.	1.9	30
16	Impact of genetic modulation of SULT1A enzymes on DNA adduct formation by aristolochic acids and 3-nitrobenzanthrone. <i>Archives of Toxicology</i> , 2017, 91, 1957-1975.	1.9	22
17	An in vitro study on the genotoxic effect of substituted furans in cells transfected with human metabolizing enzymes: 2,5-dimethylfuran and furfuryl alcohol. <i>Mutagenesis</i> , 2016, 31, 597-602.	1.0	5
18	Metabolism and excretion of 1-hydroxymethylpyrene, the proximate metabolite of the carcinogen 1-methylpyrene, in rats. <i>Toxicology</i> , 2016, 366-367, 43-52.	2.0	7

#	ARTICLE	IF	CITATIONS
19	Role of exposure/recovery schedule in micronuclei induction by several promutagens in V79-derived cells expressing human CYP2E1 and SULT1A1. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2016, 808, 27-37.	0.9	16
20	Characteristic single glucosinolates from <i>Moringa oleifera</i> : Induction of detoxifying enzymes and lack of genotoxic activity in various model systems. <i>Food and Function</i> , 2016, 7, 4660-4674.	2.1	10
21	Human CYP2E1-dependent mutagenicity of mono- and dichlorobiphenyls in Chinese hamster (V79)-derived cells. <i>Chemosphere</i> , 2016, 144, 1908-1915.	4.2	12
22	DNA adducts induced by food mutagen PhIP in a mouse model expressing human sulfotransferases 1A1 and 1A2. <i>Toxicology Letters</i> , 2016, 248, 34-38.	0.4	6
23	Modified Ames test using a strain expressing human sulfotransferase 1C2 to assess the mutagenicity of methyleugenol. <i>Genes and Environment</i> , 2016, 38, 1.	0.9	28
24	Use of genetically manipulated <i>S. almonella typhimurium</i> strains to evaluate the role of human sulfotransferases in the bioactivation of nitro- and aminotoluenes. <i>Environmental and Molecular Mutagenesis</i> , 2016, 57, 299-311.	0.9	10
25	Ethanol and 4-methylpyrazole increase DNA adduct formation of furfuryl alcohol in FVB/N wild-type mice and in mice expressing human sulfotransferases 1A1/1A2. <i>Carcinogenesis</i> , 2016, 37, 314-319.	1.3	8
26	In Silico Prediction of Human Sulfotransferase 1E1 Activity Guided by Pharmacophores from Molecular Dynamics Simulations. <i>Journal of Biological Chemistry</i> , 2016, 291, 58-71.	1.6	27
27	Conversion of Suspected Food Carcinogen 5-Hydroxymethylfurfural by Sulfotransferases and Aldehyde Dehydrogenases in Postmitochondrial Tissue Preparations of Humans, Mice, and Rats. <i>Toxicological Sciences</i> , 2016, 149, 192-201.	1.4	7
28	Bioactivation of food genotoxicants 5-hydroxymethylfurfural and furfuryl alcohol by sulfotransferases from human, mouse and rat: a comparative study. <i>Archives of Toxicology</i> , 2016, 90, 137-148.	1.9	37
29	The glucosinolate metabolite 1-methoxy- β -indolylmethyl alcohol induces a gene expression profile in mouse liver similar to the expression signature caused by known genotoxic hepatocarcinogens. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 685-697.	1.5	12
30	Genotoxicity of three food processing contaminants in transgenic mice expressing human sulfotransferases 1A1 and 1A2 as assessed by the in vivo alkaline single cell gel electrophoresis assay. <i>Environmental and Molecular Mutagenesis</i> , 2015, 56, 709-714.	0.9	30
31	Simultaneous Detection of Multiple DNA Adducts in Human Lung Samples by Isotope-Dilution UPLC-MS/MS. <i>Analytical Chemistry</i> , 2015, 87, 641-648.	3.2	59
32	Formation of DNA adducts in wild-type and transgenic mice expressing human sulfotransferases 1A1 and 1A2 after oral exposure to furfuryl alcohol. <i>Mutagenesis</i> , 2015, 30, 643-649.	1.0	9
33	Glucosinolates Are Mainly Absorbed Intact in Germfree and Human Microbiota-Associated Mice. <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 8418-8428.	2.4	22
34	Sulfotransferases. , 2015, , 1-5.		0
35	Sulfotransferases. , 2015, , 4392-4396.		0
36	The carcinogen 1-methylpyrene forms benzylic DNA adducts in mouse and rat tissues in vivo via a reactive sulphuric acid ester. <i>Archives of Toxicology</i> , 2014, 88, 815-21.	1.9	23

#	ARTICLE	IF	CITATIONS
37	A secondary metabolite of Brassicales, 1-methoxy-3-indolylmethyl glucosinolate, as well as its degradation product, 1-methoxy-3-indolylmethyl alcohol, forms DNA adducts in the mouse, but in varying tissues and cells. Archives of Toxicology, 2014, 88, 823-36.	1.9	17
38	The effect of knockout of sulfotransferases 1a1 and 1d1 and of transgenic human sulfotransferases 1A1/1A2 on the formation of DNA adducts from furfuryl alcohol in mouse models. Carcinogenesis, 2014, 35, 2339-2345.	1.3	23
39	Formation of hepatic DNA adducts by methyleugenol in mouse models: drastic decrease by Sult1a1 knockout and strong increase by transgenic human SULT1A1/2. Carcinogenesis, 2014, 35, 935-941.	1.3	50
40	Sulfotransferase-independent genotoxicity of illudin S and its acylfulvene derivatives in bacterial and mammalian cells. Archives of Toxicology, 2014, 88, 161-169.	1.9	8
41	High mutagenic activity of juice from pak choi (<i>Brassica rapa</i> ssp. <i>chinensis</i>) sprouts due to its content of 1-methoxy-3-indolylmethyl glucosinolate, and its enhancement by elicitation with methyl jasmonate. Food and Chemical Toxicology, 2014, 67, 10-16.	1.8	36
42	Determination of Sulfotransferase Forms Involved in the Metabolic Activation of the Genotoxicant 1-Hydroxymethylpyrene Using Bacterially Expressed Enzymes and Genetically Modified Mouse Models. Chemical Research in Toxicology, 2014, 27, 1060-1069.	1.7	34
43	Identification and Quantification of Protein Adducts Formed by Metabolites of 1-Methoxy-3-indolylmethyl Glucosinolate <i>in Vitro</i> and in Mouse Models. Chemical Research in Toxicology, 2014, 27, 188-199.	1.7	14
44	Glucosinolates from pak choi and broccoli induce enzymes and inhibit inflammation and colon cancer differently. Food and Function, 2014, 5, 1073-1081.	2.1	70
45	Tox-Box: securing drops of life - an enhanced health-related approach for risk assessment of drinking water in Germany. Environmental Sciences Europe, 2013, 25, .	2.6	30
46	Genotypic Variation of the Glucosinolate Profile in Pak Choi (<i>Brassica rapa</i> ssp. <i>chinensis</i>). Journal of Agricultural and Food Chemistry, 2013, 61, 1943-1953.	2.4	74
47	Optimized enzymatic hydrolysis of DNA for LC-MS/MS analyses of adducts of 1-methoxy-3-indolylmethyl glucosinolate and methyleugenol. Analytical Biochemistry, 2013, 434, 4-11.	1.1	27
48	Highly selective bioactivation of 1- and 2-hydroxy-3-methylcholanthrene to mutagens by individual human and other mammalian sulphotransferases expressed in <i>Salmonella typhimurium</i> . Mutagenesis, 2013, 28, 609-619.	1.0	15
49	Induced Production of 1-Methoxy-indol-3-ylmethyl Glucosinolate by Jasmonic Acid and Methyl Jasmonate in Sprouts and Leaves of Pak Choi (<i>Brassica rapa</i> ssp. <i>chinensis</i>). International Journal of Molecular Sciences, 2013, 14, 14996-15016.	1.8	67
50	Abundance of DNA adducts of methyleugenol, a rodent hepatocarcinogen, in human liver samples. Carcinogenesis, 2013, 34, 1025-1030.	1.3	50
51	Identification of human and murine sulfotransferases able to activate hydroxylated metabolites of methyleugenol to mutagens in <i>Salmonella typhimurium</i> and detection of associated DNA adducts using UPLC-MS/MS methods. Mutagenesis, 2012, 27, 453-462.	1.0	64
52	Hydroxymethyl-substituted furans: mutagenicity in <i>Salmonella typhimurium</i> strains engineered for expression of various human and rodent sulphotransferases. Mutagenesis, 2012, 27, 41-48.	1.0	51
53	Detection of DNA Adducts Originating from 1-Methoxy-3-indolylmethyl Glucosinolate Using Isotope-Dilution UPLC-ESI-MS/MS. Analytical Chemistry, 2012, 84, 6256-6262.	3.2	17
54	Mutagenicity of 5-Hydroxymethylfurfural in V79 Cells Expressing Human SULT1A1: Identification and Mass Spectrometric Quantification of DNA Adducts Formed. Chemical Research in Toxicology, 2012, 25, 1484-1492.	1.7	65

#	ARTICLE	IF	CITATIONS
55	Intestinal carcinogenesis of two food processing contaminants, 2-aminomethyl-6-phenylimidazo[4,5-b]pyridine and 5-hydroxymethylfurfural, in transgenic FVB mice expressing human sulfotransferases. <i>Molecular Carcinogenesis</i> , 2012, 51, 984-992.		12
56	Study of 5-hydroxymethylfurfural and its metabolite 5-sulfoxymethylfurfural on induction of colonic aberrant crypt foci in wild-type mice and transgenic mice expressing human sulfotransferases 1A1 and 1A2. <i>Molecular Nutrition and Food Research</i> , 2012, 56, 593-600.	1.5	14
57	Metabolism of Methyleugenol in Liver Microsomes and Primary Hepatocytes: Pattern of Metabolites, Cytotoxicity, and DNA-Adduct Formation. <i>Toxicological Sciences</i> , 2012, 129, 21-34.	1.4	40
58	Toxicity studies with 5-hydroxymethylfurfural and its metabolite 5-sulphoxymethylfurfural in wild-type mice and transgenic mice expressing human sulphotransferases 1A1 and 1A2. <i>Archives of Toxicology</i> , 2012, 86, 701-711.	1.9	41
59	First Chemical Synthesis and in Vitro Characterization of the Potential Human Metabolites 5-O-Feruloylquinic Acid 4-Sulfate and 4-O-Glucuronide. <i>Journal of Agricultural and Food Chemistry</i> , 2011, 59, 5671-5676.	2.4	13
60	1-Methoxy-3-indolylmethyl glucosinolate; a potent genotoxicant in bacterial and mammalian cells: Mechanisms of bioactivation. <i>Chemico-Biological Interactions</i> , 2011, 192, 81-86.	1.7	40
61	Identification of glucosinolate congeners able to form DNA adducts and to induce mutations upon activation by myrosinase. <i>Molecular Nutrition and Food Research</i> , 2011, 55, 783-792.	1.5	50
62	Metabolic activation of furfuryl alcohol: formation of 2-methylfuranyl DNA adducts in Salmonella typhimurium strains expressing human sulfotransferase 1A1 and in FVB/N mice. <i>Carcinogenesis</i> , 2011, 32, 1533-1539.	1.3	45
63	Detection of genotoxicants in Brassicales using endogenous DNA as a surrogate target and adducts determined by 32P-postlabelling as an experimental end point. <i>Mutagenesis</i> , 2011, 26, 407-413.	1.0	27
64	Altered tissue distribution of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine-DNA adducts in mice transgenic for human sulfotransferases 1A1 and 1A2. <i>Carcinogenesis</i> , 2011, 32, 1734-1740.	1.3	29
65	Sulfotransferases. , 2011, , 3558-3561.		0
66	Human cytochrome P450 2E1 and sulfotransferase 1A1 coexpressed in Chinese hamster V79 cells enhance spontaneous mutagenesis. <i>Environmental and Molecular Mutagenesis</i> , 2010, 51, 23-30.	0.9	19
67	The Friedreich's ataxia protein frataxin modulates DNA base excision repair in prokaryotes and mammals. <i>Biochemical Journal</i> , 2010, 432, 165-172.	1.7	34
68	Constitutive expression of bioactivating enzymes in normal human prostate suggests a capability to activate pro-carcinogens to DNA-damaging metabolites. <i>Prostate</i> , 2010, 70, 1586-1599.	1.2	35
69	Breakdown products of neoglucobrassicin inhibit activation of Nrf2 target genes mediated by myrosinase-derived glucoraphanin hydrolysis products. <i>Biological Chemistry</i> , 2010, 391, 1281-93.	1.2	39
70	Phase II Metabolism of Hesperetin by Individual UDP-Glucuronosyltransferases and Sulfotransferases and Rat and Human Tissue Samples. <i>Drug Metabolism and Disposition</i> , 2010, 38, 617-625.	1.7	86
71	Comparison of murine and human estrogen sulfotransferase inhibition in vitro and in silico—Implications for differences in activity, subunit dimerization and substrate inhibition. <i>Molecular and Cellular Endocrinology</i> , 2010, 317, 127-140.	1.6	18
72	The Suggested Physiologic Aryl Hydrocarbon Receptor Activator and Cytochrome P4501 Substrate 6-Formylindolo[3,2-b]carbazole Is Present in Humans. <i>Journal of Biological Chemistry</i> , 2009, 284, 2690-2696.	1.6	239

#	ARTICLE	IF	CITATIONS
73	Impact of Gut Microbiota on Intestinal and Hepatic Levels of Phase 2 Xenobiotic-Metabolizing Enzymes in the Rat. <i>Drug Metabolism and Disposition</i> , 2009, 37, 1179-1186.	1.7	84
74	Renal organic anion transporters OAT1 and OAT3 mediate the cellular accumulation of 5-sulfooxymethylfurfural, a reactive, nephrotoxic metabolite of the Maillard product 5-hydroxymethylfurfural. <i>Biochemical Pharmacology</i> , 2009, 78, 414-419.	2.0	59
75	Probenecid, an inhibitor of transmembrane organic anion transporters, alters tissue distribution of DNA adducts in 1-hydroxymethylpyrene-treated rats. <i>Toxicology</i> , 2009, 262, 80-85.	2.0	26
76	Molecular evidence for an involvement of organic anion transporters (OATs) in aristolochic acid nephropathy. <i>Toxicology</i> , 2009, 264, 74-79.	2.0	68
77	Effect of pentachlorophenol and 2,6-dichloro-4-nitrophenol on the activity of cDNA-expressed human alcohol and aldehyde dehydrogenases. <i>Toxicology Letters</i> , 2009, 191, 360-364.	0.4	5
78	Conversion of the Common Food Constituent 5-Hydroxymethylfurfural into a Mutagenic and Carcinogenic Sulfuric Acid Ester in the Mouse in Vivo. <i>Chemical Research in Toxicology</i> , 2009, 22, 1123-1128.	1.7	85
79	5-Hydroxymethylfurfural and 5-sulfooxymethylfurfural increase adenoma and flat ACF number in the intestine of Min/+ mice. <i>Anticancer Research</i> , 2009, 29, 1921-6.	0.5	38
80	Oxidation of alcohols and reduction of aldehydes derived from methyl- and dimethylpyrenes by cDNA-expressed human alcohol dehydrogenases. <i>Toxicology</i> , 2008, 245, 65-75.	2.0	12
81	Consumption of Brussels sprouts protects peripheral human lymphocytes against 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) and oxidative DNA damage: results of a controlled human intervention trial. <i>Molecular Nutrition and Food Research</i> , 2008, 52, 330-341.		50
82	Efficient oxidation of promutagenic hydroxymethylpyrenes by cDNA-expressed human alcohol dehydrogenase ADH2 and its inhibition by various agents. <i>Biochemical Pharmacology</i> , 2008, 75, 527-537.	2.0	9
83	Mutagenicity of N-nitrosodiethanolamine in a V79-derived cell line expressing two human biotransformation enzymes. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2008, 643, 64-69.	0.4	23
84	Detoxification of promutagenic aldehydes derived from methylpyrenes by human aldehyde dehydrogenases ALDH2 and ALDH3A1. <i>Archives of Biochemistry and Biophysics</i> , 2008, 477, 196-205.	1.4	22
85	SULT1C3, an orphan sequence of the human genome, encodes an enzyme activating various promutagens. <i>Food and Chemical Toxicology</i> , 2008, 46, 1249-1256.	1.8	20
86	Time Course of Hepatic 1-Methylpyrene DNA Adducts in Rats Determined by Isotope Dilution LC-MS/MS and ³² P-Postlabeling. <i>Chemical Research in Toxicology</i> , 2008, 21, 2017-2025.	1.7	36
87	Sulfotransferase Forms Expressed in Human Intestinal Caco-2 and TC7 Cells at Varying Stages of Differentiation and Role in Benzo[<i>a</i>]pyrene Metabolism. <i>Drug Metabolism and Disposition</i> , 2008, 36, 276-283.	1.7	72
88	Sulfotransferases. , 2008, , 2857-2861.		0
89	Directing Role of Organic Anion Transporters in the Excretion of Mercapturic Acids of Alkylated Polycyclic Aromatic Hydrocarbons. <i>Drug Metabolism and Disposition</i> , 2007, 35, 1824-1831.	1.7	11
90	Mutagenicity and DNA Adduct Formation by the Urban Air Pollutant 2-Nitrobenzanthrone. <i>Toxicological Sciences</i> , 2007, 98, 445-457.	1.4	42

#	ARTICLE	IF	CITATIONS
91	Cloning and Functional Characterization of Human Sodium-dependent Organic Anion Transporter (SLC10A6). <i>Journal of Biological Chemistry</i> , 2007, 282, 19728-19741.	1.6	82
92	Identification and localization of soluble sulfotransferases in the human gastrointestinal tract. <i>Biochemical Journal</i> , 2007, 404, 207-215.	1.7	151
93	Strategy for genotoxicity testing – Metabolic considerations. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2007, 627, 59-77.	0.9	100
94	Determination of alkylated and methylene-bridged polycyclic aromatic hydrocarbons in environmental matrices. <i>Toxicology Letters</i> , 2006, 164, S180.	0.4	1
95	Mutagenicity of arbutin in mammalian cells after activation by human intestinal bacteria. <i>Food and Chemical Toxicology</i> , 2006, 44, 1940-1947.	1.8	81
96	Development and validation of a fluorescence HPLC-based screening assay for inhibition of human estrogen sulfotransferase. <i>Analytical Biochemistry</i> , 2006, 357, 85-92.	1.1	21
97	Human cytochrome P450 reductase can act as a source of endogenous oxidative DNA damage and genetic instability. <i>Free Radical Biology and Medicine</i> , 2006, 40, 801-807.	1.3	10
98	Human sulphotransferases are involved in the activation of aristolochic acids and are expressed in renal target tissue. <i>International Journal of Cancer</i> , 2006, 118, 1090-1097.	2.3	79
99	STRUCTURAL ELUCIDATION OF HYDROXYLATED METABOLITES OF THE ISOFLAVAN EQUOL BY GAS CHROMATOGRAPHY-MASS SPECTROMETRY AND HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY-MASS SPECTROMETRY. <i>Drug Metabolism and Disposition</i> , 2006, 34, 51-60.	1.7	50
100	Uptake of Chemically Reactive, DNA-Damaging Sulfuric Acid Esters into Renal Cells by Human Organic Anion Transporters. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 1414-1421.	3.0	34
101	Biomarkers of Exposure, Effect, and Susceptibility in Workers Exposed to Nitrotoluenes. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2006, 15, 559-566.	1.1	36
102	Dietary resistant starch type 3 prevents tumor induction by 1,2-dimethylhydrazine and alters proliferation, apoptosis and dedifferentiation in rat colon. <i>Carcinogenesis</i> , 2006, 27, 1849-1859.	1.3	101
103	N-Acetyltransferase and sulfotransferase activity in human prostate: potential for carcinogen activation. <i>Pharmacogenetics and Genomics</i> , 2006, 16, 391-399.	0.7	16
104	Indicator Assays for Polycyclic Aromatic Hydrocarbon-Induced Genotoxicity. , 2005, , 283-314.		3
105	Polymorphisms in sulfotransferases SULT1A1 and SULT1A2 are not related to colorectal cancer. <i>International Journal of Cancer</i> , 2005, 113, 683-686.	2.3	25
106	Sulfotransferases and Acetyltransferases in Mutagenicity Testing: Technical Aspects. <i>Methods in Enzymology</i> , 2005, 400, 230-249.	0.4	30
107	V79-hCYP2E1-hSULT1A1, a cell line for the sensitive detection of genotoxic effects induced by carbohydrate pyrolysis products and other food-borne chemicals. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2005, 580, 41-52.	0.9	128
108	Activation and Inactivation of Carcinogens and Mutagens by Human Sulfotransferases. , 2005, , 279-304.		6

#	ARTICLE	IF	CITATIONS
109	Risk of Colorectal Adenomas in Relation to Meat Consumption, Meat Preparation, and Genetic Susceptibility in a Dutch Population. <i>Cancer Causes and Control</i> , 2004, 15, 225-236.	0.8	63
110	Pharmacogenetics of soluble sulfotransferases (SULTs). <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2004, 369, 55-68.	1.4	120
111	Effect of SULT1A1 and NAT2 genetic polymorphism on the association between cigarette smoking and colorectal adenomas. <i>International Journal of Cancer</i> , 2004, 108, 97-103.	2.3	56
112	Sulpho-conjugation of ethanol in humans in vivo and by individual sulphotransferase forms in vitro. <i>Biochemical Journal</i> , 2004, 383, 543-549.	1.7	51
113	Vitamin E activates gene expression via the pregnane X receptor. <i>Biochemical Pharmacology</i> , 2003, 65, 269-273.	2.0	213
114	Activation of 3-nitrobenzanthrone and its metabolites by human acetyltransferases, sulfotransferases and cytochrome P450 expressed in Chinese hamster V79 cells. <i>International Journal of Cancer</i> , 2003, 105, 583-592.	2.3	75
115	Extractionless method for the determination of urinary caffeine metabolites using high-performance liquid chromatography coupled with tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2003, 789, 227-237.	1.2	47
116	Bioactivation of the heterocyclic aromatic amine 2-amino-3-methyl-9H-pyrido [2,3-b]indole (MeA ⁺) in recombinant test systems expressing human xenobiotic-metabolizing enzymes. <i>Carcinogenesis</i> , 2003, 25, 801-807.	1.3	58
117	Use of genetically manipulated <i>Salmonella typhimurium</i> strains to evaluate the role of sulfotransferases and acetyltransferases in nitrofen mutagenicity. <i>Carcinogenesis</i> , 2003, 25, 779-786.	1.3	29
118	Characterization of rat iodothyronine sulfotransferases. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2003, 285, E592-E598.	1.8	34
119	Ethanol Enhances the Activation of 1-Hydroxymethylpyrene to DNA Adduct-Forming Species in the Rat. <i>Polycyclic Aromatic Compounds</i> , 2002, 22, 933-946.	1.4	18
120	Metabolic activation of the environmental contaminant 3-nitrobenzanthrone by human acetyltransferases and sulfotransferase. <i>Carcinogenesis</i> , 2002, 23, 1937-1945.	1.3	112
121	Stable expression of rat sulfotransferase 1B1 in V79 cells: activation of benzylic alcohols to mutagens. <i>Carcinogenesis</i> , 2002, 23, 1877-1884.	1.3	11
122	Activation of Polycyclic Aromatic Compounds by cDNA-Expressed Phase I and Phase II Enzymes. <i>Polycyclic Aromatic Compounds</i> , 2002, 22, 955-967.	1.4	9
123	Potent Inhibition of Estrogen Sulfotransferase by Hydroxylated Metabolites of Polyhalogenated Aromatic Hydrocarbons Reveals Alternative Mechanism for Estrogenic Activity of Endocrine Disrupters. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2002, 87, 1142-1150.	1.8	142
124	Differential activation of promutagens by alloenzymes of human sulfotransferase 1A2 expressed in <i>Salmonella typhimurium</i> . <i>Pharmacogenetics and Genomics</i> , 2002, 12, 677-689.	5.7	58
125	Sulphotransferases. , 2002, , 353-439.		26
126	Conjugation of 4-nitrophenol and 4-hydroxylonazolac in V79-derived cells expressing individual forms of human sulphotransferases. <i>Environmental Toxicology and Pharmacology</i> , 2002, 11, 243-250.	2.0	6

#	ARTICLE	IF	CITATIONS
127	Structure and Localization of the Human SULT1B1 Gene: Neighborhood to SULT1E1 and a SULT1D Pseudogene. <i>Biochemical and Biophysical Research Communications</i> , 2001, 288, 855-862.	1.0	28
128	N-Acetyltransferases, sulfotransferases and heterocyclic amine activation in the breast. <i>Pharmacogenetics and Genomics</i> , 2001, 11, 373-388.	5.7	44
129	Human cytosolic sulphotransferases: genetics, characteristics, toxicological aspects. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2001, 482, 27-40.	0.4	211
130	Detection of Mercapturic Acids and Nucleoside Adducts in Blood, Urine and Feces of Rats Treated with Metabolites of Methylpyrene. <i>Polycyclic Aromatic Compounds</i> , 2000, 21, 135-149.	1.4	7
131	Association between functional genetic polymorphisms of human sulfotransferases 1A1 and 1A2. <i>Pharmacogenetics and Genomics</i> , 2000, 10, 163-169.	5.7	66
132	Further studies on polycyclic arene sulfides. Preparation and mutagenic activity of 9, 10, 10a, 11a-tetrahydrotriphenylene[1,2-b]thiirene, 1a,2,3,10b-tetrahydro-5H-thiireno[3,4]benzo[1,2-b]fluorene and 1a,2,3,11b-tetrahydroacenaphtho[1,2,2':6,7]naphth[1,2-b]thiirene. <i>Journal of Heterocyclic Chemistry</i> , 2000, 37, 1109-1112.	1.4	2
133	Sulfotransferases in the bioactivation of xenobiotics. <i>Chemico-Biological Interactions</i> , 2000, 129, 141-170.	1.7	314
134	Potent Inhibition of Estrogen Sulfotransferase by Hydroxylated PCB Metabolites: A Novel Pathway Explaining the Estrogenic Activity of PCBs. <i>Endocrinology</i> , 2000, 141, 1897-1900.	1.4	322
135	Sulfotransferases: genetics and role in toxicology. <i>Toxicology Letters</i> , 2000, 112-113, 341-348.	0.4	111
136	Characterization of Human Iodothyronine Sulfotransferases1. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999, 84, 1357-1364.	1.8	73
137	Benzylic hydroxylation of 1-methylpyrene and 1-ethylpyrene by human and rat cytochromes P450 individually expressed in V79 Chinese hamster cells. <i>Carcinogenesis</i> , 1999, 20, 1777-1785.	1.3	52
138	Inhibitors of mammalian melanocyte tyrosinase: in vitro comparisons of alkyl esters of gentisic acid with other putative inhibitors. <i>Biochemical Pharmacology</i> , 1999, 57, 663-672.	2.0	301
139	Activation of propane 2-nitronate to a genotoxicant in V79-derived cell lines engineered for the expression of rat hepatic sulfotransferases. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 1999, 439, 191-197.	0.9	18
140	Direct optical resolution of trans-dihydrodiol enantiomers of fjord-region polycyclic aromatic hydrocarbons by high-performance liquid chromatography on a modified cellulose phase. <i>Journal of Chromatography A</i> , 1998, 822, 29-35.	1.8	4
141	Sulfotransferase-mediated activation of mutagens studied using heterologous expression systems. <i>Chemico-Biological Interactions</i> , 1998, 109, 195-219.	1.7	77
142	Expression of human estrogen sulfotransferase in <i>Salmonella typhimurium</i> : differences between hHST and hEST in the enantioselective activation of 1-hydroxyethylpyrene to a mutagen. <i>Chemico-Biological Interactions</i> , 1998, 109, 249-253.	1.7	31
143	Characterization of thyroid hormone sulfotransferases. <i>Chemico-Biological Interactions</i> , 1998, 109, 279-291.	1.7	65
144	Resolution of four k-region arene imines and mutagenicity of the optically pure aziridines. <i>Journal of Heterocyclic Chemistry</i> , 1998, 35, 39-43.	1.4	5

#	ARTICLE	IF	CITATIONS
145	Chiral Inversion of 1-Hydroxyethylpyrene Enantiomers Mediated by Enantioselective Sulfotransferases. <i>Biochemical and Biophysical Research Communications</i> , 1998, 247, 181-185.	1.0	17
146	Metabolic Activation of the (+)-S,S- and (âˆ’)-R,R-Enantiomers of trans-11,12-Dihydroxy-11,12-dihydrodibenzo[a,l]pyrene: A Stereoselectivity, DNA Adduct Formation, and Mutagenicity in Chinese Hamster V79 Cells. <i>Chemical Research in Toxicology</i> , 1997, 10, 1161-1170.	1.7	34
147	Bioactivation of mutagens via sulfation. <i>FASEB Journal</i> , 1997, 11, 314-321.	0.2	196
148	External Versus Internal Metabolic Activation of Polycyclic Aromatic Compounds in Mutagenicity Tests: A Comparison Using Heterologous Expression Systems. <i>Polycyclic Aromatic Compounds</i> , 1996, 10, 43-50.	1.4	4
149	Activation of benzylic alcohols to mutagens by rat and human sulfotransferases expressed in <i>Escherichia coli</i> . <i>European Journal of Pharmacology - Environmental Toxicology and Pharmacology Section</i> , 1995, 293, 173-181.	0.8	27
150	Human fetal adrenal hydroxysteroid sulphotransferase: cDNA cloning, stable expression in V79 cells and functional characterisation of the expressed enzyme. <i>Molecular and Cellular Endocrinology</i> , 1995, 112, 53-60.	1.6	41
151	Human Dehydroepiandrosterone Sulfotransferase.. <i>Annals of the New York Academy of Sciences</i> , 1995, 774, 59-72.	1.8	109
152	Salmonella strains and mammalian cells genetically engineered for expression of sulfotransferases. <i>Toxicology Letters</i> , 1995, 82-83, 829-834.	0.4	20
153	Substance-dependent sex differences in the activation of benzylic alcohols to mutagens by hepatic sulfotransferases of the rat. <i>Carcinogenesis</i> , 1994, 15, 2605-2611.	1.3	40
154	Synthesis and mutagenicity of the diastereomeric fjord-region 11,12-dihydrodiol 13,14-epoxides of dibenzo [a,l]pyrene. <i>Carcinogenesis</i> , 1994, 15, 2507-2516.	1.3	80
155	Fjord-region diol-epoxides of benzo[c]chrysene are potent inducers of micronuclei in murine bone marrow. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1994, 309, 37-43.	0.4	3
156	Mutagenicity of K-region oxides and imines of chrysene, benzo[c]phenanthrene and benzo[g]chrysene in <i>Salmonella typhimurium</i> . <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1994, 308, 135-141.	0.4	13
157	Comparison of common gene mutation tests in mammalian cells in culture: a position paper of the GUM Commission for the Development of Guidelines for Genotoxicity Testing. <i>Mutation Research - Environmental Mutagenesis and Related Subjects Including Methodology</i> , 1994, 313, 7-20.	0.4	21
158	The presence of KCl in the exposure medium strongly influences the mutagenicity of metabolites of polycyclic aromatic hydrocarbons in <i>Escherichia coli</i> . <i>Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1994, 324, 111-114.	1.2	0
159	1-hydroxymethylpyrene and its sulfuric acid ester: toxicological effects in vitro and in vivo, and metabolic aspects. <i>Chemico-Biological Interactions</i> , 1994, 92, 305-319.	1.7	20
160	Activation of promutagens by endogenous and heterologous sulfotransferases expressed in continuous cell cultures. <i>Toxicology Letters</i> , 1994, 72, 13-21.	0.4	33
161	Activation of benzylic alcohols to mutagens by human hepatic sulphotransferases. <i>Mutagenesis</i> , 1994, 9, 553-557.	1.0	34
162	K-region oxides and imines derived from alkylated benz[1±]anthracene congeners: synthesis, stability in aqueous media and mutagenicity. <i>Mutagenesis</i> , 1994, 9, 83-92.	1.0	13

#	ARTICLE	IF	CITATIONS
163	Stable expression of human cytochrome P450 1A1 cDNA in V79 Chinese hamster cells and metabolic activation of benzo[a]pyrene. <i>European Journal of Pharmacology - Environmental Toxicology and Pharmacology Section</i> , 1993, 248, 251-261.	0.8	45
164	Sulfotransferase-mediated mutagenicity of 1-hydroxymethylpyrene and 4H-cyclopenta[def]chrysen-4-ol and its enhancement by chloride anions. <i>Carcinogenesis</i> , 1993, 14, 599-602.	1.3	44
165	Development of hydroxysteroid sulfotransferase-deficient lesions during hepatocarcinogenesis in rats. <i>Carcinogenesis</i> , 1993, 14, 2267-2270.	1.3	8
166	Genotoxicity characteristics of reverse diol-epoxides of chrysene. <i>Carcinogenesis</i> , 1993, 14, 11-19.	1.3	31
167	DNA strand break induction, mutagenicity, and cytotoxicity of the mycotoxins 11- β -hydroxy-7-deoxy-rosenonolactone, rosenonolactone, and trichothecin. <i>Mycotoxin Research</i> , 1992, 8, 77-83.	1.3	1
168	Biphenyl and fluorinated derivatives: liver enzyme-mediated mutagenicity detected in <i>Salmonella typhimurium</i> and Chinese hamster V79 cells. <i>Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1992, 281, 151-156.	1.2	11
169	Influence of the level of cytosolic epoxide hydrolase on the induction of sister chromatid exchanges by trans- β -ethylstyrene 7,8-oxide in human lymphocytes. <i>Biochemical Pharmacology</i> , 1991, 42, 2147-2152.	2.0	12
170	Relationship between mutagenicity and DNA adduct formation in mammalian cells for fjord- and bay-region diol-epoxides of polycyclic aromatic hydrocarbons. <i>Chemico-Biological Interactions</i> , 1991, 80, 177-186.	1.7	35
171	Stable expression of rat cytochrome P450IA2 cDNA and hydroxylation of 17 β -estradiol and 2-aminofluorene in V79 Chinese hamster cells. <i>Molecular Carcinogenesis</i> , 1991, 4, 489-498.	1.3	40
172	Strong influence of the exposure medium on mutagenicity in the Ames test: 7-methylbenz[\pm]anthracene-5, 6-oxide. <i>Mutagenesis</i> , 1991, 6, 537-540.	1.0	7
173	Expression of xenobiotic-metabolizing enzymes in propagatable cell cultures and induction of micronuclei by 13 compounds. <i>Mutagenesis</i> , 1990, 5, 241-250.	1.0	103
174	Endogenous mutagens derived from amino acids. <i>Mutation Research - Reviews in Genetic Toxicology</i> , 1990, 238, 235-243.	3.0	25
175	Mutagenicity experiments on l-cysteine and d-penicillamine using V79 cells as indicators and for metabolic activation. <i>Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1990, 243, 187-193.	1.2	13
176	Mutagenicity spectra in <i>Salmonella typhimurium</i> strains of glutathione, L-cysteine and active oxygen species. <i>Mutagenesis</i> , 1989, 4, 221-227.	1.0	37
177	Characterization of an epithelial, nearly diploid liver cell strain, from Chinese hamster, able to activate promutagens. <i>Mutagenesis</i> , 1987, 2, 127-135.	1.0	30
178	Metabolic activation to a mutagen of 3-hydroxy-trans-7,8-dihydroxy-7,8-dihydrobenzo[a]pyrene, a secondary metabolite of benzo[a]pyrene. <i>Carcinogenesis</i> , 1987, 8, 1621-1627.	1.3	28
179	Mutagenicity of N-substituted phenanthrene 9,10-imines in <i>Salmonella typhimurium</i> and Chinese hamster V79 cells. <i>Environmental Mutagenesis</i> , 1986, 8, 829-837.	1.4	8
180	Cis- and trans-1,2-diphenylaziridines: induction of xenobiotic-metabolizing enzymes in rat liver and mutagenicity in <i>Salmonella typhimurium</i> . <i>Archives of Toxicology</i> , 1986, 59, 242-248.	1.9	7

#	ARTICLE	IF	CITATIONS
181	V79 Chinese hamster cells deacetylate Trans-N-acetoxy-4-acetylaminostilbene and Trans-N-hydroxy-4-acetylaminostilbene to mutagenic and cytotoxic metabolites. <i>Cell Biology and Toxicology</i> , 1986, 2, 213-221.	2.4	3
182	Xenobiotic metabolizing enzymes of rat liver nonparenchymal cells. <i>Toxicology and Applied Pharmacology</i> , 1986, 84, 500-511.	1.3	29
183	Determination of DNA single strand breaks and selective DNA amplification by N-nitrodimethylamine and analogs, and estimation of the indicator cells' metabolic capacities. <i>Journal of Cancer Research and Clinical Oncology</i> , 1986, 111, 123-128.	1.2	11
184	Metabolism of the bay-region diol-epoxide of chrysene to a triol-epoxide and the enzyme-catalysed conjugation of these epoxides with glutathione. <i>Carcinogenesis</i> , 1986, 7, 2095-2098.	1.3	8
185	Mutagenic potential of DNA adducts formed by diol-epoxides, triol-epoxides and the K-region epoxide of chrysene in mammalian cells. <i>Carcinogenesis</i> , 1986, 7, 1739-1743.	1.3	29
186	Propylidazine is mutagenic in <i>Salmonella typhimurium</i> and <i>Escherichia coli</i> : Distinct specificity for strains TA1537 AND TA97. <i>Teratogenesis, Carcinogenesis, and Mutagenesis</i> , 1985, 5, 339-345.	0.8	9
187	Host-mediated mutagenicity experiments with benzo[a]pyrene and two of its metabolites. <i>Mutation Research - Genetic Toxicology Testing and Biomonitoring of Environmental Or Occupational Exposure</i> , 1985, 156, 163-169.	1.2	9
188	Mutagenicity of cysteine and penicillamine and its enantiomeric selectivity. <i>Biochemical Pharmacology</i> , 1985, 34, 3725-3728.	2.0	45
189	Microsomal epoxide hydrolase in different rat strains. <i>Biochemical Pharmacology</i> , 1983, 32, 1783-1788.	2.0	18
190	Effects of various enzyme inducers on monooxygenase, glutathione S-transferase and epoxide hydrolase activities in cultured hepatoma cells. <i>Chemico-Biological Interactions</i> , 1982, 42, 27-43.	1.7	7
191	Ethanol- or acetone-pretreatment of mice strongly enhances the bacterial mutagenicity of dimethylnitrosamine in assays mediated by liver subcellular fraction, but not in host-mediated assays. <i>Carcinogenesis</i> , 1981, 2, 1057-1061.	1.3	24
192	The apparent ubiquity of epoxide hydratase in rat organs. <i>Biochemical Pharmacology</i> , 1977, 26, 603-607.	2.0	141
193	Dual role of epoxide hydratase in both activation and inactivation of benzo(a)pyrene. <i>Archives of Toxicology</i> , 1977, 39-39, 65-75.	1.9	90
194	Inactivation of electrophilic metabolites by glutathione S-transferases and limitation of the system due to subcellular localization. <i>Archives of Toxicology</i> , 1977, 39-39, 87-96.	1.9	80
195	Species differences in activating and inactivating enzymes related to the control of mutagenic metabolites. <i>Archives of Toxicology</i> , 1977, 39-39, 97-108.	1.9	52
196	Ethanol Enhances the Activation of 1-Hydroxymethylpyrene to DNA Adduct-Forming Species in the Rat. , 0, .		6
197	Activation of Polycyclic Aromatic Compounds by cDNA-Expressed Phase I and Phase II Enzymes. , 0, .		2