
Weiping Li

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5045450/publications.pdf Version: 2024-02-01



WEIDING

#	Article	IF	CITATIONS
1	High Triglyceride-Glucose Index is Associated with Poor Cardiovascular Outcomes in Nondiabetic Patients with ACS with LDL-C below 1.8 mmol/L. Journal of Atherosclerosis and Thrombosis, 2022, 29, 268-281.	0.9	32
2	Protective role of activating PPARÎ ³ in advanced glycation end products-induced impairment of coronary artery vasodilation via inhibiting p38 phosphorylation and reactive oxygen species production. Biomedicine and Pharmacotherapy, 2022, 147, 112641.	2.5	8
3	Association of Prior Statin Therapy With Cardiovascular Outcomes in Patients With Initial Diagnosis of OCAD and LDL-C Below 1.8Âmmol/L. Angiology, 2022, , 000331972210758.	0.8	0
4	Predictive effect of triglyceride‑glucose index on clinical events in patients with type 2 diabetes mellitus and acute myocardial infarction: results from an observational cohort study in China. Cardiovascular Diabetology, 2021, 20, 43.	2.7	40
5	Role of peroxisome proliferators-activated receptor-gamma in advanced glycation end product-mediated functional loss of voltage-gated potassium channel in rat coronary arteries. BMC Cardiovascular Disorders, 2020, 20, 337.	0.7	1
6	High triglyceride-glucose index is associated with adverse cardiovascular outcomes in patients with acute myocardial infarction. Nutrition, Metabolism and Cardiovascular Diseases, 2020, 30, 2351-2362.	1.1	46
7	Real-world use of ACEI/ARB in diabetic hypertensive patients before the initial diagnosis of obstructive coronary artery disease: patient characteristics andAlong-term follow-up outcome. Journal of Translational Medicine, 2020, 18, 150.	1.8	10
8	Underweight Predicts Greater Risk of Cardiac Mortality Post Acute Myocardial Infarction. International Heart Journal, 2020, 61, 658-664.	0.5	6
9	AGEs impair Kv channel-mediated vasodilation of coronary arteries by activating the NF-κB signaling pathway in ZDF rats. Biomedicine and Pharmacotherapy, 2019, 120, 109527.	2.5	9
10	Pioglitazone downregulates Twist-1 expression in the kidney and protects renal function of Zucker diabetic fatty rats. Biomedicine and Pharmacotherapy, 2019, 118, 109346.	2.5	9
11	Pregnancy-Associated Plasma Protein A Induces Inflammatory Cytokine Expression by Activating IGF-I/PI3K/Akt Pathways. Mediators of Inflammation, 2019, 2019, 1-12.	1.4	10
12	Liraglutide inhibited AGEs induced coronary smooth muscle cell phenotypic transition through inhibiting the NF-κB signal pathway. Peptides, 2019, 112, 125-132.	1.2	10
13	The Neutrophil Percentage to Albumin Ratio as a New Predictor of In-Hospital Mortality in Patients with ST-Segment Elevation Myocardial Infarction. Medical Science Monitor, 2019, 25, 7845-7852.	0.5	29
14	Telmisartan ameliorates adipoR1 and adipoR2 expression via PPAR-Î ³ activation in the coronary artery and VSMCs. Biomedicine and Pharmacotherapy, 2017, 95, 129-136.	2.5	12
15	Pioglitazone Attenuates Atherosclerosis in Diabetic Mice by Inhibition of Receptor for Advanced Glycation End-Product (RAGE) Signaling. Medical Science Monitor, 2017, 23, 6121-6131.	0.5	11
16	RhoA/rock signaling mediates peroxynitrite-induced functional impairment of Rat coronary vessels. BMC Cardiovascular Disorders, 2016, 16, 193.	0.7	10
17	Advanced Glycation End Products Impair Voltage-Gated K+ Channels-Mediated Coronary Vasodilation in Diabetic Rats. PLoS ONE, 2015, 10, e0142865.	1.1	12
18	CRP and TNF- <mml:math <br="" xmlns:mml="http://www.w3.org/1998/Math/MathML">id="M1"><mml:mrow><mml:mi mathvariant="bold-italic">α  </mml:mi></mml:mrow></mml:math> Indu PAPP-A Expression in Human Peripheral Blood Mononuclear Cells. Mediators of Inflammation, 2012, 2012, 1-9.	^{ICe} 1.4	8

#	Article	IF	CITATIONS
19	Pioglitazone prevents hyperglycemia induced decrease of AdipoR1 and AdipoR2 in coronary arteries and coronary VSMCs. Molecular and Cellular Endocrinology, 2012, 363, 27-35.	1.6	15