

Yosuke Amagai

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5056293/publications.pdf>

Version: 2024-02-01

17
papers

341
citations

1040056

9
h-index

888059

17
g-index

17
all docs

17
docs citations

17
times ranked

709
citing authors

#	ARTICLE	IF	CITATIONS
1	Skin pH Is the Master Switch of Kallikrein 5-Mediated Skin Barrier Destruction in a Murine Atopic Dermatitis Model. <i>Journal of Investigative Dermatology</i> , 2016, 136, 127-135.	0.7	92
2	Nuclear factor- κ B plays a critical role in both intrinsic and acquired resistance against endocrine therapy in human breast cancer cells. <i>Scientific Reports</i> , 2014, 4, 4057.	3.3	54
3	Dihomo- γ -linolenic acid prevents the development of atopic dermatitis through prostaglandin D1 production in NC/Tnd mice. <i>Journal of Dermatological Science</i> , 2015, 79, 30-37.	1.9	37
4	Mast cell hyperactivity underpins the development of oxygen-induced retinopathy. <i>Journal of Clinical Investigation</i> , 2017, 127, 3987-4000.	8.2	24
5	Dieckol, a phlorotannin of <i>Ecklonia cava</i> , suppresses IgE-mediated mast cell activation and passive cutaneous anaphylactic reaction. <i>Experimental Dermatology</i> , 2015, 24, 968-970.	2.9	23
6	Silencing of int6 gene restores function of the ischaemic hindlimb in a rat model of peripheral arterial disease. <i>Cardiovascular Research</i> , 2011, 92, 209-217.	3.8	18
7	Establishment of a novel high-affinity IgE receptor-positive canine mast cell line with wild-type c-kit receptors. <i>Biochemical and Biophysical Research Communications</i> , 2008, 366, 857-861.	2.1	14
8	Abnormalities in Itch Sensation and Skin Barrier Function in Atopic NC/Tnd Mice. <i>Biological and Pharmaceutical Bulletin</i> , 2013, 36, 1248-1252.	1.4	10
9	Production of stem cell factor in canine mast cell tumors. <i>Research in Veterinary Science</i> , 2014, 96, 124-126.	1.9	10
10	A point mutation in the extracellular domain of KIT promotes tumorigenesis of mastcells via ligand-independent auto-dimerization. <i>Scientific Reports</i> , 2015, 5, 9775.	3.3	10
11	The Phosphoinositide 3-Kinase Pathway Is Crucial for the Growth of Canine Mast Cell Tumors. <i>Journal of Veterinary Medical Science</i> , 2013, 75, 791-794.	0.9	9
12	Oral Administration of Fermented Probiotics Improves the Condition of Feces in Adult Horses. <i>Journal of Equine Science</i> , 2014, 25, 65-72.	0.8	8
13	Amelioration of atopic-like skin conditions in NC/Tnd mice by topical application with distilled <i>Alpinia intermedia</i> Gagnep extracts. <i>Journal of Dermatology</i> , 2017, 44, 1238-1247.	1.2	8
14	Topical Application of Ketoprofen Improves Gait Disturbance in Rat Models of Acute Inflammation. <i>BioMed Research International</i> , 2013, 2013, 1-7.	1.9	7
15	Nerve growth factor promotes killing of <i>Leishmania donovani</i> by macrophages through the induction of hydrogen peroxide. <i>Microbes and Infection</i> , 2014, 16, 702-706.	1.9	7
16	Swing time ratio, a new parameter of gait disturbance, for the evaluation of the severity of neuropathic pain in a rat model of partial sciatic nerve ligation. <i>Journal of Pharmacological and Toxicological Methods</i> , 2016, 79, 7-14.	0.7	6
17	Increased Expression of the Antiapoptotic Protein MCL1 in Canine Mast Cell Tumors. <i>Journal of Veterinary Medical Science</i> , 2013, 75, 971-974.	0.9	4