

Sheng Hu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5060313/publications.pdf>

Version: 2024-02-01

14
papers

178
citations

1478505

6
h-index

1199594

12
g-index

14
all docs

14
docs citations

14
times ranked

278
citing authors

#	ARTICLE	IF	CITATIONS
1	Basal ganglia-orbitofrontal circuits are associated with prospective memory deficits in Wilson's disease. <i>Brain Imaging and Behavior</i> , 2022, 16, 141-150.	2.1	5
2	Structural and Functional Changes Are Related to Cognitive Status in Wilson's Disease. <i>Frontiers in Human Neuroscience</i> , 2021, 15, 610947.	2.0	9
3	The degeneration changes of basal forebrain are associated with prospective memory impairment in patients with Wilson's disease. <i>Brain and Behavior</i> , 2021, 11, e2239.	2.2	5
4	Acupuncture Treatment Decreased Temporal Variability of Dynamic Functional Connectivity in Chronic Tinnitus. <i>Frontiers in Neuroscience</i> , 2021, 15, 737993.	2.8	5
5	Functional MRI Investigation of Ultrasound Stimulation at ST 36. <i>Evidence-based Complementary and Alternative Medicine</i> , 2020, 2020, 1-7.	1.2	4
6	Comprehensive Analysis of CDCAs Methylation and Immune Infiltrates in Hepatocellular Carcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 566183.	2.8	8
7	Longitudinal Changes in Functional Connectivity of the Caudate Is Associated With Recovery From Bell's Palsy. <i>Frontiers in Aging Neuroscience</i> , 2019, 11, 295.	3.4	1
8	Aberrant Coupling Between Resting-State Cerebral Blood Flow and Functional Connectivity in Wilson's Disease. <i>Frontiers in Neural Circuits</i> , 2019, 13, 25.	2.8	13
9	Acute and Sustained Effects on Human Brain Induced by Different Modalities of Acupuncture: An fMRI Study. , 2019, , .		1
10	The P2P Solution Research and Design Based on NAT Traversing Technology. , 2019, , .		5
11	Functional Connectivity Modulation by Acupuncture in Patients with Bell's Palsy. <i>Evidence-based Complementary and Alternative Medicine</i> , 2016, 2016, 1-10.	1.2	12
12	Long-term cognitive impairment of breast cancer patients after chemotherapy: A functional MRI study. <i>European Journal of Radiology</i> , 2016, 85, 1053-1057.	2.6	47
13	Functional connectivity change of brain default mode network in breast cancer patients after chemotherapy. <i>Neuroradiology</i> , 2016, 58, 921-928.	2.2	46
14	Increasing functional connectivity of the anterior cingulate cortex during the course of recovery from Bell's palsy. <i>NeuroReport</i> , 2015, 26, 6-12.	1.2	17