

Leena Sharma

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5062935/leena-sharma-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

84
papers

7,925
citations

47
h-index

89
g-index

89
ext. papers

8,869
ext. citations

5.1
avg, IF

6.13
L-index

#	Paper	IF	Citations
84	Osteoarthritis of the Knee. <i>New England Journal of Medicine</i> , 2021 , 384, 51-59	59.2	91
83	Is Lamellar Cartilage Composition as Determined by T2 Relaxometry Associated with Incident and Worsening of Cartilage or Bone Marrow Abnormalities?. <i>Cartilage</i> , 2020 , 1947603520932197	3	1
82	Examining Timeliness of Total Knee Replacement Among Patients with Knee Osteoarthritis in the U.S.: Results from the OAI and MOST Longitudinal Cohorts. <i>Journal of Bone and Joint Surgery - Series A</i> , 2020 , 102, 468-476	5.6	23
81	Development and validation of risk stratification trees for incident slow gait speed in persons at high risk for knee osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 1412-1419	2.4	3
80	Hip muscle strength and protection against structural worsening and poor function and disability outcomes in knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2019 , 27, 885-894	6.2	8
79	Association of Varus Knee Thrust During Walking With Worsening Western Ontario and McMaster Universities Osteoarthritis Index Knee Pain: A Prospective Cohort Study. <i>Arthritis Care and Research</i> , 2019 , 71, 1353-1359	4.7	6
78	Physical Activity and Worsening of Radiographic Findings in Persons With or at Higher Risk of Knee Osteoarthritis. <i>Arthritis Care and Research</i> , 2019 , 71, 198-206	4.7	7
77	Varus thrust during walking and the risk of incident and worsening medial tibiofemoral MRI lesions: the Multicenter Osteoarthritis Study. <i>Osteoarthritis and Cartilage</i> , 2017 , 25, 839-845	6.2	25
76	Knee tissue lesions and prediction of incident knee osteoarthritis over 7 years in a cohort of persons at higher risk. <i>Osteoarthritis and Cartilage</i> , 2017 , 25, 1068-1075	6.2	20
75	Predictive and concurrent validity of cartilage thickness change as a marker of knee osteoarthritis progression: data from the Osteoarthritis Initiative. <i>Osteoarthritis and Cartilage</i> , 2017 , 25, 2063-2071	6.2	25
74	Varus Thrust and Incident and Progressive Knee Osteoarthritis. <i>Arthritis and Rheumatology</i> , 2017 , 69, 2136-2143	9.5	37
73	Association of baseline knee sagittal dynamic joint stiffness during gait and 2-year patellofemoral cartilage damage worsening in knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2017 , 25, 242-248	6.2	11
72	Clinical significance of worsening versus stable preradiographic MRI lesions in a cohort study of persons at higher risk for knee osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1630-6	2.4	26
71	Osteoarthritis year in review 2015: clinical. <i>Osteoarthritis and Cartilage</i> , 2016 , 24, 36-48	6.2	59
70	Comparison of radiographic joint space width and magnetic resonance imaging for prediction of knee replacement: A longitudinal case-control study from the Osteoarthritis Initiative. <i>European Radiology</i> , 2016 , 26, 1942-51	8	26
69	Baseline radiographic osteoarthritis and semi-quantitatively assessed meniscal damage and extrusion and cartilage damage on MRI is related to quantitatively defined cartilage thickness loss in knee osteoarthritis: the Multicenter Osteoarthritis Study. <i>Osteoarthritis and Cartilage</i> , 2015 , 23, 2191-2198	6.2	39
68	Sedentary behavior and physical function: objective evidence from the Osteoarthritis Initiative. <i>Arthritis Care and Research</i> , 2015 , 67, 366-73	4.7	89

67	Knee Instability and Basic and Advanced Function Decline in Knee Osteoarthritis. <i>Arthritis Care and Research</i> , 2015 , 67, 1095-102	4.7	17
66	External knee adduction and flexion moments during gait and medial tibiofemoral disease progression in knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2015 , 23, 1099-106	6.2	144
65	Trajectory of cartilage loss within 4 years of knee replacement--a nested case-control study from the osteoarthritis initiative. <i>Osteoarthritis and Cartilage</i> , 2014 , 22, 1542-9	6.2	30
64	Factors associated with pain experience outcome in knee osteoarthritis. <i>Arthritis Care and Research</i> , 2014 , 66, 1828-35	4.7	31
63	Significance of preradiographic magnetic resonance imaging lesions in persons at increased risk of knee osteoarthritis. <i>Arthritis and Rheumatology</i> , 2014 , 66, 1811-9	9.5	64
62	Relationship of meeting physical activity guidelines with health-related utility. <i>Arthritis Care and Research</i> , 2014 , 66, 1041-7	4.7	18
61	Physical activity, alignment and knee osteoarthritis: data from MOST and the OAI. <i>Osteoarthritis and Cartilage</i> , 2013 , 21, 789-95	6.2	38
60	Obesity and other modifiable factors for physical inactivity measured by accelerometer in adults with knee osteoarthritis. <i>Arthritis Care and Research</i> , 2013 , 65, 53-61	4.7	63
59	Racial and ethnic differences in physical activity guidelines attainment among people at high risk of or having knee osteoarthritis. <i>Arthritis Care and Research</i> , 2013 , 65, 195-202	4.7	28
58	Excess body weight and four-year function outcomes: comparison of African Americans and whites in a prospective study of osteoarthritis. <i>Arthritis Care and Research</i> , 2013 , 65, 5-14	4.7	23
57	Valgus malalignment is a risk factor for lateral knee osteoarthritis incidence and progression: findings from the Multicenter Osteoarthritis Study and the Osteoarthritis Initiative. <i>Arthritis and Rheumatism</i> , 2013 , 65, 355-62		159
56	Varus thrust and knee frontal plane dynamic motion in persons with knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2013 , 21, 1668-73	6.2	52
55	The role of varus and valgus alignment in the initial development of knee cartilage damage by MRI: the MOST study. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 235-40	2.4	125
54	How do short-term rates of femorotibial cartilage change compare to long-term changes? Four year follow-up data from the osteoarthritis initiative. <i>Osteoarthritis and Cartilage</i> , 2012 , 20, 1250-7	6.2	28
53	Knee malalignment is associated with an increased risk for incident and enlarging bone marrow lesions in the more loaded compartments: the MOST study. <i>Osteoarthritis and Cartilage</i> , 2012 , 20, 1227-33	6.2	62
52	Patterns of compartment involvement in tibiofemoral osteoarthritis in men and women and in whites and African Americans. <i>Arthritis Care and Research</i> , 2012 , 64, 847-52	4.7	100
51	Knee confidence as it relates to physical function outcome in persons with or at high risk of knee osteoarthritis in the osteoarthritis initiative. <i>Arthritis and Rheumatism</i> , 2012 , 64, 1437-46		36
50	Breaking the Law of Valgus: the surprising and unexplained prevalence of medial patellofemoral cartilage damage. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1827-32	2.4	37

49	Superficial femoral artery plaque and functional performance in peripheral arterial disease: walking and leg circulation study (WALCS III). <i>JACC: Cardiovascular Imaging</i> , 2011 , 4, 730-9	8.4	25
48	Physical activity levels and functional performance in the osteoarthritis initiative: a graded relationship. <i>Arthritis and Rheumatism</i> , 2011 , 63, 127-36		111
47	Varus-valgus alignment: reduced risk of subsequent cartilage loss in the less loaded compartment. <i>Arthritis and Rheumatism</i> , 2011 , 63, 1002-9		30
46	Objective physical activity measurement in the osteoarthritis initiative: Are guidelines being met?. <i>Arthritis and Rheumatism</i> , 2011 , 63, 3372-82		155
45	Risk factors for medial meniscal pathology on knee MRI in older US adults: a multicentre prospective cohort study. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 1733-9	2.4	83
44	Subregional effects of meniscal tears on cartilage loss over 2 years in knee osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 74-9	2.4	53
43	Subchondral bone attrition may be a reflection of compartment-specific mechanical load: the MOST Study. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 841-4	2.4	55
42	Varus and valgus alignment and incident and progressive knee osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1940-5	2.4	254
41	Moving to maintain function in knee osteoarthritis: evidence from the osteoarthritis initiative. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010 , 91, 714-21	2.8	42
40	Quadriceps weakness predicts risk for knee joint space narrowing in women in the MOST cohort. <i>Osteoarthritis and Cartilage</i> , 2010 , 18, 769-75	6.2	158
39	Comment on: Varus malalignment negates the structure-modifying benefits of doxycycline in obese women with knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2010 , 18, 1006-7	6.2	4
38	Within-subregion relationship between bone marrow lesions and subsequent cartilage loss in knee osteoarthritis. <i>Arthritis Care and Research</i> , 2010 , 62, 198-203	4.7	34
37	Frequency of varus and valgus thrust and factors associated with thrust presence in persons with or at higher risk of developing knee osteoarthritis. <i>Arthritis and Rheumatism</i> , 2010 , 62, 1403-11		62
36	Denuded subchondral bone and knee pain in persons with knee osteoarthritis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 3703-10		58
35	Medial-to-lateral ratio of tibiofemoral subchondral bone area is adapted to alignment and mechanical load. <i>Calcified Tissue International</i> , 2009 , 84, 186-94	3.9	42
34	Associations of borderline and low normal ankle-brachial index values with functional decline at 5-year follow-up: the WALCS (Walking and Leg Circulation Study). <i>Journal of the American College of Cardiology</i> , 2009 , 53, 1056-62	15.1	142
33	Knee extensor strength does not protect against incident knee symptoms at 30 months in the multicenter knee osteoarthritis (MOST) cohort. <i>PM and R</i> , 2009 , 1, 459-65	2.2	33
32	Asymptomatic peripheral arterial disease is associated with more adverse lower extremity characteristics than intermittent claudication. <i>Circulation</i> , 2008 , 117, 2484-91	16.7	113

31	Relationship of meniscal damage, meniscal extrusion, malalignment, and joint laxity to subsequent cartilage loss in osteoarthritic knees. <i>Arthritis and Rheumatism</i> , 2008 , 58, 1716-26		214
30	Full-limb and knee radiography assessments of varus-valgus alignment and their relationship to osteoarthritis disease features by magnetic resonance imaging. <i>Arthritis and Rheumatism</i> , 2007 , 57, 398-406		74
29	Lower extremity ischemia, calf skeletal muscle characteristics, and functional impairment in peripheral arterial disease. <i>Journal of the American Geriatrics Society</i> , 2007 , 55, 400-6	5.6	110
28	The ratio of type II collagen breakdown to synthesis and its relationship with the progression of knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2007 , 15, 819-23	6.2	71
27	The relationship between toe-out angle during gait and progression of medial tibiofemoral osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2007 , 66, 1271-5	2.4	131
26	Epidemiology of osteoarthritis: an update. <i>Current Rheumatology Reports</i> , 2006 , 8, 7-15	4.9	109
25	The relationship between specific tissue lesions and pain severity in persons with knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2006 , 14, 1033-40	6.2	269
24	dGEMRIC as a function of BMI. <i>Osteoarthritis and Cartilage</i> , 2006 , 14, 1091-7	6.2	67
23	Femorotibial and patellar cartilage loss in patients prior to total knee arthroplasty, heterogeneity, and correlation with alignment of the knee. <i>Annals of the Rheumatic Diseases</i> , 2006 , 65, 69-73	2.4	34
22	Epidemiology of osteoarthritis: an update. <i>Current Opinion in Rheumatology</i> , 2006 , 18, 147-56	5.3	193
21	Effects of doxycycline on progression of osteoarthritis: results of a randomized, placebo-controlled, double-blind trial. <i>Arthritis and Rheumatism</i> , 2005 , 52, 2015-25		211
20	The natural history of anteroposterior laxity and its role in knee osteoarthritis progression. <i>Arthritis and Rheumatism</i> , 2005 , 52, 2343-9		38
19	Delayed gadolinium-enhanced magnetic resonance imaging of cartilage in knee osteoarthritis: findings at different radiographic stages of disease and relationship to malalignment. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3528-35		142
18	Hip abduction moment and protection against medial tibiofemoral osteoarthritis progression. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3515-9		197
17	Potential strategies to reduce medial compartment loading in patients with knee osteoarthritis of varying severity: reduced walking speed. <i>Arthritis and Rheumatism</i> , 2004 , 50, 1172-8		312
16	Varus-valgus alignment in the progression of patellofemoral osteoarthritis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 2184-90		118
15	Thrust during ambulation and the progression of knee osteoarthritis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 3897-903		163
14	The role of proprioceptive deficits, ligamentous laxity, and malalignment in development and progression of knee osteoarthritis. <i>Journal of rheumatology Supplement, The</i> , 2004 , 70, 87-92		21

13	Quadriceps strength and osteoarthritis progression in malaligned and lax knees. <i>Annals of Internal Medicine</i> , 2003 , 138, 613-9	8	223
12	Examination of exercise effects on knee osteoarthritis outcomes: why should the local mechanical environment be considered?. <i>Arthritis and Rheumatism</i> , 2003 , 49, 255-60		24
11	Physical functioning over three years in knee osteoarthritis: role of psychosocial, local mechanical, and neuromuscular factors. <i>Arthritis and Rheumatism</i> , 2003 , 48, 3359-70		358
10	Depressive symptoms and lower extremity functioning in men and women with peripheral arterial disease. <i>Journal of General Internal Medicine</i> , 2003 , 18, 461-7	4	85
9	Statin use and leg functioning in patients with and without lower-extremity peripheral arterial disease. <i>Circulation</i> , 2003 , 107, 757-61	16.7	179
8	The influence of alignment on risk of knee osteoarthritis progression according to baseline stage of disease. <i>Arthritis and Rheumatism</i> , 2002 , 46, 2632-6		220
7	Lower extremity performance is associated with daily life physical activity in individuals with and without peripheral arterial disease. <i>Journal of the American Geriatrics Society</i> , 2002 , 50, 247-55	5.6	68
6	Nonpharmacologic management of osteoarthritis. <i>Current Opinion in Rheumatology</i> , 2002 , 14, 603-7	5.3	17
5	The role of knee alignment in disease progression and functional decline in knee osteoarthritis. <i>JAMA - Journal of the American Medical Association</i> , 2001 , 286, 188-95	27.4	980
4	Local factors in osteoarthritis. <i>Current Opinion in Rheumatology</i> , 2001 , 13, 441-6	5.3	68
3	Effect of knee pain on joint loading in patients with osteoarthritis. <i>Current Opinion in Rheumatology</i> , 1999 , 11, 422-6	5.3	49
2	Impaired proprioception and osteoarthritis. <i>Current Opinion in Rheumatology</i> , 1997 , 9, 253-8	5.3	80
1	Decline of plasma growth hormone binding protein in old age. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1996 , 81, 995-997	5.6	21