Andrew R Tee

List of Publications by Citations

Source: https://exaly.com/author-pdf/5065976/andrew-r-tee-publications-by-citations.pdf

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

59 11,953 38 69 g-index

69 13,258 7.2 5.99 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
59	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016 , 12, 1-222	10.2	3838
58	Identification of the tuberous sclerosis complex-2 tumor suppressor gene product tuberin as a target of the phosphoinositide 3-kinase/akt pathway. <i>Molecular Cell</i> , 2002 , 10, 151-62	17.6	1247
57	Tuberous sclerosis complex gene products, Tuberin and Hamartin, control mTOR signaling by acting as a GTPase-activating protein complex toward Rheb. <i>Current Biology</i> , 2003 , 13, 1259-68	6.3	923
56	mTOR controls cell cycle progression through its cell growth effectors S6K1 and 4E-BP1/eukaryotic translation initiation factor 4E. <i>Molecular and Cellular Biology</i> , 2004 , 24, 200-16	4.8	680
55	Tuberous sclerosis complex-1 and -2 gene products function together to inhibit mammalian target of rapamycin (mTOR)-mediated downstream signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 13571-6	11.5	661
54	Hypoxia-inducible factor 1alpha is regulated by the mammalian target of rapamycin (mTOR) via an mTOR signaling motif. <i>Journal of Biological Chemistry</i> , 2007 , 282, 20534-43	5.4	370
53	The tuberous sclerosis protein TSC2 is not required for the regulation of the mammalian target of rapamycin by amino acids and certain cellular stresses. <i>Journal of Biological Chemistry</i> , 2005 , 280, 18717	7- 2 : 1	288
52	Activity of TSC2 is inhibited by AKT-mediated phosphorylation and membrane partitioning. <i>Journal of Cell Biology</i> , 2006 , 173, 279-89	7.3	268
51	Regulation of targets of mTOR (mammalian target of rapamycin) signalling by intracellular amino acid availability. <i>Biochemical Journal</i> , 2003 , 372, 555-66	3.8	254
50	mTOR, translational control and human disease. <i>Seminars in Cell and Developmental Biology</i> , 2005 , 16, 29-37	7.5	248
49	Mammalian target of rapamycin complex 1: signalling inputs, substrates and feedback mechanisms. <i>Cellular Signalling</i> , 2009 , 21, 827-35	4.9	204
48	A tuberous sclerosis complex signalling node at the peroxisome regulates mTORC1 and autophagy in response to ROS. <i>Nature Cell Biology</i> , 2013 , 15, 1186-96	23.4	182
47	Reactive nitrogen species regulate autophagy through ATM-AMPK-TSC2-mediated suppression of mTORC1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, E2950-7	11.5	181
46	mTOR Ser-2481 autophosphorylation monitors mTORC-specific catalytic activity and clarifies rapamycin mechanism of action. <i>Journal of Biological Chemistry</i> , 2010 , 285, 7866-79	5.4	175
45	Inactivation of the tuberous sclerosis complex-1 and -2 gene products occurs by phosphoinositide 3-kinase/Akt-dependent and -independent phosphorylation of tuberin. <i>Journal of Biological Chemistry</i> , 2003 , 278, 37288-96	5.4	170
44	Leucine and mTORC1: a complex relationship. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012 , 302, E1329-42	6	168
43	mTORC1 drives HIF-1[and VEGF-A signalling via multiple mechanisms involving 4E-BP1, S6K1 and STAT3. <i>Oncogene</i> , 2015 , 34, 2239-50	9.2	157

(2005-2011)

42	ULK1 inhibits mTORC1 signaling, promotes multisite Raptor phosphorylation and hinders substrate binding. <i>Autophagy</i> , 2011 , 7, 737-47	10.2	151
41	The extracellular signal-regulated kinase pathway regulates the phosphorylation of 4E-BP1 at multiple sites. <i>Journal of Biological Chemistry</i> , 2002 , 277, 11591-6	5.4	149
40	Caspase cleavage of initiation factor 4E-binding protein 1 yields a dominant inhibitor of cap-dependent translation and reveals a novel regulatory motif. <i>Molecular and Cellular Biology</i> , 2002 , 22, 1674-83	4.8	116
39	Structure-activity analysis of niclosamide reveals potential role for cytoplasmic pH in control of mammalian target of rapamycin complex 1 (mTORC1) signaling. <i>Journal of Biological Chemistry</i> , 2012 , 287, 17530-17545	5.4	110
38	DNA-damaging agents cause inactivation of translational regulators linked to mTOR signalling. <i>Oncogene</i> , 2000 , 19, 3021-31	9.2	108
37	Control of TSC2-Rheb signaling axis by arginine regulates mTORC1 activity. <i>ELife</i> , 2016 , 5,	8.9	102
36	The tumor suppressor folliculin regulates AMPK-dependent metabolic transformation. <i>Journal of Clinical Investigation</i> , 2014 , 124, 2640-50	15.9	101
35	The kinase triad, AMPK, mTORC1 and ULK1, maintains energy and nutrient homoeostasis. <i>Biochemical Society Transactions</i> , 2013 , 41, 939-43	5.1	92
34	Impairment of Angiogenesis by Fatty Acid Synthase Inhibition Involves mTOR Malonylation. <i>Cell Metabolism</i> , 2018 , 28, 866-880.e15	24.6	83
33	Analysis of mTOR signaling by the small G-proteins, Rheb and RhebL1. FEBS Letters, 2005, 579, 4763-8	3.8	79
32	Absence of the Birt-Hogg-Dub[gene product is associated with increased hypoxia-inducible factor transcriptional activity and a loss of metabolic flexibility. <i>Oncogene</i> , 2011 , 30, 1159-73	9.2	62
31	Mammalian target of rapamycin complex 1-mediated phosphorylation of eukaryotic initiation factor 4E-binding protein 1 requires multiple protein-protein interactions for substrate recognition. <i>Cellular Signalling</i> , 2009 , 21, 1073-84	4.9	62
30	Birt-Hogg-Dube syndrome is a novel ciliopathy. Human Molecular Genetics, 2013, 22, 4383-97	5.6	56
29	Tertiary active transport of amino acids reconstituted by coexpression of System A and L transporters in Xenopus oocytes. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2009 , 297, E822-9	6	56
28	Neurofibromatosis type 1: Fundamental insights into cell signalling and cancer. <i>Seminars in Cell and Developmental Biology</i> , 2016 , 52, 39-46	7.5	53
27	FLCN, a novel autophagy component, interacts with GABARAP and is regulated by ULK1 phosphorylation. <i>Autophagy</i> , 2014 , 10, 1749-60	10.2	48
26	The role of mTOR signalling in neurogenesis, insights from tuberous sclerosis complex. <i>Seminars in Cell and Developmental Biology</i> , 2016 , 52, 12-20	7.5	46
25	Characterization of a conserved C-terminal motif (RSPRR) in ribosomal protein S6 kinase 1 required for its mammalian target of rapamycin-dependent regulation. <i>Journal of Biological Chemistry</i> , 2005 , 280, 11101-6	5.4	46

24	cAMP inhibits mammalian target of rapamycin complex-1 and -2 (mTORC1 and 2) by promoting complex dissociation and inhibiting mTOR kinase activity. <i>Cellular Signalling</i> , 2011 , 23, 1927-35	4.9	39
23	Staurosporine inhibits phosphorylation of translational regulators linked to mTOR. <i>Cell Death and Differentiation</i> , 2001 , 8, 841-9	12.7	39
22	Bidirectional regulation of nuclear factor- B and mammalian target of rapamycin signaling functionally links Bnip3 gene repression and cell survival of ventricular myocytes. <i>Circulation: Heart Failure</i> , 2013 , 6, 335-43	7.6	38
21	The Target of Rapamycin and Mechanisms of Cell Growth. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	36
20	Oncogenic Signalling through Mechanistic Target of Rapamycin (mTOR): A Driver of Metabolic Transformation and Cancer Progression. <i>Cancers</i> , 2018 , 10,	6.6	34
19	Possible targets for nonimmunosuppressive therapy of GravesVorbitopathy. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, E1183-90	5.6	27
18	Endoplasmic reticulum stress and cell death in mTORC1-overactive cells is induced by nelfinavir and enhanced by chloroquine. <i>Molecular Oncology</i> , 2015 , 9, 675-88	7.9	25
17	Exploiting cancer vulnerabilities: mTOR, autophagy, and homeostatic imbalance. <i>Essays in Biochemistry</i> , 2017 , 61, 699-710	7.6	22
16	Characterizing the interaction of the mammalian eIF4E-related protein 4EHP with 4E-BP1. <i>FEBS Letters</i> , 2004 , 564, 58-62	3.8	21
15	STAT3 and HIF1 ignaling Drives Oncogenic Cellular Phenotypes in Malignant Peripheral Nerve Sheath Tumors. <i>Molecular Cancer Research</i> , 2015 , 13, 1149-60	6.6	19
14	Localisation and regulation of the eIF4E-binding protein 4E-BP3. FEBS Letters, 2002, 532, 319-23	3.8	17
13	Evaluation of copy number variation and gene expression in neurofibromatosis type-1-associated malignant peripheral nerve sheath tumours. <i>Human Genomics</i> , 2015 , 9, 3	6.8	15
12	Birt-Hogg-Dub[]tumour suppressor function and signalling dynamics central to folliculin. <i>Familial Cancer</i> , 2013 , 12, 367-72	3	14
11	STAT3 and mTOR: co-operating to drive HIF and angiogenesis. <i>Oncoscience</i> , 2015 , 2, 913-4	0.8	14
10	Finding a cure for tuberous sclerosis complex: From genetics through to targeted drug therapies. <i>Advances in Genetics</i> , 2019 , 103, 91-118	3.3	13
9	Targeting protein homeostasis with nelfinavir/salinomycin dual therapy effectively induces death of mTORC1 hyperactive cells. <i>Oncotarget</i> , 2017 , 8, 48711-48724	3.3	8
8	Determining the pathogenicity of patient-derived TSC2 mutations by functional characterization and clinical evidence. <i>European Journal of Human Genetics</i> , 2011 , 19, 789-95	5.3	8
7	Loss of tuberous sclerosis complex 2 sensitizes tumors to nelfinavir-bortezomib therapy to intensify endoplasmic reticulum stress-induced cell death. <i>Oncogene</i> , 2018 , 37, 5913-5925	9.2	6

LIST OF PUBLICATIONS

1	The zinc finger/RING domain protein Unkempt regulates cognitive flexibility. <i>Scientific Reports</i> , 2021 , 11, 16299	4.9	0
2	The Role of Mitochondria-Linked Fatty-Acid Uptake-Driven Adipogenesis in Graves Orbitopathy. <i>Endocrinology</i> , 2021 , 162,	4.8	1
3	Energy Stress-Mediated Cytotoxicity in Tuberous Sclerosis Complex 2-Deficient Cells with Nelfinavir and Mefloquine Treatment. <i>Cancers</i> , 2018 , 10,	6.6	2
4	Exploring transcriptional regulators Ref-1 and STAT3 as therapeutic targets in malignant peripheral nerve sheath tumours. <i>British Journal of Cancer</i> , 2021 , 124, 1566-1580	8.7	3
5	Distinctive Features of Orbital Adipose Tissue (OAT) in GravesVOrbitopathy. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	3
6	Reciprocal signaling between mTORC1 and MNK2 controls cell growth and oncogenesis. <i>Cellular and Molecular Life Sciences</i> , 2021 , 78, 249-270	10.3	5