

Tetsukazu Yahara

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/50691/publications.pdf>

Version: 2024-02-01

159
papers

6,350
citations

117625
34
h-index

76900
74
g-index

160
all docs

160
docs citations

160
times ranked

9826
citing authors

#	ARTICLE	IF	CITATIONS
1	The IPBES Conceptual Framework – connecting nature and people. <i>Current Opinion in Environmental Sustainability</i> , 2015, 14, 1-16.	6.3	1,658
2	A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny: The Legume Phylogeny Working Group (LPWG). <i>Taxon</i> , 2017, 66, 44-77.	0.7	803
3	Biodiversity, climate change, and ecosystem services. <i>Current Opinion in Environmental Sustainability</i> , 2009, 1, 46-54.	6.3	337
4	Ecosystem services: an evolutionary perspective on the links between biodiversity and human well-being. <i>Current Opinion in Environmental Sustainability</i> , 2010, 2, 66-74.	6.3	168
5	Monitoring biodiversity change through effective global coordination. <i>Current Opinion in Environmental Sustainability</i> , 2017, 29, 158-169.	6.3	147
6	Understanding and monitoring the consequences of human impacts on intraspecific variation. <i>Evolutionary Applications</i> , 2017, 10, 121-139.	3.1	145
7	Chromosomal Evolution in the Genus <i>Brachyscome</i> (Asteraceae, Astereae): Statistical Tests Regarding Correlation Between Changes in Karyotype and Habit Using Phylogenetic Information. <i>Journal of Plant Research</i> , 1999, 112, 145-161.	2.4	133
8	Building a global observing system for biodiversity. <i>Current Opinion in Environmental Sustainability</i> , 2012, 4, 139-146.	6.3	125
9	Floral scents of hawkmoth-pollinated flowers in Japan. <i>Journal of Plant Research</i> , 1998, 111, 199-205.	2.4	107
10	The earliest recorded plant virus disease. <i>Nature</i> , 2003, 422, 831-831.	27.8	101
11	Trends in site-number change of rDNA loci during polyploid evolution in <i>Sanguisorba</i> (Rosaceae). <i>Chromosoma</i> , 2002, 110, 550-558.	2.2	91
12	Effects of variation in flower number on pollinator visits in <i>Cirsium purpuratum</i> (Asteraceae). <i>American Journal of Botany</i> , 1998, 85, 219-224.	1.7	90
13	EVOLUTIONARY BIOLOGY IN BIODIVERSITY SCIENCE, CONSERVATION, AND POLICY: A CALL TO ACTION. <i>Evolution; International Journal of Organic Evolution</i> , 2010, 64, 1517-28.	2.3	87
14	Towards a global terrestrial species monitoring program. <i>Journal for Nature Conservation</i> , 2015, 25, 51-57.	1.8	86
15	Global legume diversity assessment: Concepts, key indicators, and strategies. <i>Taxon</i> , 2013, 62, 249-266.	0.7	85
16	Trade-off between light interception efficiency and light use efficiency: implications for species coexistence in one-sided light competition. <i>Journal of Ecology</i> , 2014, 102, 167-175.	4.0	82
17	Biodiversity and ecosystem services science for a sustainable planet: the DIVERSITAS vision for 2012–20. <i>Current Opinion in Environmental Sustainability</i> , 2012, 4, 101-105.	6.3	62
18	Why does the flower of <i>Lonicera japonica</i> open at dusk?. <i>Canadian Journal of Botany</i> , 1998, 76, 1806-1811.	1.1	59

#	ARTICLE	IF	CITATIONS
19	How Long to Stay on, and How Often to Visit a Flowering Plant?: A Model for Foraging Strategy When Floral Displays Vary in Size. <i>Oikos</i> , 1999, 86, 386.	2.7	53
20	The founding charter of the Genomic Observatories Network. <i>GigaScience</i> , 2014, 3, 2.	6.4	51
21	First molecular phylogeny of the pantropical genus <i>Dalbergia</i> : implications for infrageneric circumscription and biogeography. <i>South African Journal of Botany</i> , 2013, 89, 143-149.	2.5	50
22	Relative Role of Flower Color and Scent on Pollinator Attraction: Experimental Tests using F1 and F2 Hybrids of Daylily and Nightlily. <i>PLoS ONE</i> , 2012, 7, e39010.	2.5	48
23	Convergence, Consilience, and the Evolution of Temperate Deciduous Forests. <i>American Naturalist</i> , 2017, 190, S87-S104.	2.1	47
24	Why does the flower of <i>Lonicera japonica</i> open at dusk?. <i>Canadian Journal of Botany</i> , 1998, 76, 1806-1811.	1.1	46
25	Theoretical Evaluation of Pollen Transfer by Nocturnal and Diurnal Pollinators: When Should a Flower Open?. <i>Oikos</i> , 1999, 86, 233.	2.7	45
26	Chromosomal Cytology and Evolution in Eupatorieae (Asteraceae). <i>Annals of the Missouri Botanical Garden</i> , 1995, 82, 581.	1.3	41
27	Effects of virus infection and growth irradiance on fitness components and photosynthetic properties of <i>Eupatorium makinoi</i> (Compositae). <i>American Journal of Botany</i> , 1997, 84, 823-829.	1.7	41
28	Molecular phylogeny of geminivirus infecting wild plants in Japan. <i>Journal of Plant Research</i> , 1997, 110, 247-257.	2.4	41
29	Spatial and temporal variation of fish assemblages and their associations to habitat variables in a mountain stream of north Tiaoxi River, China. <i>Environmental Biology of Fishes</i> , 2012, 93, 403-417.	1.0	41
30	Effects of virus infection on demographic traits of an agamospermous population of <i>Eupatorium chinense</i> (Asteraceae). <i>Oecologia</i> , 1993, 96, 310-315.	2.0	38
31	Effects of time-dependent competition for oviposition sites on clutch sizes and offspring sex ratios in a fig wasp. <i>Oikos</i> , 2002, 96, 31-35.	2.7	37
32	Trade-Offs between Flower Number and Investment to a Flower in Selfing and Outcrossing Varieties of <i>Impatiens hypophylla</i> (Balsaminaceae). <i>American Journal of Botany</i> , 1999, 86, 1699.	1.7	36
33	Bimodal distribution of flowering time in a natural hybrid population of daylily (<i>Hemerocallis fulva</i>) and nightlily (<i>Hemerocallis citrina</i>). <i>Journal of Plant Research</i> , 2006, 119, 63-68.	2.4	36
34	Evolution of Agamospermous Races in Boehmeria and <i>Eupatorium</i> . <i>Plant Species Biology</i> , 1990, 5, 183-196.	1.0	35
35	Reproductive ecology of a cleistogamous annual, <i>Impatiens noli-tangere</i> L., occurring under different environmental conditions. <i>Ecological Research</i> , 1994, 9, 67-75.	1.5	33
36	FEMALE CONTROL OF PATERNITY DURING COPULATION: INBREEDING AVOIDANCE IN FERAL CATS. <i>Behaviour</i> , 2001, 138, 235-250.	0.8	32

#	ARTICLE	IF	CITATIONS
37	Variation of flower opening and closing times in F1 and F2 hybrids of daylily (<i>Hemerocallis</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 107 261-267.	1.7	31
38	Effects of logging and recruitment on community phylogenetic structure in 32 permanent forest plots of Kampong Thom, Cambodia. Philosophical Transactions of the Royal Society B: Biological Sciences, 2015, 370, 20140008.	4.0	31
39	Phylogeny and Phytogeography of Eupatorium (Eupatorieae, Asteraceae): Insights from Sequence Data of the nrDNA ITS Regions and cpDNA RFLP. Journal of Plant Research, 2000, 113, 79-89.	2.4	30
40	Assessing the impact of the Japanese 2005 World Exposition Project on vascular plantsâ€™ risk of extinction. Chemosphere, 2003, 53, 325-336.	8.2	28
41	Changes in Defense of an Alien Plant Ambrosia artemisiifolia before and after the Invasion of a Native Specialist Enemy Ophraella communis. PLoS ONE, 2012, 7, e49114.	2.5	28
42	Climatic Gradients of Arms Race Coevolution. American Naturalist, 2011, 177, 562-573.	2.1	27
43	GRAPHICAL ANALYSIS OF MATING SYSTEM EVOLUTION IN PLANTS. Evolution; International Journal of Organic Evolution, 1992, 46, 557-561.	2.3	26
44	Extinction risk assessment of declining wild populations: The case of the southern Bluefin Tuna. Researches on Population Ecology, 1998, 40, 271-278.	0.9	26
45	Facultative sex ratio adjustment in response to male tarsus length in the Varied Tit Parus varius. Ibis, 2003, 146, 108-113.	1.9	25
46	Pollinator-Mediated Selection on Flower Color, Flower Scent and Flower Morphology of Hemerocallis: Evidence from Genotyping Individual Pollen Grains On the Stigma. PLoS ONE, 2013, 8, e85601.	2.5	25
47	Range shift and introgression of the rear and leading populations in two ecologically distinct Rubus species. BMC Evolutionary Biology, 2014, 14, 209.	3.2	25
48	Low genetic differentiation among populations of <i>Arabis serrata</i> (Brassicaceae) along an altitudinal gradient. Journal of Plant Research, 1993, 106, 143-148.	2.4	23
49	Is tuna critically endangered? Extinction risk of a large and overexploited population. Ecological Research, 1997, 12, 345-356.	1.5	23
50	Pollination biology of <i>Lilium japonicum</i> var. <i>abeanicum</i> and var. <i>japonicum</i>: evidence of adaptation to the different availability of diurnal and nocturnal pollinators. Plant Species Biology, 2012, 27, 96-105.	1.0	22
51	Molecular Phylogeny of Eupatorieae (Asteraceae) Estimated from cpDNA RFLP and its Implication for the Polyploid Origin Hypothesis of the Tribe. Journal of Plant Research, 2000, 113, 91-96.	2.4	20
52	More highly female-biased sex ratio in the fig wasp, <i>Blastophaga nipponica</i> Grandi (Agaonidae). Researches on Population Ecology, 1998, 40, 239-242.	0.9	19
53	EXTENSIVE GENE DUPLICATIONS IN DIPLOID EUPATORIUM (ASTERACEAE). American Journal of Botany, 1989, 76, 1247-1253.	1.7	18
54	Mexican species of the genus Stevia (Eupatorieae, Asteraceae): Chromosome numbers and geographical distribution. Plant Species Biology, 2001, 16, 49-68.	1.0	18

#	ARTICLE	IF	CITATIONS
55	Extra-pair mate choice in the female great tit <i>Parus major</i> : good males or compatible males. <i>Journal of Ethology</i> , 2009, 27, 349-359.	0.8	18
56	A taxonomic study of <i>Quercus langbianensis</i> complex based on morphology and DNA barcodes of classic and next generation sequences. <i>PhytoKeys</i> , 2018, 95, 37-70.	1.0	18
57	Effects of virus infection and light environment on population dynamics of <i>Eupatorium makinoi</i> (Asteraceae). <i>American Journal of Botany</i> , 2001, 88, 616-622.	1.7	17
58	Post-pollination reproductive isolation between diurnally and nocturnally flowering daylilies, <i>Hemerocallis fulva</i> and <i>Hemerocallis citrina</i> . <i>Journal of Plant Research</i> , 2006, 119, 617-623.	2.4	17
59	Mitochondrial phylogeny certified PGL (Paternal Genome Loss) is of single origin and haplodiploidy sensu stricto (arrhenotoky) did not evolve from PGL in the scale insects (Hemiptera: Coccoidea). <i>Genes and Genetic Systems</i> , 2009, 84, 57-66.	0.7	17
60	Crisis of Japanese Vascular Flora Shown By Quantifying Extinction Risks for 1618 Taxa. <i>PLoS ONE</i> , 2014, 9, e98954.	2.5	17
61	Genetic diversity assessments in the century of genome science. <i>Current Opinion in Environmental Sustainability</i> , 2010, 2, 43-49.	6.3	16
62	New perspectives on habitat selection by the Black-faced Spoonbill <i>Platalea minor</i> based upon satellite telemetry. <i>Bird Conservation International</i> , 2013, 23, 495-501.	1.3	16
63	Size advantage for male function and size-dependent sex allocation in <i>< i>Ambrosia artemisiifolia</i></i> , a wind-pollinated plant. <i>Ecology and Evolution</i> , 2018, 8, 1159-1170.	1.9	16
64	Comparative phylogeography of two closely related <i>Viola</i> species occurring in contrasting habitats in the Japanese archipelago. <i>Journal of Plant Research</i> , 2009, 122, 389-401.	2.4	15
65	Distribution of Sexual and Agamospermous Populations of <i>Eupatorium</i> (Compositae) in Asia. <i>Plant Species Biology</i> , 1989, 4, 37-46.	1.0	14
66	Ant-Attendance in Extrafloral Nectar-Bearing Plants Promotes Growth and Decreases the Expression of Traits Related to Direct Defenses. <i>Evolutionary Biology</i> , 2015, 42, 191-198.	1.1	14
67	UV bullseye contrast of <i>< i>Hemerocallis</i></i> flowers attracts hawkmoths but not swallowtail butterflies. <i>Ecology and Evolution</i> , 2019, 9, 52-64.	1.9	14
68	Evolutionary history of <i>Hemerocallis</i> in Japan inferred from chloroplast and nuclear phylogenies and levels of interspecific gene flow. <i>Molecular Phylogenetics and Evolution</i> , 2021, 164, 107264.	2.7	14
69	Spatial structure of genetic variation in a population of the endangered plant <i>Cerastium fischerianum</i> var. <i>molle</i> (Caryophyllaceae).. <i>Genes and Genetic Systems</i> , 1997, 72, 239-242.	0.7	13
70	Evolution of floral dimorphism in a cleistogamous annual, <i>Impatiens noli-tangere</i> L. occurring under different environmental conditions. <i>Ecological Research</i> , 2004, 19, 571-580.	1.5	13
71	CHROMOSOME NUMBERS AND KARYOTYPES IN ASTERACEAE ¹ . <i>Annals of the Missouri Botanical Garden</i> , 2007, 94, 643-654.	1.3	13
72	Genetic structure and putative selective sweep in the pioneer tree, <i>Zanthoxylum ailanthoides</i> . <i>Journal of Plant Research</i> , 2010, 123, 607-616.	2.4	13

#	ARTICLE	IF	CITATIONS
73	Morphology and genome size of the widespread weed <i>Cardamine occulta</i> : how it differs from cleistogamic <i>C. kokaiensis</i> and other closely related taxa in Europe and Asia. <i>Botanical Journal of the Linnean Society</i> , 2018, 187, 456-482.	1.6	13
74	Molecular evolution of a host-range gene in geminiviruses infecting asexual populations of <i>Eupatorium makinoi</i> . <i>Genes and Genetic Systems</i> , 1998, 73, 137-141.	0.7	12
75	Directional selection for early flowering is imposed by a re-associated herbivore - but no evidence of directional evolution. <i>Basic and Applied Ecology</i> , 2013, 14, 387-395.	2.7	12
76	Distribution pattern, threats and conservation of fish biodiversity in the East Tiaoxi, China. <i>Environmental Biology of Fishes</i> , 2013, 96, 519-533.	1.0	12
77	Formation of a hybrid triploid agamosperm on a sexual diploid plant: evidence from progeny tests in <i>Taraxacum platycarpum</i> Dahlst.. <i>Plant Systematics and Evolution</i> , 2014, 300, 863-870.	0.9	12
78	Phylogeny and biogeography of the genus <i>Stevia</i> (Asteraceae: Eupatorieae): an example of diversification in the Asteraceae in the new world. <i>Journal of Plant Research</i> , 2017, 130, 953-972.	2.4	12
79	Quantitative comparison of flowering phenology traits among trees, perennial herbs, and annuals in a temperate plant community. <i>American Journal of Botany</i> , 2019, 106, 1545-1557.	1.7	12
80	Museomics for reconstructing historical floristic exchanges: Divergence of stone oaks across Wallacea. <i>PLoS ONE</i> , 2020, 15, e0232936.	2.5	12
81	Natural Hybrid Populations between Chasmogamous and Cleistogamous Species, <i>Ainsliaea faurieana</i> and <i>A. apiculata</i> (Asteraceae; Mutisiae): Morphology, Cytology, Reproductive Mode and Allozyme Variation. <i>Plant Species Biology</i> , 1992, 7, 49-59.	1.0	11
82	Thirteen new species and two new combinations of <i>Stevia</i> (Asteraceae: Eupatorieae) from Mexico. <i>Brittonia</i> , 2001, 53, 377-395.	0.2	11
83	Distribution and variation of sexual and agamospermous populations of <i>Stevia</i> (Asteraceae:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 TF		
84	<i>Thismia brunneomitroides</i> (Thismiaceae), a new mycoheterotrophic species from southern Thailand. <i>Phytotaxa</i> , 2017, 314, 103.	0.3	11
85	The <sc>Asia-Pacific</sc> Biodiversity Observation Network: 10-year achievements and new strategies to 2030. <i>Ecological Research</i> , 2021, 36, 232-257.	1.5	11
86	Dispersal of chasmogamous and cleistogamous seeds in <i>Viola hondoensis</i> W. Backer et H. Boiss. <i>Botanical Magazine</i> , 1992, 105, 323-326.	0.6	10
87	Temporally changing male reproductive success and resource allocation strategy in protandrous <i>Heracleum lanatum</i> (Apiaceae). <i>Journal of Plant Research</i> , 1997, 110, 227-234.	2.4	10
88	Effects of Salinity and Temperature on Seed Germination in a Japanese Endangered Halophyte <i>Triglochin maritimum</i> (Juncaginaceae). <i>Journal of Plant Research</i> , 1999, 112, 457-461.	2.4	10
89	Reproductive isolation on interspecific backcross of F1 pollen to parental species, <i>Hemerocallis fulva</i> and <i>H. citrina</i> (Hemerocallidaceae). <i>Journal of Plant Research</i> , 2008, 121, 287-291.	2.4	10
90	Genetic variation and population structure of a threatened timber tree <i>Dalbergia cochinchinensis</i> in Cambodia. <i>Tree Genetics and Genomes</i> , 2017, 13, 1.	1.6	10

#	ARTICLE	IF	CITATIONS
91	A new species and two new records of <i>Quercus</i> (Fagaceae) from northern Vietnam. <i>PhytoKeys</i> , 2018, 92, 1-15.	1.0	10
92	Self-pollination of <i>Asarum caulescens</i> Maxim. (Aristolochiaceae) in Japan. <i>Plant Species Biology</i> , 1987, 2, 133-136.	1.0	9
93	Difference in flowering time as an isolating barrier. <i>Journal of Theoretical Biology</i> , 2013, 317, 161-167.	1.7	9
94	Biodiversity only makes sense in the light of evolution. <i>Journal of Biosciences</i> , 2014, 39, 333-337.	1.1	9
95	Flora of Bokor National Park VII: <i>Thismia bokorensis</i> (Burmanniaceae), a new species representing a new generic record. <i>Phytotaxa</i> , 2018, 334, 65.	0.3	9
96	Flora of Nam Kading National Protected Area I: a new species of yellow-flowered <i>Strobilanthes</i> (Acanthaceae), <i>S. namkadingensis</i> . <i>PhytoKeys</i> , 2017, 81, 11-17.	1.0	9
97	Studies on the asian eupatoria. <i>Botanical Magazine</i> , 1984, 97, 87-105.	0.6	8
98	VERY LOW GENETIC HETEROZYGOSITIES IN SEXUAL AND AGAMOSPERMOUS POPULATIONS OF <i>EUPATORIUM ALTISSIMUM</i> (ASTERACEAE). <i>American Journal of Botany</i> , 1991, 78, 706-710.	1.7	8
99	Difference in flowering time can initiate speciation of nocturnally flowering species. <i>Journal of Theoretical Biology</i> , 2015, 370, 61-71.	1.7	8
100	Strategies to Observe and Assess Changes of Terrestrial Biodiversity in the Asia-Pacific Regions. <i>Structure and Function of Mountain Ecosystems in Japan</i> , 2012, , 3-19.	0.5	8
101	Peroxidase Phenotypes of <i>Eupatorium mohrii</i> and <i>E. scabridum</i> (Compositae), Widespread All-agamospermous "Species" in the Southeastern United States. <i>Plant Species Biology</i> , 1986, 1, 27-34.	1.0	7
102	Intron length variation of the <i>Adh</i> gene in <i>Brachyscome</i> (Asteraceae). <i>Plant Molecular Biology</i> , 1995, 28, 1067-1073.	3.9	7
103	Pollinator trapping in selfing carnivorous plants, <i>< i>Drosera makinoi</i></i> and <i>< i>D. toyoakensis</i></i> (Droseraceae). <i>Ecological Research</i> , 2018, 33, 487-494.	1.5	7
104	Extensive Gene Duplications in Diploid <i>Eupatorium</i> (Asteraceae). <i>American Journal of Botany</i> , 1989, 76, 1247.	1.7	7
105	<i>Cryptocarya kaengkrachanensis</i> , a new species of Lauraceae from Kaeng Krachan National Park, southwest Thailand. <i>PhytoKeys</i> , 2020, 140, 139-157.	1.0	7
106	<i>Popowia bachmaensis</i> (Annonaceae), a new species from Bach Ma National Park, Central Vietnam. <i>PhytoKeys</i> , 2016, 65, 125-131.	1.0	7
107	Factors causing variation in flock size: Decision making to join a foraging flock. <i>Ecological Research</i> , 2002, 17, 361-371.	1.5	6
108	Soil disturbances can suppress the invasion of alien plants under plantâ€“soil feedback. <i>Ecological Modelling</i> , 2013, 260, 42-49.	2.5	6

#	ARTICLE	IF	CITATIONS
109	Lithocarpus dahuoaiensis (Fagaceae), a new species from Lam Dong Province, Vietnam. <i>PhytoKeys</i> , 2016, 69, 23-30.	1.0	6
110	A sensitive flower: mechanical stimulation induces rapid flower closure in <i>Drosera</i> spp. (Droseraceae). <i>Plant Species Biology</i> , 2018, 33, 153-157.	1.0	6
111	Flora of Nam Kading National Protected Area III: <i>Begonia namkadingensis</i> (Begoniaceae), a new species in limestone area. <i>Phytotaxa</i> , 2018, 334, 195.	0.3	6
112	The effects of water pollution on the phylogenetic community structure of aquatic plants in the East Tiaoxi River, China. <i>Freshwater Biology</i> , 2020, 65, 632-645.	2.4	6
113	Validation of <i>Hosta alata</i> (Asparagaceae) as a new species and its phylogenetic affinity. <i>PhytoKeys</i> , 2021, 181, 79-93.	1.0	6
114	Molecular phylogeny and taxonomy of the <i>Hydrangea serrata</i> complex (Hydrangeaceae) in western Japan, including a new subspecies of <i>H. acuminata</i> from Yakushima. <i>PhytoKeys</i> , 2022, 188, 49-71.	1.0	6
115	A new subspecies of <i>Stellaria alsine</i> (Caryophyllaceae) from Yakushima, Japan. <i>PhytoKeys</i> , 2021, 187, 177-188.	1.0	6
116	Morphological and molecular evidence reveals three new species of <i>Lithocarpus</i> (Fagaceae) from Bidoup-Nui Ba National Park, Vietnam. <i>PhytoKeys</i> , 2021, 186, 73-92.	1.0	6
117	Distinction in morphology and esterase isozyme between <i>Eupatorium glehni</i> (= <i>E. chinense</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 165-179.	0.6	5
118	Analysis on Pollen Flow in a Hybridizing Population between <i>Farfugium hiberniflorum</i> and <i>F. japonicum</i> (Asteraceae: Senecioneae): Derivation of an Empirical Formula Describing Frequency of Hybridization as a Function of Interspecific Plant Distance. <i>Plant Species Biology</i> , 1989, 4, 101-106.	1.0	5
119	Isolation of polymorphic microsatellite loci in <i>Hemerocallis fulva</i> and <i>Hemerocallis citrina</i> (Hemerocallidaceae). <i>Molecular Ecology Notes</i> , 2006, 6, 909-911.	1.7	5
120	<i>Goniothalamus flagellistylus</i> Tagane & V. S. Dang (Annonaceae), a new species from Mt. Hon Ba, Vietnam. <i>PhytoKeys</i> , 2015, 50, 1-8.	1.0	5
121	Constant tree species richness along an elevational gradient of Mt. Bokor, a table-shaped mountain in southwestern Cambodia. <i>Ecological Research</i> , 2016, 31, 495-504.	1.5	5
122	<i>Lasianthus honbaensis</i> (Rubiaceae), a New Species from Southern Vietnam. <i>Annales Botanici Fennici</i> , 2016, 53, 263-266.	0.1	5
123	Hoverflies can sense the risk of being trapped by carnivorous plants: An empirical study using <i>Sphaerophoria menthastris</i> and <i>Drosera toyoakensis</i> . <i>Journal of Asia-Pacific Entomology</i> , 2018, 21, 944-946.	0.9	5
124	<i>Macrosolen bidouensis</i> (Loranthaceae), a new species from Bidoup Nui Ba National Park, southern Vietnam. <i>PhytoKeys</i> , 2017, 80, 113-120.	1.0	5
125	Effects of flower production and predispersal seed predation on reproduction in <i>Cirsium purpuratum</i> . <i>Canadian Journal of Botany</i> , 2000, 78, 230-236.	1.1	4
126	Flora of Nam Kading National Protected Area IV: Two new species of Annonaceae, <i>Monoon namkadingense</i> and <i>Neo-uvaria laosensis</i> . <i>Phytotaxa</i> , 2018, 336, 82.	0.3	4

#	ARTICLE	IF	CITATIONS
127	Developing a Regional Network of Biodiversity Observation in the Asia-Pacific Region: Achievements and Challenges of AP BON. Structure and Function of Mountain Ecosystems in Japan, 2014, , 3-28.	0.5	4
128	Very Low Genetic Heterozygosities in Sexual and Agamospermous Populations of <i>Eupatorium altissimum</i> (Asteraceae). American Journal of Botany, 1991, 78, 706.	1.7	4
129	<i>Callicarpa bachmaensis</i> Soejima & Tagane (Lamiaceae), a new species from Bach Ma National Park in Thua Thien Hue Province, Central Vietnam. PhytoKeys, 2016, 62, 33-39.	1.0	4
130	Effects of biparental inbreeding on the evolution of gynodioecy: A model and a case study in <i>Chionographis japonica</i> var. <i>kurohimensis</i> . Journal of Plant Research, 1993, 106, 279-281.	2.4	3
131	Amplified fragment length polymorphism analysis of the genetic variation of an endangered plant <i>Lysimachia tashiroi</i> (Myrsinaceae). Plant Species Biology, 2007, 22, 33-39.	1.0	3
132	A genome-wide AFLP replacement in a hybrid population derived from two closely related <i>Viola</i> species from contrasting habitats. Plant Systematics and Evolution, 2015, 301, 1073-1084.	0.9	3
133	<i>Lysimachia kraduengensis</i> (Primulaceae), a new species from northeastern Thailand. Phytotaxa, 2016, 289, 69.	0.3	3
134	Effects of density, season, and food intake on sika deer nutrition on Yakushima Island, Japan. Ecological Research, 2017, 32, 369-378.	1.5	3
135	<i>Mussaenda recurvata</i> (Rubiaceae), a new species from southern Vietnam with observations on its heterostyly. Phytotaxa, 2017, 328, 167.	0.3	3
136	Effects of apical damage on plant growth and male and female reproductive investments in <i>Ambrosia artemisiifolia</i> , a wind-pollinated plant. Plant Ecology, 2018, 219, 853-862.	1.6	3
137	<i>Quercus ngochoaensis</i> (Fagaceae), a new species from Ba Vi National Park, northern, Vietnam. Phytotaxa, 2021, 516, .	0.3	3
138	Phylogeny and phytogeography in the genus <i>Eupatorium</i> (Asteraceae). Korean Journal of Plant Taxonomy, 1998, 28, 83-98.	0.7	3
139	<i>Garcinia hopii</i> (Clusiaceae), a new species from Bidoup Nui Ba National Park, southern Vietnam. PhytoKeys, 2017, 77, 63-70.	1.0	3
140	<i>Homalium glandulosum</i> (Salicaceae), a new species from Vu Quang National Park, North Central Vietnam. PhytoKeys, 2016, 58, 97-104.	1.0	3
141	<i>Lithocarpus vuquangensis</i> (Fagaceae), a new species from Vu Quang National Park, Vietnam. PhytoKeys, 2018, 95, 15-25.	1.0	3
142	Geographical and seasonal variation of plant taxa detected in faces of <i>Cervus nippon yakushimae</i> based on plant DNA analysis in Yakushima Island. Ecological Research, 2022, 37, 582-597.	1.5	3
143	Recurved <i>Taraxacum</i> phyllaries function as a floral defense: experimental evidence and its implication for <i>Taraxacum</i> evolutionary history. Ecological Research, 2017, 32, 313-329.	1.5	2
144	<i>Lecanorchis sarawakensis</i> (Orchidaceae, Vanilloideae), a new mycoheterotrophic species from Sarawak, Borneo. Phytotaxa, 2018, 338, 135.	0.3	2

#	ARTICLE	IF	CITATIONS
145	Erythroxylum calypratum (Erythroxylaceae), a new species from Mt. Fansipan, northern Vietnam. Phytotaxa, 2018, 347, 279.	0.3	2
146	Decision Science for Future Earth: A Conceptual Framework. , 2021, , 3-64.		2
147	Aporosa tetragona Tagane & V. S. Dang (Phyllanthaceae), a new species from Mt. Hon Ba, Vietnam. PhytoKeys, 2015, 57, 51-60.	1.0	2
148	Glycosmis suberosa (Rutaceae), a new species from Khao Luang National Park, Peninsular Thailand. Thai Forest Bulletin (Botany), 2016, 44, 116-121.	0.2	2
149	Flora of Bokor National Park V: Two new species of Machilus (Lauraceae), M. bokorensis and M. brevipaniculata. PhytoKeys, 2016, 65, 35-46.	1.0	2
150	Asplenium minutifolium (Aspleniaceae), a new species from Thailand. Thai Forest Bulletin (Botany), 2017, 45, 29-34.	0.2	2
151	Adaptive Species Differentiation and Population Uniformity in Viola Species Sharing Similar Geographical Distribution but Differing Habitat Preferences. Journal of Plant Studies, 2012, 1, .	0.3	1
152	Flora of Bokor National Park, Cambodia VI: A new species of Wikstroemia (Thymelaeaceae), W. bokorensis. Phytotaxa, 2017, 317, 280.	0.3	1
153	Five new species of Syzygium (Myrtaceae) from Indochina and Thailand. Phytotaxa, 2018, 375, 247.	0.3	1
154	Evaluating the genetic diversity in two tropical leguminous trees, <i><i>Dalbergia cochinchinensis</i></i> and <i><i>D. nigrescens</i></i> , in lowland forests in Cambodia and Thailand using MiG-seq. Genes and Genetic Systems, 2021, 96, 41-53.	0.7	1
155	A new species of Eustigma (Hamamelidaceae) from Hon Ba Nature Reserve, Vietnam. PhytoKeys, 2016, 65, 47-55.	1.0	1
156	<p>Flora of Nam Kading National Protected Area VII: a new species of Diospyros</p> (Ebenaceae), D. laoensis</p>. Phytotaxa, 2020, 477, 90-96.	0.3	1
157	A new species of Gynochthodes (Rubiaceae), G. honbaensis from Hon Ba Nature Reserve, southern Vietnam. Phytotaxa, 2019, 406, 213-217.	0.3	0
158	<p>Resurrection and emended description ofSciaphila major(Triuridaceae)</p>. Phytotaxa, 2020, 459, 25-38.	0.3	0
159	Gentiana laotica, a new species of Gentianaceae from Laos. Thai Forest Bulletin (Botany), 2018, 46, 72-75.	0.2	0