

Hunjoo Ha

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5077191/hunjoo-ha-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

131
papers

5,258
citations

37
h-index

70
g-index

133
ext. papers

5,793
ext. citations

5.1
avg. IF

5.62
L-index

#	Paper	IF	Citations
131	Kidney protective potential of lactoferrin: pharmacological insights and therapeutic advances.. <i>Korean Journal of Physiology and Pharmacology</i> , 2022 , 26, 1-13	1.8	1
130	Exercise-Induced Irisin Decreases Inflammation and Improves NAFLD by Competitive Binding with MD2.. <i>Cells</i> , 2021 , 10,	7.9	4
129	Heukcha, naturally post-fermented green tea extract, ameliorates diet-induced hypercholesterolemia and NAFLD in hamster. <i>Journal of Food Science</i> , 2021 , 86, 5016-5025	3.4	2
128	Prospects of Marine Sterols against Pathobiology of Alzheimer's Disease: Pharmacological Insights and Technological Advances. <i>Marine Drugs</i> , 2021 , 19,	6	3
127	Dual Actions of A and A Adenosine Receptor Ligand Prevents Obstruction-Induced Kidney Fibrosis in Mice. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	2
126	Prospective Pharmacological Potential of Resveratrol in Delaying Kidney Aging. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	7
125	CO-Releasing Molecule-2 Prevents Acute Kidney Injury through Suppression of ROS-Fyn-ER Stress Signaling in Mouse Model. <i>Oxidative Medicine and Cellular Longevity</i> , 2021 , 2021, 9947772	6.7	4
124	Activation of Adrenergic receptor signaling modulates inflammation: a target limiting the progression of kidney diseases. <i>Archives of Pharmacal Research</i> , 2021 , 44, 49-62	6.1	9
123	Dojuksan ameliorates tubulointerstitial fibrosis through irisin-mediated muscle-kidney crosstalk. <i>Phytomedicine</i> , 2021 , 80, 153393	6.5	5
122	KF-1607, a Novel Pan Src Kinase Inhibitor, Attenuates Obstruction-Induced Tubulointerstitial Fibrosis in Mice. <i>Biomolecules and Therapeutics</i> , 2021 , 29, 41-51	4.2	2
121	Associations of Circulating Irisin with FNDC5 Expression in Fat and Muscle in Type 1 and Type 2 Diabetic Mice. <i>Biomolecules</i> , 2021 , 11,	5.9	6
120	Pharmacotherapy against Oxidative Stress in Chronic Kidney Disease: Promising Small Molecule Natural Products Targeting Nrf2-HO-1 Signaling. <i>Antioxidants</i> , 2021 , 10,	7.1	19
119	Protective Effects of Black Cumin () and Its Bioactive Constituent, Thymoquinone against Kidney Injury: An Aspect on Pharmacological Insights. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
118	Urinary Metabolomic Profiling in Streptozotocin-Induced Diabetic Mice after Treatment with Losartan. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	3
117	Inhibition of Src Family Kinases Ameliorates LPS-Induced Acute Kidney Injury and Mitochondrial Dysfunction in Mice. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	5
116	Evaluation of Anti-Tumor Effects of Whole-Body Low-Dose Irradiation in Metastatic Mouse Models. <i>Cancers</i> , 2020 , 12,	6.6	2
115	Recent Insights Into SREBP as a Direct Mediator of Kidney Fibrosis via Lipid-Independent Pathways. <i>Frontiers in Pharmacology</i> , 2020 , 11, 265	5.6	23

114	Fyn Kinase: A Potential Therapeutic Target in Acute Kidney Injury. <i>Biomolecules and Therapeutics</i> , 2020 , 28, 213-221	4.2	7
113	The impaired redox balance in peroxisomes of catalase knockout mice accelerates nonalcoholic fatty liver disease through endoplasmic reticulum stress. <i>Free Radical Biology and Medicine</i> , 2020 , 148, 22-32	7.8	20
112	P0719SRC KINASES AGGRAVATE DIABETIC KIDNEY INJURY THROUGH ACTIVATION OF ENDOPLASMIC RETICULUM STRESS. <i>Nephrology Dialysis Transplantation</i> , 2020 , 35,	4.3	1
111	Autophagy attenuates tubulointerstitial fibrosis through regulating transforming growth factor- β and NLRP3 inflammasome signaling pathway. <i>Cell Death and Disease</i> , 2019 , 10, 78	9.8	42
110	Inhibition of Karyopherin- β Augments Radiation-Induced Cell Death by Perturbing BRCA1-Mediated DNA Repair. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	7
109	PGC-1 α potential therapeutic target against kidney aging. <i>Aging Cell</i> , 2019 , 18, e12994	9.9	22
108	Impaired Peroxisomal Fitness in Obese Mice, a Vicious Cycle Exacerbating Adipocyte Dysfunction Oxidative Stress. <i>Antioxidants and Redox Signaling</i> , 2019 , 31, 1339-1351	8.4	6
107	Urinary myo-inositol is associated with the clinical outcome in focal segmental glomerulosclerosis. <i>Scientific Reports</i> , 2019 , 9, 14707	4.9	1
106	Integrative Omics Reveals Metabolic and Transcriptomic Alteration of Nonalcoholic Fatty Liver Disease in Catalase Knockout Mice. <i>Biomolecules and Therapeutics</i> , 2019 , 27, 134-144	4.2	6
105	Enrichment of Short-Chain Ceramides and Free Fatty Acids in the Skin Epidermis, Liver, and Kidneys of db/db Mice, a Type 2 Diabetes Mellitus Model. <i>Biomolecules and Therapeutics</i> , 2019 , 27, 457-465	4.2	5
104	Network-based integrated analysis of omics data reveal novel players of TGF- β -induced EMT in human peritoneal mesothelial cells. <i>Scientific Reports</i> , 2019 , 9, 1497	4.9	7
103	Cigarette smoke inhalation aggravates diabetic kidney injury in rats. <i>Toxicology Research</i> , 2019 , 8, 964-976	4.3	2
102	Ablation of catalase promotes non-alcoholic fatty liver via oxidative stress and mitochondrial dysfunction in diet-induced obese mice. <i>Pflugers Archiv European Journal of Physiology</i> , 2019 , 471, 829-843	4.6	13
101	Peroxiredoxin 3 deficiency accelerates chronic kidney injury in mice through interactions between macrophages and tubular epithelial cells. <i>Free Radical Biology and Medicine</i> , 2019 , 131, 162-172	7.8	14
100	Correlation study between A adenosine receptor binding affinity and anti-renal interstitial fibrosis activity of truncated adenosine derivatives. <i>Archives of Pharmacal Research</i> , 2019 , 42, 773-779	6.1	3
99	Orally active, species-independent novel A adenosine receptor antagonist protects against kidney injury in db/db mice. <i>Experimental and Molecular Medicine</i> , 2018 , 50, 1-14	12.8	11
98	Metabolic changes in urine and serum during progression of diabetic kidney disease in a mouse model. <i>Archives of Biochemistry and Biophysics</i> , 2018 , 646, 90-97	4.1	5
97	A pan-NADPH Oxidase Inhibitor Ameliorates Kidney Injury in Type 1 Diabetic Rats. <i>Pharmacology</i> , 2018 , 102, 180-189	2.3	13

96	SJB-003-085, a newly-synthesized Src kinase inhibitor, attenuates the progression of renal interstitial fibrosis. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2018</i> , WCP2018, PO1-3-23	0	
95	Carbon monoxide reduces ER stress through suppression of Fyn in acute kidney injury. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2018</i> , WCP2018, PO1-3-6	0	
94	The importance of peroxisome in obesity-related adipocyte injury. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2018</i> , WCP2018, PO3-6-17	0	
93	Carbon monoxide releasing molecule-2 protects mice against acute kidney injury through inhibition of ER stress. <i>Korean Journal of Physiology and Pharmacology, 2018</i> , 22, 567-575	1.8	16
92	Delayed treatment with fenofibrate protects against high-fat diet-induced kidney injury in mice: the possible role of AMPK autophagy. <i>American Journal of Physiology - Renal Physiology, 2017</i> , 312, F323-F334	4.3	34
91	Endogenous catalase delays high-fat diet-induced liver injury in mice. <i>Korean Journal of Physiology and Pharmacology, 2017</i> , 21, 317-325	1.8	21
90	A novel pan-Nox inhibitor, APX-115, protects kidney injury in streptozotocin-induced diabetic mice: possible role of peroxisomal and mitochondrial biogenesis. <i>Oncotarget, 2017</i> , 8, 74217-74232	3.3	23
89	Effects of low-dose irradiation on mice with Escherichia coli-induced sepsis. <i>Toxicology and Applied Pharmacology, 2017</i> , 333, 17-25	4.6	11
88	8-Hydroxy-2-deoxyguanosine ameliorates high-fat diet-induced insulin resistance and adipocyte dysfunction in mice. <i>Biochemical and Biophysical Research Communications, 2017</i> , 491, 890-896	3.4	12
87	TM5441, a plasminogen activator inhibitor-1 inhibitor, protects against high fat diet-induced non-alcoholic fatty liver disease. <i>Oncotarget, 2017</i> , 8, 89746-89760	3.3	12
86	Novel Plasminogen Activator Inhibitor-1 Inhibitors Prevent Diabetic Kidney Injury in a Mouse Model. <i>PLoS ONE, 2016</i> , 11, e0157012	3.7	28
85	Novel Role of Endogenous Catalase in Macrophage Polarization in Adipose Tissue. <i>Mediators of Inflammation, 2016</i> , 2016, 8675905	4.3	14
84	A novel plasminogen activator inhibitor-1 inhibitor, TM5441, protects against high-fat diet-induced obesity and adipocyte injury in mice. <i>British Journal of Pharmacology, 2016</i> , 173, 2622-32	8.6	22
83	Metformin Radiosensitizes p53-Deficient Colorectal Cancer Cells through Induction of G2/M Arrest and Inhibition of DNA Repair Proteins. <i>PLoS ONE, 2015</i> , 10, e0143596	3.7	33
82	Short-term Treatment of Daumone Improves Hepatic Inflammation in Aged Mice. <i>Korean Journal of Physiology and Pharmacology, 2015</i> , 19, 269-74	1.8	8
81	18β-glycyrrhetic acid attenuates anandamide-induced adiposity and high-fat diet induced obesity. <i>Molecular Nutrition and Food Research, 2014</i> , 58, 1436-46	5.9	18
80	Effect of biocompatible peritoneal dialysis solution on residual renal function: a systematic review of randomized controlled trials. <i>Peritoneal Dialysis International, 2014</i> , 34, 724-31	2.8	27
79	Clinical implication of allogeneic implantation of adipogenic differentiated adipose-derived stem cells. <i>Stem Cells Translational Medicine, 2014</i> , 3, 1312-21	6.9	27

78	Synthesis and anti-renal fibrosis activity of conformationally locked truncated 2-hexynyl-N(6)-substituted-(N)-methanocarba-nucleosides as A3 adenosine receptor antagonists and partial agonists. <i>Journal of Medicinal Chemistry</i> , 2014 , 57, 1344-54	8.3	19
77	Inhibitory role of the KEAP1-NRF2 pathway in TGF β -stimulated renal epithelial transition to fibroblastic cells: a modulatory effect on SMAD signaling. <i>PLoS ONE</i> , 2014 , 9, e93265	3.7	49
76	Daumone fed late in life improves survival and reduces hepatic inflammation and fibrosis in mice. <i>Aging Cell</i> , 2014 , 13, 709-18	9.9	14
75	Real-time monitoring of adipocyte differentiation using a capacitance sensor array. <i>Lab on A Chip</i> , 2013 , 13, 3410-6	7.2	10
74	Autologous adipose tissue-derived stem cells treatment demonstrated favorable and sustainable therapeutic effect for Crohn's fistula. <i>Stem Cells</i> , 2013 , 31, 2575-81	5.8	190
73	Fractalkine and its receptor mediate extracellular matrix accumulation in diabetic nephropathy in mice. <i>Diabetologia</i> , 2013 , 56, 1661-9	10.3	33
72	Where are we now in diabetic research?. <i>Archives of Pharmacal Research</i> , 2013 , 36, 142-4	6.1	2
71	Carnosic acid, a phenolic diterpene from rosemary, prevents UV-induced expression of matrix metalloproteinases in human skin fibroblasts and keratinocytes. <i>Experimental Dermatology</i> , 2013 , 22, 336-41	4	59
70	The selective A3AR antagonist LJ-1888 ameliorates UUO-induced tubulointerstitial fibrosis. <i>American Journal of Pathology</i> , 2013 , 183, 1488-1497	5.8	34
69	Fibrin glue improves the therapeutic effect of MSCs by sustaining survival and paracrine function. <i>Tissue Engineering - Part A</i> , 2013 , 19, 2373-81	3.9	46
68	Sorafenib Acts Synergistically in Combination with Radiotherapy without Causing Intestinal Damage in Colorectal Cancer. <i>Tumori</i> , 2013 , 99, 176-182	1.7	9
67	Sorafenib acts synergistically in combination with radiotherapy without causing intestinal damage in colorectal cancer. <i>Tumori</i> , 2013 , 99, 176-82	1.7	6
66	Role of reactive oxygen species in transforming growth factor-beta1-induced extracellular matrix accumulation in renal tubular epithelial cells. <i>Transplantation Proceedings</i> , 2012 , 44, 625-8	1.1	29
65	Adenosine receptors are up-regulated in unilateral ureteral obstructed rat kidneys. <i>Transplantation Proceedings</i> , 2012 , 44, 1166-8	1.1	8
64	High glucose increases mesangial lipid accumulation via impaired cholesterol transporters. <i>Transplantation Proceedings</i> , 2012 , 44, 1021-5	1.1	12
63	Fractalkine increases mesangial cell proliferation through reactive oxygen species and mitogen-activated protein kinases. <i>Transplantation Proceedings</i> , 2012 , 44, 1026-8	1.1	11
62	Lipopolysaccharide increases monocyte binding to mesangial cells through fractalkine and its receptor. <i>Transplantation Proceedings</i> , 2012 , 44, 1029-31	1.1	5
61	Determination of daumone in mouse plasma by HPLC/MS-MS. <i>Biomedical Chromatography</i> , 2012 , 26, 152-5	1.7	3

60	Peroxiredoxin 3 is a key molecule regulating adipocyte oxidative stress, mitochondrial biogenesis, and adipokine expression. <i>Antioxidants and Redox Signaling</i> , 2012 , 16, 229-43	8.4	114
59	Human umbilical cord blood-derived mesenchymal stem cells prevent diabetic renal injury through paracrine action. <i>Diabetes Research and Clinical Practice</i> , 2012 , 98, 465-73	7.4	79
58	Catalase deficiency accelerates diabetic renal injury through peroxisomal dysfunction. <i>Diabetes</i> , 2012 , 61, 728-38	0.9	120
57	8-Hydroxy-2-deoxyguanosine prevents plaque formation and inhibits vascular smooth muscle cell activation through Rac1 inactivation. <i>Free Radical Biology and Medicine</i> , 2012 , 53, 109-21	7.8	21
56	Glucose-based peritoneal dialysis solution suppresses adiponectin synthesis through oxidative stress in an experimental model of peritoneal dialysis. <i>Peritoneal Dialysis International</i> , 2012 , 32, 20-8	2.8	13
55	Reactive oxygen species and oxidative stress. <i>Contributions To Nephrology</i> , 2011 , 170, 102-112	1.6	25
54	Antioxidant treatment may protect pancreatic beta cells through the attenuation of islet fibrosis in an animal model of type 2 diabetes. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 414, 397-402	3.4	44
53	Rapid and reliable measurement for evaluating directly the reactivity of N-acetylcysteine with glucose degradation products in peritoneal dialysis fluids. <i>Analytical Chemistry</i> , 2011 , 83, 1518-22	7.8	2
52	Quantitative determination of daumone in rat plasma by liquid chromatography-mass spectrometry. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2011 , 56, 114-7	3.5	11
51	Renoprotective antioxidant effect of alagebrium in experimental diabetes. <i>Nephrology Dialysis Transplantation</i> , 2011 , 26, 3474-84	4.3	16
50	Plasminogen activator inhibitor-1 antisense oligodeoxynucleotides abrogate mesangial fibronectin accumulation. <i>Korean Journal of Physiology and Pharmacology</i> , 2010 , 14, 385-90	1.8	6
49	Stability of N-acetylcysteine in peritoneal dialysis solution. <i>Peritoneal Dialysis International</i> , 2010 , 30, 105-8	2.8	3
48	The role of plasminogen activator inhibitor 1 in renal and cardiovascular diseases. <i>Nature Reviews Nephrology</i> , 2009 , 5, 203-11	14.9	109
47	Positive feedback loop between plasminogen activator inhibitor-1 and transforming growth factor-beta1 during renal fibrosis in diabetes. <i>American Journal of Nephrology</i> , 2009 , 30, 481-90	4.6	49
46	Histone deacetylase-2 is a key regulator of diabetes- and transforming growth factor-beta1-induced renal injury. <i>American Journal of Physiology - Renal Physiology</i> , 2009 , 297, F729-39	4.3	202
45	Wnt/beta-catenin signaling: a novel target for therapeutic intervention of fibrotic kidney disease. <i>Archives of Pharmacal Research</i> , 2009 , 32, 1653-62	6.1	51
44	Lithospermic acid B ameliorates the development of diabetic nephropathy in OLETF rats. <i>European Journal of Pharmacology</i> , 2008 , 579, 418-25	5.3	26
43	Mycophenolic acid inhibits cell proliferation and extracellular matrix synthesis in rat vascular smooth muscle cells through direct and indirect inhibition of cellular reactive oxygen species. <i>Journal of Surgical Research</i> , 2008 , 150, 17-23	2.5	12

42	Role of reactive oxygen species in the pathogenesis of diabetic nephropathy. <i>Diabetes Research and Clinical Practice</i> , 2008 , 82 Suppl 1, S42-5	7.4	168
41	Inducible nitric oxide synthase-nitric oxide plays an important role in acute and severe hypoxic injury to pancreatic beta cells. <i>Transplantation</i> , 2008 , 85, 323-30	1.8	35
40	Protective Effects of Lithospermic Acid B on Diabetic Nephropathy in OLETF Rats Comparing with Amlodipine and Losartan. <i>Korean Diabetes Journal</i> , 2008 , 32, 10		
39	Antifibrotic effect of globular adiponectin in human hepatocyte. <i>FASEB Journal</i> , 2008 , 22, 978.11	0.9	
38	Histone deacetylase 2 plays an important role in the development and progression of diabetic renal injury. <i>FASEB Journal</i> , 2008 , 22, 944.5	0.9	
37	Mechanisms of epithelial-mesenchymal transition of peritoneal mesothelial cells during peritoneal dialysis. <i>Journal of Korean Medical Science</i> , 2007 , 22, 943-5	4.7	28
36	Mycophenolic acid inhibits oleic acid-induced vascular smooth muscle cell activation by inhibiting cellular reactive oxygen species. <i>Transplantation</i> , 2007 , 84, 634-8	1.8	4
35	Mycophenolic acid inhibits mesangial cell activation through p38 MAPK inhibition. <i>Life Sciences</i> , 2006 , 79, 1561-7	6.8	26
34	Carvedilol inhibits platelet-derived growth factor-induced extracellular matrix synthesis by inhibiting cellular reactive oxygen species and mitogen-activated protein kinase activation. <i>Journal of Heart and Lung Transplantation</i> , 2006 , 25, 683-9	5.8	13
33	Reactive oxygen species mediate high glucose-induced plasminogen activator inhibitor-1 up-regulation in mesangial cells and in diabetic kidney. <i>Kidney International</i> , 2005 , 67, 1762-71	9.9	101
32	Reactive oxygen species amplify glucose signalling in renal cells cultured under high glucose and in diabetic kidney. <i>Nephrology</i> , 2005 , 10 Suppl, S7-10	2.2	102
31	Plasminogen activator inhibitor-1 and diabetic nephropathy. <i>Nephrology</i> , 2005 , 10 Suppl, S11-3	2.2	49
30	Angiotensin II Mediates High Glucose-Induced TGF- β 1 and Fibronectin Upregulation in HPMC through Reactive Oxygen Species. <i>Peritoneal Dialysis International</i> , 2005 , 25, 38-47	2.8	81
29	Role of reactive oxygen species in TGF-beta1-induced mitogen-activated protein kinase activation and epithelial-mesenchymal transition in renal tubular epithelial cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2005 , 16, 667-75	12.7	430
28	Oxidative stress and chronic allograft nephropathy. <i>Yonsei Medical Journal</i> , 2004 , 45, 1049-52	3	24
27	Reactive oxygen species amplify protein kinase C signaling in high glucose-induced fibronectin expression by human peritoneal mesothelial cells. <i>Kidney International</i> , 2004 , 65, 1170-9	9.9	125
26	Mycophenolic acid inhibits platelet-derived growth factor-induced reactive oxygen species and mitogen-activated protein kinase activation in rat vascular smooth muscle cells. <i>American Journal of Transplantation</i> , 2004 , 4, 1982-90	8.7	46
25	Evidence for heme oxygenase-1 association with caveolin-1 and -2 in mouse mesangial cells. <i>IUBMB Life</i> , 2003 , 55, 525-32	4.7	37

24	Reactive oxygen species mediate TGF-beta1-induced plasminogen activator inhibitor-1 upregulation in mesangial cells. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 309, 961-6	3.4	89
23	Reactive oxygen species-regulated signaling pathways in diabetic nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2003 , 14, S241-5	12.7	377
22	Delayed treatment with lithospermate B attenuates experimental diabetic renal injury. <i>Journal of the American Society of Nephrology: JASN</i> , 2003 , 14, 709-20	12.7	59
21	Agmatine reduces hydrogen peroxide in mesangial cells under high glucose conditions. <i>BMB Reports</i> , 2003 , 36, 251-7	5.5	12
20	Peritoneal mesothelial cell biology in peritoneal dialysis. <i>Nephrology</i> , 2002 , 7, 220-226	2.2	2
19	Effects of carvedilol alone and in the presence of cyclosporine A on the DNA synthesis of cultured vascular smooth muscle cells. <i>Surgery Today</i> , 2002 , 32, 230-5	3	5
18	High glucose increases inducible NO production in cultured rat mesangial cells. Possible role in fibronectin production. <i>Nephron</i> , 2002 , 90, 78-85	3.3	38
17	Effects of Peritoneal Dialysis Solutions on the Secretion of Growth Factors and Extracellular Matrix Proteins by Human Peritoneal Mesothelial Cells. <i>Peritoneal Dialysis International</i> , 2002 , 22, 171-177	2.8	75
16	Role of high glucose-induced nuclear factor-kappaB activation in monocyte chemoattractant protein-1 expression by mesangial cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2002 , 13, 894-902	12.7	218
15	Oxidative stress in diabetic nephropathy: basic and clinical information. <i>Current Diabetes Reports</i> , 2001 , 1, 282-7	5.6	81
14	High glucose-induced PKC activation mediates TGF-beta 1 and fibronectin synthesis by peritoneal mesothelial cells. <i>Kidney International</i> , 2001 , 59, 463-70	9.9	126
13	Peritoneal dialysis in diabetic patients. <i>American Journal of Kidney Diseases</i> , 2001 , 38, S200-3	7.4	16
12	Activation of protein kinase c-delta and c-epsilon by oxidative stress in early diabetic rat kidney. <i>American Journal of Kidney Diseases</i> , 2001 , 38, S204-7	7.4	50
11	Reactive oxygen species as glucose signaling molecules in mesangial cells cultured under high glucose. <i>Kidney International</i> , 2000 , 77, S19-25	9.9	215
10	Effect of High Glucose on Peritoneal Mesothelial Cell Biology. <i>Peritoneal Dialysis International</i> , 2000 , 20, 15-18	2.8	52
9	Biocompatibility of New Peritoneal Dialysis Solutions. <i>Peritoneal Dialysis International</i> , 2000 , 20, 3-4	2.8	4
8	Effects of Conventional and New Peritoneal Dialysis Solutions on Human Peritoneal Mesothelial Cell Viability and Proliferation. <i>Peritoneal Dialysis International</i> , 2000 , 20, 10-18	2.8	44
7	Future of Interventions in Diabetic Nephropathy: Antioxidants. <i>Peritoneal Dialysis International</i> , 1999 , 19, 228-233	2.8	3

6	Melatonin and taurine reduce early glomerulopathy in diabetic rats. <i>Free Radical Biology and Medicine</i> , 1999 , 26, 944-50	7.8	120
5	Pathogenesis of diabetic nephropathy: the role of oxidative stress and protein kinase C. <i>Diabetes Research and Clinical Practice</i> , 1999 , 45, 147-51	7.4	145
4	Sequential effects of high glucose on mesangial cell transforming growth factor-beta 1 and fibronectin synthesis. <i>Kidney International</i> , 1998 , 54, 1872-8	9.9	103
3	A high glucose concentration stimulates the expression of monocyte chemotactic peptide 1 in human mesangial cells. <i>Nephron</i> , 1998 , 79, 33-7	3.3	79
2	High glucose can induce lipid peroxidation in the isolated rat glomeruli. <i>Kidney International</i> , 1994 , 46, 1620-6	9.9	34
1	Amelioration of diabetic microalbuminuria and lipid peroxidation by captopril. <i>Yonsei Medical Journal</i> , 1992 , 33, 217-23	3	11