

# Giammaria Fiorentini

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5084290/publications.pdf>

Version: 2024-02-01

46  
papers

1,091  
citations

331642

21  
h-index

414395

32  
g-index

46  
all docs

46  
docs citations

46  
times ranked

1188  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Intra-arterial infusion of irinotecan-loaded drug-eluting beads (DEBIRI) versus intravenous therapy (FOLFIRI) for hepatic metastases from colorectal cancer: final results of a phase III study. <i>Anticancer Research</i> , 2012, 32, 1387-95.      | 1.1 | 236       |
| 2  | Trans-arterial chemoembolization of metastatic colorectal carcinoma to the liver adopting DC Bead® <sup>®</sup> , drug-eluting bead loaded with irinotecan: results of a phase II clinical study. <i>Anticancer Research</i> , 2011, 31, 4581-7.      | 1.1 | 65        |
| 3  | Intraarterial hepatic chemoembolization of liver metastases from colorectal cancer adopting irinotecan-eluting beads: results of a phase II clinical study. <i>In Vivo</i> , 2007, 21, 1085-91.   | 1.3 | 64        |
| 4  | Hyperthermia today: Electric energy, a new opportunity in cancer treatment. <i>Journal of Cancer Research and Therapeutics</i> , 2006, 2, 41.   | 0.9 | 53        |
| 5  | Quo Vadis Oncological Hyperthermia (2020)? <i>Frontiers in Oncology</i> , 2020, 10, 1690.   | 2.8 | 40        |
| 6  | A phase II clinical study on relapsed malignant gliomas treated with electro-hyperthermia. <i>In Vivo</i> , 2006, 20, 721-4.  | 1.3 | 40        |
| 7  | Multidisciplinary approach of colorectal cancer liver metastases. <i>World Journal of Clinical Oncology</i> , 2017, 8, 190.   | 2.3 | 37        |
| 8  | Chemoembolization in colorectal liver metastases: the rebirth. <i>Anticancer Research</i> , 2014, 34, 575-84.   | 1.1 | 35        |
| 9  | Modulated Electro-Hyperthermia as Palliative Treatment for Pancreatic Cancer: A Retrospective Observational Study on 106 Patients. <i>Integrative Cancer Therapies</i> , 2019, 18, 153473541987850.   | 2.0 | 33        |
| 10 | Irinotecan Hepatic Arterial Infusion Chemotherapy for Hepatic Metastases from Colorectal Cancer: A Phase II Clinical Study. <i>Tumori</i> , 2003, 89, 382-384.  | 1.1 | 32        |
| 11 | Transarterial chemoembolization with DC Bead LUMI® <sup>®</sup> , radiopaque beads for primary liver cancer treatment: preliminary experience. <i>Future Oncology</i> , 2017, 13, 2243-2252.  | 2.4 | 32        |
| 12 | Chemoembolization with Drug-eluting Microspheres Loaded with Doxorubicin for the Treatment of Cholangiocarcinoma. <i>Anticancer Research</i> , 2017, 37, 1859-1863.   | 1.1 | 30        |
| 13 | Modulated Electrohyperthermia in Integrative Cancer Treatment for Relapsed Malignant Glioblastoma and Astrocytoma: Retrospective Multicenter Controlled Study. <i>Integrative Cancer Therapies</i> , 2019, 18, 153473541881269.                       | 2.0 | 28        |
| 14 | Updates of colorectal cancer liver metastases therapy: review on DEBIRI. <i>Hepatic Oncology</i> , 2020, 7, HEP16.  | 4.2 | 27        |
| 15 | Deliberate hypoxic pelvic and limb chemoperfusion in the treatment of recurrent melanoma. <i>American Journal of Surgery</i> , 2002, 183, 28-36.  | 1.8 | 25        |
| 16 | Oxaliplatin hepatic arterial infusion chemotherapy for hepatic metastases from colorectal cancer: a phase I-II clinical study. <i>Anticancer Research</i> , 2004, 24, 2093-6.   | 1.1 | 25        |
| 17 | MGMT methylation correlates with melphalan pelvic perfusion survival in stage III melanoma patients: a pilot study. <i>Melanoma Research</i> , 2017, 27, 439-447.   | 1.2 | 24        |
| 18 | Hepatic arterial chemotherapy in combination with systemic chemotherapy compared with hepatic arterial chemotherapy alone for liver metastases from colorectal cancer: results of a multi-centric randomized study. <i>In Vivo</i> , 2006, 20, 707-9. | 1.3 | 23        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Hepatic Arterial Infusion of Polyethylene Glycol Drug-eluting Beads for Primary and Metastatic Liver Cancer Therapy. <i>Anticancer Research</i> , 2016, 36, 3515-21.   | 1.1 | 22        |
| 20 | Hypoxic pelvic and limb perfusion with melphalan and mitomycin C for recurrent limb melanoma. <i>Melanoma Research</i> , 2003, 13, 51-58.  | 1.2 | 21        |
| 21 | TACE of liver metastases from colorectal cancer adopting irinotecan-eluting beads: beneficial effect of palliative intra-arterial lidocaine and post-procedure supportive therapy on the control of side effects. <i>Hepato-Gastroenterology</i> , 2008, 55, 2077-82.                                  | 0.5 | 21        |
| 22 | Locoregional therapy and systemic cetuximab to treat colorectal liver metastases. <i>World Journal of Gastrointestinal Oncology</i> , 2015, 7, 47.   | 2.0 | 17        |
| 23 | A Narrative Review of Regional Hyperthermia: Updates From 2010 to 2019. <i>Integrative Cancer Therapies</i> , 2020, 19, 153473542093264.   | 2.0 | 17        |
| 24 | Chemoembolization Alone or Associated With Bevacizumab for Therapy of Colorectal Cancer Metastases: Preliminary Results of a Randomized Study. <i>In Vivo</i> , 2020, 34, 683-686.   | 1.3 | 17        |
| 25 | Real-life multidisciplinary treatment for unresectable colorectal cancer liver metastases including hepatic artery infusion with chemo-filtration and liquid biopsy precision oncotherapy: observational cohort study. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 1273-1290. | 2.5 | 14        |
| 26 | Mitomycin C hypoxic pelvic perfusion for unresectable recurrent rectal cancer: pharmacokinetic comparison of surgical and percutaneous techniques. <i>Updates in Surgery</i> , 2017, 69, 403-410.  | 2.0 | 12        |
| 27 | Polyethylene glycol microspheres loaded with irinotecan for arterially directed embolic therapy of metastatic liver cancer. <i>World Journal of Gastrointestinal Oncology</i> , 2017, 9, 379.  | 2.0 | 12        |
| 28 | Irinotecan hepatic arterial infusion chemotherapy for hepatic metastases from colorectal cancer: a phase II clinical study. <i>Tumori</i> , 2003, 89, 382-4.   | 1.1 | 11        |
| 29 | Thyroid Metastases from Colorectal Cancer: No Longer a Rare Entity. <i>Tumori</i> , 2006, 92, 465-466.   | 1.1 | 10        |
| 30 | Surgical versus percutaneous isolated pelvic perfusion (IPP) for advanced melanoma: comparison in terms of melphalan pharmacokinetic pelvic bio-availability. <i>BMC Research Notes</i> , 2017, 10, 411.   | 1.4 | 10        |
| 31 | Does Locoregional Chemotherapy Still Matter in the Treatment of Advanced Pelvic Melanoma?. <i>International Journal of Molecular Sciences</i> , 2017, 18, 2382.  | 4.1 | 8         |
| 32 | Melphalan hypoxic perfusion with hemofiltration for melanoma locoregional metastases in the pelvis. <i>Journal of Surgical Research</i> , 2017, 215, 114-124.  | 1.6 | 7         |
| 33 | A review discussing the use of polyethylene glycol microspheres in the treatment of hepatocellular carcinoma. <i>Future Oncology</i> , 2019, 15, 695-703.  | 2.4 | 7         |
| 34 | Circulating tumour cell liquid biopsy in selecting therapy for recurrent cutaneous melanoma with locoregional pelvic metastases: a pilot study. <i>BMC Research Notes</i> , 2020, 13, 176.   | 1.4 | 7         |
| 35 | Complete response of colorectal liver metastases after intra-arterial chemotherapy. <i>Tumori</i> , 2008, 94, 489-92.  | 1.1 | 7         |
| 36 | Chemoembolization in Conjunction with Bevacizumab: Preliminary Results. <i>Journal of Vascular and Interventional Radiology</i> , 2018, 29, 1236-1239.   | 0.5 | 5         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Modulated electro-hyperthermia in stage III and IV pancreatic cancer: Results of an observational study on 158 patients. <i>World Journal of Clinical Oncology</i> , 2021, 12, 1064-1071.                                | 2.3 | 5         |
| 38 | Isolated thoracic perfusion in lung metastases from breast cancer: a retrospective observational study. <i>Updates in Surgery</i> , 2019, 71, 165-177.   | 2.0 | 4         |
| 39 | Immune response activation following hyperthermic intraperitoneal chemotherapy for peritoneal metastases: A pilot study. <i>World Journal of Clinical Oncology</i> , 2020, 11, 397-404.                                  | 2.3 | 3         |
| 40 | Transarterial chemoembolization alone or followed by bevacizumab for treatment of colorectal liver metastases. <i>Hepatic Oncology</i> , 2022, 9, HEP40.   | 4.2 | 2         |
| 41 | A Prospective Study of Intraarterial Infusion Chemotherapy in Advanced Wild-Type BRAF Melanoma Patients. <i>Journal of Surgical Research</i> , 2021, 268, 737-747.   | 1.6 | 1         |
| 42 | Thoracic stop-flow perfusion for refractory lymphoma: a phase I-II evaluation trial. <i>In Vivo</i> , 2009, 23, 447-57.  | 1.3 | 1         |
| 43 | Immune response and locoregional treatments for peritoneal carcinomatosis. <i>International Review of Cell and Molecular Biology</i> , 2022, , 97-116.   | 3.2 | 1         |
| 44 | The Way Forward to Develop Locoregional Treatments in Oligometastatic Colorectal Cancer. <i>CardioVascular and Interventional Radiology</i> , 2020, 43, 1484-1485.   | 2.0 | 0         |
| 45 | Re: Response to Recommendation of Regional Hyperthermia in the Treatment of Breast Cancer. <i>Integrative Cancer Therapies</i> , 2021, 20, 153473542098858.  | 2.0 | 0         |
| 46 | Imatinib mesylate induces responses in patients with liver metastases from gastrointestinal stromal tumor failing intra-arterial hepatic chemotherapy. <i>Journal of Cancer Research and Therapeutics</i> , 2006, 2, 68. | 0.9 | 0         |