

# Chang-Sheng Xia

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5099709/publications.pdf>

Version: 2024-02-01

25  
papers

238  
citations

1040056

9  
h-index

1058476

14  
g-index

25  
all docs

25  
docs citations

25  
times ranked

325  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Imbalance of Circulating Follicular Helper T Cells and Follicular Regulatory T Cells Is Associated With Disease Activity in Patients With Ulcerative Colitis. <i>Frontiers in Immunology</i> , 2020, 11, 104.	4.8	28
2	Relapse predictors and serologically unstable condition of IgG4-related disease: a large Chinese cohort. <i>Rheumatology</i> , 2020, 59, 2115-2123.	1.9	24
3	Salivary gland involvement disparities in clinical characteristics of IgG4-related disease: a retrospective study of 428 patients. <i>Rheumatology</i> , 2020, 59, 634-640.	1.9	23
4	Elevated circulating pro-inflammatory low-density granulocytes in adult-onset Still's disease. <i>Rheumatology</i> , 2021, 60, 297-303.	1.9	22
5	Diagnostic performances of serum IgG4 concentration and IgG4/IgG ratio in IgG4-related disease. <i>Clinical Rheumatology</i> , 2017, 36, 2769-2774.	2.2	20
6	Recovery of CD226-TIGIT <sup>+</sup> FoxP3 <sup>+</sup> and CD226-TIGIT <sup>+</sup> FoxP3 <sup>+</sup> regulatory T cells contributes to clinical remission from active stage in ulcerative colitis patients. <i>Immunology Letters</i> , 2020, 218, 30-39.	2.5	13
7	Upregulated IL-17A secretion and CCR6 expression in Treg subsets are related to the imbalance of Treg/Th17 cells in active UC patients. <i>Scandinavian Journal of Immunology</i> , 2020, 91, e12842.	2.7	12
8	Clinical features of IgG4-related retroperitoneal fibrosis among 407 patients with IgG4-related disease: a retrospective study. <i>Rheumatology</i> , 2021, 60, 767-772.	1.9	12
9	Increased circulating PD-1 <sup>hi</sup> CXCR5 <sup>+</sup> peripheral helper T cells are associated with disease severity of active ulcerative colitis patients. <i>Immunology Letters</i> , 2021, 233, 2-10.	2.5	12
10	Increased proportion of functional subpopulations in circulating regulatory T cells in patients with chronic hepatitis B. <i>Hepatology Research</i> , 2020, 50, 439-452.	3.4	10
11	Activated inducible co-stimulator-positive programmed cell death 1-positive follicular helper T cells indicate disease activity and severity in ulcerative colitis patients. <i>Clinical and Experimental Immunology</i> , 2020, 202, 106-118.	2.6	10
12	Analysis of human cytomegalovirus glycoprotein N genotypes in Chinese hematopoietic stem cell transplant recipients. <i>Archives of Virology</i> , 2011, 156, 17-23.	2.1	8
13	Human Cytomegalovirus Glycoprotein B Genotypes in Chinese Hematopoietic Stem Cell Transplant Recipients. <i>Intervirology</i> , 2012, 55, 342-348.	2.8	7
14	Infection of Epstein-Barr Virus is Associated with the Decrease of Helios <sup>+</sup> FoxP3 <sup>+</sup> Regulatory T Cells in Active Ulcerative Colitis Patients. <i>Immunological Investigations</i> , 2021, 50, 23-36.	2.0	5
15	Circulating CD4 <sup>+</sup> FoxP3 <sup>+</sup> CXCR5 <sup>+</sup> CXCR3 <sup>+</sup> cells are elevated in active rheumatoid arthritis and reflect the severity of the disease. <i>International Journal of Rheumatic Diseases</i> , 2021, 24, 1032-1039.	1.9	5
16	Evaluation of three automated <i>Treponema pallidum</i> antibody assays for syphilis screening. <i>Journal of Infection and Chemotherapy</i> , 2018, 24, 887-891.	1.7	4
17	Performance evaluation of the mindray anti-HCV assay for the detection of hepatitis C virus infection. <i>Journal of Clinical Laboratory Analysis</i> , 2018, 32, e22600.	2.1	4
18	Increased Circulating Th1 and Tfh1 Cell Numbers Are Associated with Disease Activity in Glucocorticoid-Treated Patients with IgG4-Related Disease. <i>Journal of Immunology Research</i> , 2020, 2020, 1-8.	2.2	4

#	ARTICLE	IF	CITATIONS
19	Use of the Nuclear Matrix Protein 22 BladderChek Test for the Detection of Primary and Recurrent Urothelial Carcinoma. <i>Disease Markers</i> , 2020, 2020, 1-6.	1.3	4
20	Serum Human Epididymis Protein 4 as a Novel Biomarker in Identifying Patients With Interstitial Lung Disease in Rheumatoid Arthritis. <i>Frontiers in Medicine</i> , 2021, 8, 755268.	2.6	4
21	Elevated Granulocyte Colony-stimulating Factor Levels in Patients With Active Phase of Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2021, 48, 664-668.	2.0	3
22	IL-7 Promotes the Expansion of Circulating CD28- Cytotoxic T Lymphocytes in Patients With IgG4-Related Disease via the JAK Signaling. <i>Frontiers in Immunology</i> , 0, 13, .	4.8	3
23	The altered HLAâ€DQ expression in peripheral blood T cells of chronic hepatitis B patients characterizes the function of T cells. <i>Journal of Viral Hepatitis</i> , 2022, 29, 340-351.	2.0	1
24	THU0591â€...SALIVARY GLAND INVOLVEMENT DISPARITIES IN CLINICAL AND LABORATORY CHARACTERISTICS OF IGG4-RELATED DISEASE: A RETROSPECTIVE STUDY OF 428 PATIENTS. , 2019, , .		0
25	Interleukin 10 inhibits the rat VSMC proliferation and collagen secretion stimulated by angiotensin II. <i>Chinese Medical Sciences Journal</i> , 2001, 16, 125-8.	0.4	0