

# Andrew J Gooday

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5106076/publications.pdf>

Version: 2024-02-01

212  
papers

13,783  
citations

28274

55  
h-index

24982

109  
g-index

212  
all docs

212  
docs citations

212  
times ranked

7909  
citing authors

#	ARTICLE	IF	CITATIONS
1	Scientific and budgetary trade-offs between morphological and molecular methods for deep-sea biodiversity assessment. <i>Integrated Environmental Assessment and Management</i> , 2022, 18, 655-663.	2.9	7
2	Patterns of eukaryotic diversity from the surface to the deep-ocean sediment. <i>Science Advances</i> , 2022, 8, eabj9309.	10.3	52
3	New species of <i>Gromia</i> (Protista, Rhizaria) from South Georgia and the Falkland Islands. <i>Polar Biology</i> , 2022, 45, 647-666.	1.2	3
4	Assessment of scientific gaps related to the effective environmental management of deep-seabed mining. <i>Marine Policy</i> , 2022, 138, 105006.	3.2	67
5	New observations on test architecture and construction of <i>Jullienella foetida</i> Schlumberger, 1890, the largest shallow-water agglutinated foraminifer in modern oceans. <i>PeerJ</i> , 2022, 10, e12884.	2.0	0
6	Taxon-rich transcriptomics supports higher-level phylogeny and major evolutionary trends in Foraminifera. <i>Molecular Phylogenetics and Evolution</i> , 2022, 174, 107546.	2.7	6
7	Molecular and morphological diversity of monothalamous foraminifera from South Georgia and the Falkland Islands: Description of four new species. <i>European Journal of Protistology</i> , 2022, 85, 125909.	1.5	10
8	Ecological variables for deep-ocean monitoring must include microbiota and meiofauna for effective conservation. <i>Nature Ecology and Evolution</i> , 2021, 5, 27-29.	7.8	22
9	Three new species of <i>Gromia</i> (Protista, Rhizaria) from western Greenland fjords. <i>Polar Biology</i> , 2021, 44, 1037-1053.	1.2	2
10	The Biodiversity and Distribution of Abyssal Benthic Foraminifera and Their Possible Ecological Roles: A Synthesis Across the Clarion-Clipperton Zone. <i>Frontiers in Marine Science</i> , 2021, 8, .	2.5	18
11	Megafaunal Ecology of the Western Clarion Clipperton Zone. <i>Frontiers in Marine Science</i> , 2021, 8, .	2.5	13
12	Megafauna of the German exploration licence area for seafloor massive sulphides along the Central and South East Indian Ridge (Indian Ocean). <i>Biodiversity Data Journal</i> , 2021, 9, e69955.	0.8	5
13	Environment, ecology, and potential effectiveness of an area protected from deep-sea mining (Clarion) Tj ETQq1 1 0.784314 r gBT /Ov	3.2	36
14	Cenozoic climatic changes drive evolution and dispersal of coastal benthic foraminifera in the Southern Ocean. <i>Scientific Reports</i> , 2021, 11, 19869.	3.3	8
15	Review: Freshwater and Soil Foraminifera – A Story of Long-Forgotten Relatives. <i>Journal of Foraminiferal Research</i> , 2021, 51, 318-331.	0.5	16
16	Discovery of Paleodictyon in the Indian Ocean. <i>Marine Biodiversity</i> , 2021, 51, 1.	1.0	1
17	Giant, highly diverse protists in the abyssal Pacific: vulnerability to impacts from seabed mining and potential for recovery. <i>Communicative and Integrative Biology</i> , 2020, 13, 189-197.	1.4	15
18	Xenophyophores (Rhizaria, Foraminifera), including four new species and two new genera, from the western Clarion-Clipperton Zone (abyssal equatorial Pacific). <i>European Journal of Protistology</i> , 2020, 75, 125715.	1.5	14

#	ARTICLE	IF	CITATIONS
19	Protist diversity and function in the dark ocean – Challenging the paradigms of deep-sea ecology with special emphasis on foraminiferans and naked protists. <i>European Journal of Protistology</i> , 2020, 75, 125721.	1.5	40
20	Loricifera inhabiting spherical agglutinated structures in the abyssal eastern equatorial Pacific nodule fields. <i>Marine Biodiversity</i> , 2019, 49, 2455-2466.	1.0	5
21	Diversity and spatial patterns of foraminiferal assemblages in the eastern Clarion-Clipperton zone (abyssal eastern equatorial Pacific). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2019, 149, 103036.	1.4	18
22	Soft-walled monothalamids (Rhizaria: foraminifera) of the Crimean shelf (Black Sea): taxonomic composition and inter-regional patterns of species diversity and distribution. <i>Marine Biology Research</i> , 2019, 15, 36-48.	0.7	0
23	The Contribution of Fine Sieve Fractions (63-150 µm) to Foraminiferal Abundance and Diversity in an Area of the Eastern Pacific Ocean Licensed for Polymetallic Nodule Exploration. <i>Frontiers in Marine Science</i> , 2019, 6, .	2.5	17
24	Bathyal benthic megafauna from the Mid-Atlantic Ridge in the region of the Charlie-Gibbs fracture zone based on remotely operated vehicle observations. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2019, 145, 1-12.	1.4	13
25	Key role of bacteria in the short-term cycling of carbon at the abyssal seafloor in a low particulate organic carbon flux region of the eastern Pacific Ocean. <i>Limnology and Oceanography</i> , 2019, 64, 694-713.	3.1	50
26	The chemical composition of a new –œmica sandwich– foraminiferal species from the East Coast of Korea: <i>Capsamina crassa</i> sp. nov.. <i>PeerJ</i> , 2019, 7, e6642.	2.0	1
27	Eyes of the Deep-sea Floor: The Integrative Taxonomy of the Foraminiferal Genus <i>Vanhoeffenella</i> . <i>Protist</i> , 2018, 169, 235-267.	1.5	6
28	Five new species and two new genera of xenophyophores (Foraminifera: Rhizaria) from part of the abyssal equatorial Pacific licensed for polymetallic nodule exploration. <i>Zoological Journal of the Linnean Society</i> , 2018, 183, 723-748.	2.3	20
29	Characteristics of meiofauna in extreme marine ecosystems: a review. <i>Marine Biodiversity</i> , 2018, 48, 35-71.	1.0	153
30	Xenophyophores (Rhizaria, Foraminifera) from the Eastern Clarion-Clipperton Zone (equatorial) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 30	1.5	9
31	New species of the xenophyophore genus <i>Aschemonella</i> (Rhizaria: Foraminifera) from areas of the abyssal eastern Pacific licensed for polymetallic nodule exploration. <i>Zoological Journal of the Linnean Society</i> , 2018, 182, 479-499.	2.3	14
32	First record of a Xenophyophore (Rhizaria: Foraminifera) on the Chilean margin. <i>Zootaxa</i> , 2018, 4455, 589.	0.5	1
33	Micro-CT 3D imaging reveals the internal structure of three abyssal xenophyophore species (Protista,) Tj ETQq1 1 0,784314 rgBT /Overlock 18	3.3	18
34	Relationship between –live–™ and dead benthic foraminiferal assemblages in the abyssal NE Atlantic. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2017, 121, 190-201.	1.4	13
35	Giant protists (xenophyophores, Foraminifera) are exceptionally diverse in parts of the abyssal eastern Pacific licensed for polymetallic nodule exploration. <i>Biological Conservation</i> , 2017, 207, 106-116.	4.1	60
36	Macrofaunal nematodes of the deep Whittard Canyon (<sc>NE</sc> Atlantic): assemblage characteristics and comparison with polychaetes. <i>Marine Ecology</i> , 2017, 38, e12408.	1.1	4

#	ARTICLE	IF	CITATIONS
37	Novel benthic foraminifera are abundant and diverse in an area of the abyssal equatorial Pacific licensed for polymetallic nodule exploration. <i>Scientific Reports</i> , 2017, 7, 45288.	3.3	35
38	Abundance and morphology of <i>Paleodictyon nodosum</i> , observed at the Clarion-Clipperton Zone. <i>Marine Biodiversity</i> , 2017, 47, 265-269.	1.0	13
39	Xenophyophores (Rhizaria, Foraminifera) from the Russian license area of the Clarion-Clipperton Zone (eastern equatorial Pacific), with the description of three new species. <i>Marine Biodiversity</i> , 2017, 47, 299-306.	1.0	11
40	From the Surface to the Deep-Sea: Bacterial Distributions across Polymetallic Nodule Fields in the Clarion-Clipperton Zone of the Pacific Ocean. <i>Frontiers in Microbiology</i> , 2017, 8, 1696.	3.5	54
41	Major impacts of climate change on deep-sea benthic ecosystems. <i>Elementa</i> , 2017, 5, .	3.2	252
42	Worldwide Analysis of Sedimentary DNA Reveals Major Gaps in Taxonomic Knowledge of Deep-Sea Benthos. <i>Frontiers in Marine Science</i> , 2016, 3, .	2.5	138
43	Abyssal hills: Influence of topography on benthic foraminiferal assemblages. <i>Progress in Oceanography</i> , 2016, 148, 44-55.	3.2	26
44	Insights into the abundance and diversity of abyssal megafauna in a polymetallic-nodule region in the eastern Clarion-Clipperton Zone. <i>Scientific Reports</i> , 2016, 6, 30492.	3.3	173
45	The Whittard Canyon – A case study of submarine canyon processes. <i>Progress in Oceanography</i> , 2016, 146, 38-57.	3.2	68
46	Formation of agglutinated cysts by the foraminiferan <i>Sphaeroidina bulloides</i> on the Porcupine Abyssal Plain (NE Atlantic). <i>Marine Biodiversity</i> , 2016, 46, 747-749.	1.0	5
47	Agglutination of benthic foraminifera in relation to mesoscale bathymetric features in the abyssal NE Atlantic (Porcupine Abyssal Plain). <i>Marine Micropaleontology</i> , 2016, 123, 15-28.	1.2	15
48	Taxonomic composition and distribution of soft-walled monothalamid foraminifera in the area of Zernovâ€™s<i>Phyllophora</i>Field (NW Black Sea). <i>Marine Biology Research</i> , 2016, 12, 640-646.	0.7	4
49	MEMORIAL TO MARTIN DAVID BRASIER (1947â€™2014). <i>Journal of Foraminiferal Research</i> , 2016, 46, 221-223.	0.5	2
50	The trophic and metabolic pathways of foraminifera in the Arabian Sea: evidence from cellular stable isotopes. <i>Biogeosciences</i> , 2015, 12, 1781-1797.	3.3	13
51	Living (Rose-Bengal-stained) benthic foraminiferal faunas along a strong bottom-water oxygen gradient on the Indian margin (Arabian Sea). <i>Biogeosciences</i> , 2015, 12, 5005-5019.	3.3	26
52	Abyssal foraminifera attached to polymetallic nodules from the eastern Clarion Clipperton Fracture Zone: a preliminary description and comparison with North Atlantic dropstone assemblages. <i>Marine Biodiversity</i> , 2015, 45, 391-412.	1.0	39
53	Macrofaunal abundance and community composition at lower bathyal depths in different branches of the Whittard Canyon and on the adjacent slope (3500 m; NE Atlantic). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2015, 97, 29-39.	1.4	26
54	Is the meiofauna a good indicator for climate change and anthropogenic impacts?. <i>Marine Biodiversity</i> , 2015, 45, 505-535.	1.0	209

#	ARTICLE	IF	CITATIONS
55	Xenophyophores (Protista, Foraminifera) from the Clarion-Clipperton Fracture Zone with description of three new species. <i>Marine Biodiversity</i> , 2015, 45, 581-593.	1.0	22
56	Radiolarian tests as microhabitats for novel benthic foraminifera: Observations from the abyssal eastern equatorial Pacific (Clarion-Clipperton Fracture Zone). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2015, 103, 73-85.	1.4	13
57	Benthic polychaete diversity patterns and community structure in the Whittard Canyon system and adjacent slope (NE Atlantic). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2015, 106, 42-54.	1.4	14
58	Basal monothalamous and pseudo-chambered benthic foraminifera associated with planktonic foraminiferal shells and mineral grains from the Porcupine Abyssal Plain, NE Atlantic. <i>Marine Biodiversity</i> , 2015, 45, 357-369.	1.0	4
59	Uptake of algal carbon and the likely synthesis of an 'essential' fatty acid by <i>Uvigerina</i> , ex. gr. <i>semiornata</i> (Foraminifera) within the Pakistan margin oxygen minimum zone: evidence from fatty acid biomarker and $\delta^{13}C$ tracer experiments. <i>Biogeosciences</i> , 2014, 11, 3729-3738.	3.3	19
60	Agglutinated foraminifera (superfamily Hormosinacea) across the Indian margin oxygen minimum zone (Arabian Sea). <i>Marine Biodiversity</i> , 2014, 44, 5-25.	1.0	7
61	Basal foraminifera and gromiids (Protista) at the Håkon-Mosby Mud Volcano (Barents Sea slope). <i>Marine Biodiversity</i> , 2013, 43, 205-225.	1.0	9
62	Giant protists (xenophyophores and komokiaceans) from the Clarion-Clipperton ferromanganese nodule field (eastern Pacific). <i>Biology Bulletin Reviews</i> , 2013, 3, 388-398.	0.9	27
63	Genetic Diversity and Environmental Preferences of Monothalamous Foraminifers Revealed through Clone Analysis of Environmental Small-Subunit Ribosomal DNA Sequences. <i>Journal of Foraminiferal Research</i> , 2013, 43, 3-13.	0.5	6
64	New and poorly known benthic foraminifera (Protista, Rhizaria) inhabiting the shells of planktonic foraminifera on the bathyal Mid-Atlantic Ridge. <i>Marine Biology Research</i> , 2013, 9, 447-461.	0.7	14
65	Trawled megafaunal invertebrate assemblages from bathyal depth of the Mid-Atlantic Ridge (48°N-54°N). <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2013, 98, 326-340.	1.4	18
66	Deep-sea surface-dwelling enteropneusts from the Mid-Atlantic Ridge: Their ecology, distribution and mode of life. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2013, 98, 374-387.	1.4	22
67	The ecology and biogeography of <i>Discospirina tenuissima</i> (Foraminifera) in the Atlantic and Indian Oceans. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2013, 98, 301-314.	1.4	8
68	Two new genera and species of the monothalamous foraminifera from coastal waters of the Black Sea. <i>Marine Biodiversity</i> , 2013, 43, 473-479.	1.0	7
69	2012 Cushman Award to Khadyzhat M. Saidova. <i>Journal of Foraminiferal Research</i> , 2013, 43, 1-2.	0.5	0
70	Biotic and Human Vulnerability to Projected Changes in Ocean Biogeochemistry over the 21st Century. <i>PLoS Biology</i> , 2013, 11, e1001682.	5.6	194
71	Does Presence of a Mid-Ocean Ridge Enhance Biomass and Biodiversity?. <i>PLoS ONE</i> , 2013, 8, e61550.	2.5	68
72	Macrofaunal colonization across the Indian margin oxygen minimum zone. <i>Biogeosciences</i> , 2013, 10, 7161-7177.	3.3	9

#	ARTICLE	IF	CITATIONS
73	The Porcupine Abyssal Plain fixed-point sustained observatory (PAP-SO): variations and trends from the Northeast Atlantic fixed-point time-series. <i>ICES Journal of Marine Science</i> , 2012, 69, 776-783.	2.5	50
74	Possible early foraminiferans in post-Sturtian (716-635 Ma) cap carbonates. <i>Geology</i> , 2012, 40, 67-70.	4.4	66
75	Benthic Foraminiferal Biogeography: Controls on Global Distribution Patterns in Deep-Water Settings. <i>Annual Review of Marine Science</i> , 2012, 4, 237-262.	11.6	102
76	Foraminiferal biodiversity associated with cold-water coral carbonate mounds and open slope of SE Rockall Bank (Irish continental margin, NE Atlantic). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2012, 59, 54-71.	1.4	17
77	Meiobenthos of the Oxidic/Anoxic Interface in the Southwestern Region of the Black Sea: Abundance and Taxonomic Composition. <i>Cellular Origin and Life in Extreme Habitats</i> , 2012, , 369-401.	0.3	15
78	The influence of productivity on abyssal foraminiferal biodiversity. <i>Marine Biodiversity</i> , 2012, 42, 415-431.	1.0	18
79	Large, enigmatic foraminiferan-like protists in the eastern part of the Clarion-Clipperton Fracture Zone (abyssal north-eastern subequatorial Pacific): biodiversity and vertical distribution in the sediment. <i>Marine Biodiversity</i> , 2012, 42, 311-327.	1.0	14
80	Possible effects of global environmental changes on Antarctic benthos: a synthesis across five major taxa. <i>Ecology and Evolution</i> , 2012, 2, 453-485.	1.9	88
81	Intracellular mineral grains in the xenophyophore <i>Nazareammina tenera</i> (Rhizaria, Foraminifera) from the Nazaré Canyon (Portuguese margin, NE Atlantic). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2011, 58, 1189-1195.	1.4	10
82	Xenophyophores (Rhizaria, Foraminifera) from the Nazaré Canyon (Portuguese margin, NE Atlantic). <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2011, 58, 2401-2419.	1.4	36
83	A new <i>saccamminid</i> ™ genus (Rhizaria: Foraminifera), from 4400 m water depth in the Nazaré Canyon (NE) Tj ETQq1 1 0.7843 0.5 1	0.5	1
84	Testing the protozoan hypothesis for Ediacaran fossils: a developmental analysis of <i>Palaeopascichnus</i> . <i>Palaeontology</i> , 2011, 54, 1157-1175.	2.2	66
85	Grazing of intertidal benthic foraminifera on bacteria: Assessment using pulse-chase radiotracing. <i>Journal of Experimental Marine Biology and Ecology</i> , 2011, 399, 25-34.	1.5	43
86	Biodiversity and distribution of the genus <i>Gromia</i> (Protista, Rhizaria) in the deep Weddell Sea (Southern Ocean). <i>Polar Biology</i> , 2011, 34, 69-81.	1.2	16
87	New genera and species of monothalamous Foraminifera from Balaclava and Kazachâ€™ya Bays (Crimean) Tj ETQq1 1 0.7843 1.0 24 rgBT	1.0	24
88	The organic-walled genera <i>Resigella</i> and <i>Conicotheca</i> (Protista, Foraminifera) at two Arctic deep-sea sites (North Pole and Barents Sea), including the description of a new species of <i>Resigella</i> . <i>Marine Biodiversity</i> , 2010, 40, 33-44.	1.0	5
89	Ontogenetic effects on stable carbon and oxygen isotopes in tests of live (Rose Bengal stained) benthic foraminifera from the Pakistan continental margin. <i>Marine Micropaleontology</i> , 2010, 76, 92-103.	1.2	17
90	Habitat heterogeneity and its influence on benthic biodiversity in oxygen minimum zones. <i>Marine Ecology</i> , 2010, 31, 125-147.	1.1	126

#	ARTICLE	IF	CITATIONS
91	The roles of habitat heterogeneity in generating and maintaining biodiversity on continental margins: an introduction. <i>Marine Ecology</i> , 2010, 31, 1-5.	1.1	116
92	Biological structures as a source of habitat heterogeneity and biodiversity on the deep ocean margins. <i>Marine Ecology</i> , 2010, 31, 21-50.	1.1	490
93	Deep-Sea Biodiversity in the Mediterranean Sea: The Known, the Unknown, and the Unknowable. <i>PLoS ONE</i> , 2010, 5, e11832.	2.5	321
94	Natural and human-induced hypoxia and consequences for coastal areas: synthesis and future development. <i>Biogeosciences</i> , 2010, 7, 1443-1467.	3.3	358
95	Traces of dissolved particles, including coccoliths, in the tests of agglutinated foraminifera from the Challenger Deep (10,897m water depth, western equatorial Pacific). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2010, 57, 239-247.	1.4	12
96	Temporal changes (1989–1999) in deep-sea metazoan meiofaunal assemblages on the Porcupine Abyssal Plain, NE Atlantic. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2010, 57, 1383-1395.	1.4	44
97	Decadal-scale changes in shallow-infaunal foraminiferal assemblages at the Porcupine Abyssal Plain, NE Atlantic. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2010, 57, 1362-1382.	1.4	51
98	Global genetic homogeneity in the deep-sea foraminiferan <i>Epistominella exigua</i> (Rotaliida). <i>Journal of Foraminiferal Research</i> , 2010, 40, 462-477.	0.5	47
99	Europe's Grand Canyon: Nazaré Submarine Canyon. <i>Oceanography</i> , 2009, 22, 46-57.	1.0	86
100	Historical records of coastal eutrophication-induced hypoxia. <i>Biogeosciences</i> , 2009, 6, 1707-1745.	3.3	134
101	Effects of natural and human-induced hypoxia on coastal benthos. <i>Biogeosciences</i> , 2009, 6, 2063-2098.	3.3	525
102	A new genus and two new species of saccamminid foraminiferans (Protista, Rhizaria) from the deep Southern Ocean. <i>Zootaxa</i> , 2009, 2096, 9-22.	0.5	17
103	A minute new species of <i>Saccamina</i> (monothalamous Foraminifera). <i>Journal of Foraminiferal Research</i> , 2009, 39, 17-21.	3.6	17
104	Precambrian Biota: Protistan Origin of Trace Fossils?. <i>Current Biology</i> , 2009, 19, R28-R30.	3.9	9
105	A new genus of xenophyophores (Foraminifera) from Japan Trench: morphological description, molecular phylogeny and elemental analysis. <i>Zoological Journal of the Linnean Society</i> , 2009, 156, 455-464.	2.3	31
106	Megafaunal responses to strong oxygen gradients on the Pakistan margin of the Arabian Sea. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2009, 56, 472-487.	1.4	40
107	Foraminiferal faunal responses to monsoon-driven changes in organic matter and oxygen availability at 140 and 300m water depth in the NE Arabian Sea. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2009, 56, 403-421.	1.4	50
108	Faunal responses to oxygen gradients on the Pakistan margin: A comparison of foraminiferans, macrofauna and megafauna. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2009, 56, 488-502.	1.4	148

#	ARTICLE	IF	CITATIONS
109	Large organic-walled Protista (Gromia) in the Arabian Sea: Density, diversity, distribution and ecology. Deep-Sea Research Part II: Topical Studies in Oceanography, 2009, 56, 422-433.	1.4	19
110	Paleodictyon nodosum: A living fossil on the deep-sea floor. Deep-Sea Research Part II: Topical Studies in Oceanography, 2009, 56, 1700-1712.	1.4	56
111	Foraminiferal faunal responses to monsoon-driven changes in organic matter and oxygen availability at 140 and 300m water depth in the NE Arabian Sea. Deep-Sea Research Part II: Topical Studies in Oceanography, 2009, 56, 403-421.	1.4	11
112	New species of Leptohalysis (Rhizaria, Foraminifera) from an extreme hadal site in the western Pacific Ocean. Zootaxa, 2009, 2059, 23-32.	0.5	2
113	Genetic differentiation between Arctic and Antarctic monothalamous foraminiferans. Polar Biology, 2008, 31, 1205-1216.	1.2	43
114	Modern deep-sea benthic foraminifera: a brief review of their morphology-based biodiversity and trophic diversity. Geological Society Special Publication, 2008, 303, 97-119.	1.3	45
115	New organic-walled Foraminifera (Protista) from the ocean's deepest point, the Challenger Deep (western Pacific Ocean). Zoological Journal of the Linnean Society, 2008, 153, 399-423.	2.3	28
116	Exponential Decline of Deep-Sea Ecosystem Functioning Linked to Benthic Biodiversity Loss. Current Biology, 2008, 18, 1-8.	3.9	641
117	The enigmatic, deep-sea, organic-walled genera <i>Chitinosiphon</i> , <i>Nodellum</i> and <i>Resigella</i> (Protista, Foraminifera): A taxonomic re-evaluation. Systematics and Biodiversity, 2008, 6, 385-404.	1.2	8
118	Trophic modes of large Antarctic Foraminifera: roles of carnivory, omnivory, and detritivory. Marine Ecology - Progress Series, 2008, 371, 155-164.	1.9	31
119	DISTRIBUTION AND BIODIVERSITY OF STAINED MONOTHALAMOUS FORAMINIFERA FROM TEMPELFJORD, SVALBARD. Journal of Foraminiferal Research, 2007, 37, 93-106.	0.5	34
120	Oxygen as a control on sea floor biological communities and their roles in sedimentary carbon cycling. Limnology and Oceanography, 2007, 52, 1698-1709.	3.1	146
121	Ferromanganese nodule fauna in the Tropical North Pacific Ocean: Species richness, faunal cover and spatial distribution. Deep-Sea Research Part I: Oceanographic Research Papers, 2007, 54, 1912-1935.	1.4	73
122	Influence of surface texture and microhabitat heterogeneity in structuring nodule faunal communities. Deep-Sea Research Part I: Oceanographic Research Papers, 2007, 54, 1936-1943.	1.4	32
123	Introduction to ANDEEP, summary and outlook. Deep-Sea Research Part II: Topical Studies in Oceanography, 2007, 54, 1645-1651.	1.4	24
124	The biodiversity and biogeography of komokiaceans and other enigmatic foraminiferan-like protists in the deep Southern Ocean. Deep-Sea Research Part II: Topical Studies in Oceanography, 2007, 54, 1691-1719.	1.4	25
125	A note on the genetic similarity between shallow- and deep-water <i>Epistominella vitrea</i> (Foraminifera) in the Antarctic. Deep-Sea Research Part II: Topical Studies in Oceanography, 2007, 54, 1720-1726.	1.4	28
126	First insights into the biodiversity and biogeography of the Southern Ocean deep sea. Nature, 2007, 447, 307-311.	27.8	417



#	ARTICLE	IF	CITATIONS
127	New and little-known Komokiacea (Foraminifera) from the bathyal and abyssal Weddell Sea and adjacent areas. <i>Zoological Journal of the Linnean Society</i> , 2007, 151, 219-251.	2.3	15
128	Bipolar gene flow in deep-sea benthic foraminifera. <i>Molecular Ecology</i> , 2007, 16, 4089-4096.	3.9	132
129	Live (Rose Bengal stained) and dead benthic foraminifera from the oxygen minimum zone of the Pakistan continental margin (Arabian Sea). <i>Marine Micropaleontology</i> , 2007, 62, 45-73.	1.2	116
130	Siliceous scales of filose-amoebae (Pompholyxophryidae, Rotosphaerida) from deep Southern Ocean sediments, including first records for the Southern Hemisphere. <i>Polar Biology</i> , 2007, 30, 945-950.	1.2	7
131	Live benthic foraminifera at an abyssal site in the equatorial Pacific nodule province: Abundance, diversity and taxonomic composition. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2006, 53, 1406-1422.	1.4	47
132	High diversity of deep-sea <i>Gromia</i> from the Arabian Sea revealed by small subunit rDNA sequence analysis. <i>Marine Biology</i> , 2006, 148, 769-777.	1.5	10
133	BENTHIC FORAMINIFERAL TRENDS IN RELATION TO AN ORGANIC ENRICHMENT GRADIENT ON THE CONTINENTAL SLOPE (850 M WATER DEPTH) OFF NORTH CAROLINA (USA). <i>Journal of Foraminiferal Research</i> , 2006, 36, 34-43.	0.5	13
134	<i>Tinogullmia lukyanovae</i> sp. nov. a monothalamous, organic-walled foraminiferan from the coastal Black Sea. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2006, 86, 43-49.	0.8	12
135	Organic and siliceous protistan scales in north-east Atlantic abyssal sediments. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2006, 86, 679-688.	0.8	4
136	The Second Species of <i>Gromia</i> (Protista) from the Deep Sea: its Natural History and Association with the Pakistan Margin Oxygen Minimum Zone. <i>Protist</i> , 2005, 156, 113-126.	1.5	17
137	Monothalamous foraminiferans and gromiids (Protista) from western Svalbard: A preliminary survey. Published in collaboration with the University of Bergen and the Institute of Marine Research, Norway, and the Marine Biological Laboratory, University of Copenhagen, Denmark. <i>Marine Biology Research</i> , 2005, 1, 290-312.	0.7	52
138	Simple Foraminifera Flourish at the Ocean's Deepest Point. <i>Science</i> , 2005, 307, 689-689.	12.6	63
139	The monothalamous foraminiferan <i>Tinogullmia</i> in the Black Sea. <i>Journal of Micropalaeontology</i> , 2005, 24, 191-192.	3.6	11
140	<i>Hyperammina micaceus</i> sp. nov.: a new foraminiferan species (Protista) from the Porcupine Abyssal Plain, northeast Atlantic. <i>Journal of Micropalaeontology</i> , 2004, 23, 171-179.	3.6	2
141	Soft-walled, monothalamous benthic foraminiferans in the Pacific, Indian and Atlantic Oceans: aspects of biodiversity and biogeography. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2004, 51, 33-53.	1.4	47
142	Associations between living benthic foraminifera and dead tests of <i>Syringammina fragilissima</i> (Xenophyophorea) in the Darwin Mounds region (NE Atlantic). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2004, 51, 1741-1758.	1.4	49
143	Live (stained) deep-sea benthic foraminiferans in the western Weddell Sea: trends in abundance, diversity and taxonomic composition along a depth transect. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2004, 51, 1571-1602.	1.4	70
144	A new monothalamous foraminiferan from 1000 to 6300m water depth in the Weddell Sea: morphological and molecular characterisation. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2004, 51, 1603-1616.	1.4	47

#	ARTICLE	IF	CITATIONS
145	Introduction to ANDEEP (ANTarctic benthic DEEP-sea biodiversity: colonization history and recent) Tj ETQq1 1 0.784314 rgBT /Overlook Oceanography, 2004, 51, 1457-1465.	1.4	74
146	Soft-shelled monothalamous foraminifera at an intertidal site on the south coast of England. Journal of Micropalaeontology, 2004, 23, 135-137.	3.6	12
147	Conqueria laevis gen. and sp. nov., a new soft-walled, monothalamous foraminiferan from the deep Weddell Sea. Journal of the Marine Biological Association of the United Kingdom, 2004, 84, 919-924.	0.8	33
148	Vellaria zucchellii sp. nov. a new monothalamous foraminifer from Terra Nova Bay, Antarctica. Antarctic Science, 2004, 16, 307-312.	0.9	17
149	Associations between living benthic foraminifera and dead tests of Syringammina fragilissima (Xenophyophorea) in the Darwin Mounds region (NE Atlantic). Deep-Sea Research Part I: Oceanographic Research Papers, 2004, 51, 1741-1758.	1.4	2
150	Benthic foraminifera (protista) as tools in deep-water palaeoceanography: Environmental influences on faunal characteristics. Advances in Marine Biology, 2003, 46, 1-90.	1.4	377
151	The evolution of early Foraminifera. Proceedings of the National Academy of Sciences of the United States of America, 2003, 100, 11494-11498.	7.1	227
152	Selective feeding by benthic foraminifera on phytodetritus on the western Antarctic Peninsula shelf: evidence from fatty acid biomarker analysis. Marine Ecology - Progress Series, 2003, 262, 153-162.	1.9	127
153	A THEME ISSUE ON THE BIOLOGY AND BIODIVERSITY OF 'ALLOGROMIID' FORAMINIFERA. Journal of Foraminiferal Research, 2002, 32, 331-333.	0.5	12
154	STRUCTURE, TAXONOMY AND ECOLOGY OF ASTRAMMINA TRIANGULARIS (EARLAND), AN ALLOGROMIID-LIKE AGGLUTINATED FORAMINIFER FROM EXPLORERS COVE, ANTARCTICA. Journal of Foraminiferal Research, 2002, 32, 364-374.	0.5	27
155	Soft-shelled benthic foraminifera from a hadal site (7800 m water depth) in the Atacama Trench (SE) Tj ETQq1 1 0.784314 rgBT /Overlook 3.6 22	3.6	22
156	ORGANIC-WALLED ALLOGROMIIDS: ASPECTS OF THEIR OCCURRENCE, DIVERSITY AND ECOLOGY IN MARINE HABITATS. Journal of Foraminiferal Research, 2002, 32, 384-399.	0.5	95
157	Foraminifera associated with phytodetritus deposits at a bathyal site in the northern Rockall Trough (NE Atlantic): seasonal contrasts and a comparison of stained and dead assemblages. Marine Micropaleontology, 2002, 46, 83-110.	1.2	134
158	Biological Responses to Seasonally Varying Fluxes of Organic Matter to the Ocean Floor: A Review. Journal of Oceanography, 2002, 58, 305-332.	1.7	247
159	Ecology and nutrition of the large agglutinated foraminiferan Bathysiphon capillare in the bathyal NE Atlantic: distribution within the sediment profile and lipid biomarker composition. Marine Ecology - Progress Series, 2002, 245, 69-82.	1.9	45
160	The foraminiferan macrofauna from three North Carolina (USA) slope sites with contrasting carbon flux: a comparison with the metazoan macrofauna. Deep-Sea Research Part I: Oceanographic Research Papers, 2001, 48, 1709-1739.	1.4	53
161	Environmental Influences on Regional Deep-Sea Species Diversity. Annual Review of Ecology, Evolution, and Systematics, 2001, 32, 51-93.	6.7	607
162	Title is missing!. Journal of Oceanography, 2001, 57, 377-384.	1.7	22

#	ARTICLE	IF	CITATIONS
163	Morphological and ecological parallels between sublittoral and abyssal foraminiferal species in the NE Atlantic: a comparison of <i>Stainforthia fusiformis</i> and <i>Stainforthia</i> sp.. <i>Progress in Oceanography</i> , 2001, 50, 261-283.	3.2	33
164	Evidence for episodic recruitment in a small opheliid polychaete species from the abyssal NE Atlantic. <i>Progress in Oceanography</i> , 2001, 50, 285-301.	3.2	26
165	Temporal patterns among meiofauna and macrofauna taxa related to changes in sediment geochemistry at an abyssal NE Atlantic site. <i>Progress in Oceanography</i> , 2001, 50, 303-324.	3.2	85
166	Dressing up for the deep: agglutinated protists adorn an irregular urchin. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2001, 81, 881-882.	0.8	15
167	Impact of large-scale natural physical disturbance on the diversity of deep-sea North Atlantic nematodes. <i>Marine Ecology - Progress Series</i> , 2001, 214, 121-126.	1.9	49
168	Foraminifera in the Arabian Sea oxygen minimum zone and other oxygen-deficient settings: taxonomic composition, diversity, and relation to metazoan faunas. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2000, 47, 25-54.	1.4	240
169	A large testate protist, <i>Gromia sphaerica</i> sp. nov. (Order Filosea), from the bathyal Arabian Sea. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2000, 47, 55-73.	1.4	24
170	Temporal variability in living deep-sea benthic foraminifera: a review. <i>Earth-Science Reviews</i> , 1999, 46, 187-212.	9.1	168
171	Physical reworking by near-bottom flow alters the metazoan meiofauna of Fieberling Guyot (northeast Pacific). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 1999, 46, 2041-2052.	1.4	27
172	Deep-sea benthic foraminiferal species diversity in the NE Atlantic and NW Arabian sea: a synthesis. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 1998, 45, 165-201.	1.4	82
173	Recent benthic foraminifera in the abyssal Northeast Atlantic Ocean; relation to phytodetrital inputs. <i>Journal of Foraminiferal Research</i> , 1997, 27, 85-92.	0.5	65
174	Large, deep-sea agglutinated Foraminifera; two differing kinds of organization and their possible ecological significance. <i>Journal of Foraminiferal Research</i> , 1997, 27, 278-291.	0.5	50
175	Epifaunal and shallow infaunal foraminiferal communities at three abyssal NE Atlantic sites subject to differing phytodetritus input regimes. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 1996, 43, 1395-1421.	1.4	124
176	Cenozoic deep-sea benthic foraminifers: Tracers for changes in oceanic productivity?. <i>Geology</i> , 1996, 24, 355.	4.4	210
177	Xenophyophores (Protista) including two new species, from two abyssal sites in the Northeast Atlantic Ocean. <i>Journal of Foraminiferal Research</i> , 1996, 26, 193-208.	0.5	15
178	Benthic foraminiferal assemblages in Explorers Cove, Antarctica: A shallow-water site with deep-sea characteristics. <i>Progress in Oceanography</i> , 1996, 37, 117-166.	3.2	78
179	Micro- and Nanoforaminifera from Abyssal Northeast Atlantic Sediments: A Preliminary Report. <i>International Review of Hydrobiology</i> , 1995, 80, 361-383.	0.6	35
180	Larger agglutinated foraminifera of McMurdo Sound, Antarctica: Are <i>Astrammmina rara</i> and <i>Notodendrodes antarctikos</i> allogromiids incognito?. <i>Marine Micropaleontology</i> , 1995, 26, 75-88.	1.2	53

#	ARTICLE	IF	CITATIONS
181	The taxonomy and ecology of <i>Crithionina delacai</i> sp. nov., and abundant large agglutinated foraminifer from Explorers Cove, Antarctica. <i>Journal of Foraminiferal Research</i> , 1995, 25, 290-298.	0.5	19
182	A benthic foraminiferal proxy of pulsed organic matter paleofluxes. <i>Marine Micropaleontology</i> , 1994, 23, 89-99.	1.2	153
183	The Biology of Deep-Sea Foraminifera: A Review of Some Advances and Their Applications in Paleoceanography. <i>Palaios</i> , 1994, 9, 14.	1.3	264
184	A new large agglutinated foraminifer ( <i>Arboramminidae</i> n. fam.) from an oligotrophic site in the abyssal Northeast Atlantic. <i>Journal of Foraminiferal Research</i> , 1994, 24, 149-157.	0.5	9
185	The morphology and ecology of an abundant new komokiacean mudball ( <i>Komokiacea</i> , <i>Foraminiferida</i> ) from the bathyal and abyssal NE Atlantic. <i>Journal of Foraminiferal Research</i> , 1994, 24, 214-225.	0.5	13
186	Microforaminifera – Perspective on a Neglected Group of Foraminifera. <i>Archiv für Protistenkunde</i> , 1993, 143, 271-284.	0.8	13
187	Deep-sea benthic foraminiferal species which exploit phytodetritus: Characteristic features and controls on distribution. <i>Marine Micropaleontology</i> , 1993, 22, 187-205.	1.2	368
188	Direct observation of episodic growth in an abyssal xenophyophore (Protista). <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 1993, 40, 2131-2143.	1.4	54
189	Maintenance of abyssal benthic foraminifera under high pressure and low temperature; some preliminary results. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 1993, 40, 643-652.	1.4	39
190	A New Allogromiid Genus ( <i>Rhizopoda</i> : <i>Foraminiferida</i> ) from the Vellar Estuary, Bay of Bengal. <i>Journal of Micropalaeontology</i> , 1992, 11, 233-239.	3.6	18
191	Some recent advances in the study of deep-sea foraminiferal biology and their palaeoecological significance. <i>The Paleontological Society Special Publications</i> , 1992, 6, 111-111.	0.0	0
192	The distribution and ecology of <i>Bathysiphon filiformis</i> Sars and <i>B. major de Folin</i> (Protista). <i>Journal of Foraminiferal Research</i> , 1992, 22, 129-146.	0.5	60
193	The Role of Benthic Foraminifera in Deep-Sea Food Webs and Carbon Cycling. , 1992, , 63-91.		121
194	Xenophyophores (Protista, Rhizopoda) in box-core samples from the abyssal Northeast Atlantic Ocean, BIOTRANS area; their taxonomy, morphology, and ecology. <i>Journal of Foraminiferal Research</i> , 1991, 21, 197-212.	0.5	22
195	Recent Deep-Sea Agglutinated Foraminifera: A Brief Review. , 1990, , 271-304.		23
196	The impact of seasonally deposited phytodetritus on epifaunal and shallow infaunal benthic foraminiferal populations in the bathyal northeast Atlantic: the assemblage response. <i>Deep-sea Research Part A, Oceanographic Research Papers</i> , 1990, 37, 1263-1283.	1.5	70
197	The morphology, internal organization, and taxonomic position of <i>Rhizammina algaeformis</i> Brady, a large, agglutinated, deep-sea foraminifer. <i>Journal of Foraminiferal Research</i> , 1989, 19, 115-125.	0.5	37
198	The genus <i>Bathysiphon</i> (Protista, Foraminiferida) in the northeast Atlantic: SEM observations on the wall structure of seven species. <i>Journal of Natural History</i> , 1989, 23, 591-611.	0.5	23

#	ARTICLE	IF	CITATIONS
199	A response by benthic Foraminifera to the deposition of phytodetritus in the deep sea. <i>Nature</i> , 1988, 332, 70-73.	27.8	560
200	The genus <i>Bathysiphon</i> (Protista, Foraminiferida) in the north-east Atlantic: a neotype for <i>B. filiformis</i> G.O. & M. Sars, 1872 and the description of a new species. <i>Journal of Natural History</i> , 1988, 22, 95-105.	0.5	13
201	The genus <i>Bathysiphon</i> (Protista, Foraminiferida) in the NE Atlantic: revision of some species described by de Folin (1886). <i>Journal of Natural History</i> , 1988, 22, 71-93.	0.5	19
202	New xenophyophores (Protista) from the bathyal and abyssal north-east Atlantic Ocean. <i>Journal of Natural History</i> , 1988, 22, 413-434.	0.5	21
203	Soft-shelled foraminifera in meiofaunal samples from the bathyal northeast Atlantic. <i>Sarsia</i> , 1986, 71, 275-287.	0.5	58
204	Meiofaunal foraminiferans from the bathyal Porcupine Seabight (northeast Atlantic): size structure, standing stock, taxonomic composition, species diversity and vertical distribution in the sediment. <i>Deep-sea Research Part A, Oceanographic Research Papers</i> , 1986, 33, 1345-1373.	1.5	238
205	The genus <i>Rhabdammina</i> in the Northeast Atlantic; a new species, a redescription of <i>R. major</i> de Folin, 1887, and some speculations on species relationships. <i>Journal of Foraminiferal Research</i> , 1986, 16, 150-160.	0.5	11
206	An association between komokiacean foraminifers (Protozoa) and paludicelline ctenostomes (Bryozoa) from the abyssal Northeast Atlantic. <i>Journal of Natural History</i> , 1984, 18, 765-784.	0.5	23
207	Records of deep-sea rhizopod tests inhabited by metazoans in the North-east Atlantic. <i>Sarsia</i> , 1984, 69, 45-53.	0.5	51
208	<i>Bathysiphon rusticus</i> de Folin, 1886 and <i>Bathysiphon folini</i> n. sp.; two large agglutinated foraminifers abundant in abyssal NE Atlantic epibenthic sledge samples. <i>Journal of Foraminiferal Research</i> , 1983, 13, 262-276.	0.5	23
209	Abyssal foraminifers, including two new genera, encrusting the interior of <i>Bathysiphon rusticus</i> tubes. <i>Deep-sea Research Part A, Oceanographic Research Papers</i> , 1983, 30, 591-614.	1.5	27
210	Distribution of Planktonic Ostracoda (Halocyprididae) in the North Adriatic With the Description of a New Subspecies, <i>Conchoecia Porrecta Adriatica</i> . <i>Crustaceana</i> , 1977, 32, 139-154.	0.3	6
211	The sinking velocities of some halocyprid ostracods. <i>Journal of Experimental Marine Biology and Ecology</i> , 1975, 19, 105-116.	1.5	10
212	Three new species of deep-sea <i>Gromia</i> (Protista, Rhizaria) from the bathyal and abyssal Weddell Sea, Antarctica. <i>Zoological Journal of the Linnean Society</i> , 0, 157, 451-469.	2.3	11