

# Marisol Veny

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5125414/publications.pdf>

Version: 2024-02-01

16  
papers

378  
citations

1163117

8  
h-index

1281871

11  
g-index

16  
all docs

16  
docs citations

16  
times ranked

802  
citing authors

#	ARTICLE	IF	CITATIONS
1	Btla signaling in conventional and regulatory lymphocytes coordinately tempers humoral immunity in the intestinal mucosa. <i>Cell Reports</i> , 2022, 38, 110553.	6.4	9
2	Dissecting Common and Unique Effects of Anti-Î±4Î²7 and Anti-Tumor Necrosis Factor Treatment in Ulcerative Colitis. <i>Journal of Crohn's and Colitis</i> , 2021, 15, 441-452.	1.3	17
3	Expression Levels of 4 Genes in Colon Tissue Might Be Used to Predict Which Patients Will Enter Endoscopic Remission After Vedolizumab Therapy for Inflammatory Bowel Diseases. <i>Clinical Gastroenterology and Hepatology</i> , 2020, 18, 1142-1151.e10.	4.4	50
4	Contactin-1 Is Required for Peripheral Innervation and Immune Homeostasis Within the Intestinal Mucosa. <i>Frontiers in Immunology</i> , 2020, 11, 1268.	4.8	9
5	Controlling leukocyte trafficking in IBD. <i>Pharmacological Research</i> , 2020, 159, 105050.	7.1	14
6	Differences in Peripheral and Tissue Immune Cell Populations Following Haematopoietic Stem Cell Transplantation in Crohn's Disease Patients. <i>Journal of Crohn's and Colitis</i> , 2019, 13, 634-647.	1.3	13
7	TNF Superfamily in Inflammation. , 2018, , 1-50.		1
8	Sphingosine-1-phosphate receptor-1 (S1P1) is expressed by lymphocytes, dendritic cells, and endothelium and modulated during inflammatory bowel disease. <i>Mucosal Immunology</i> , 2017, 10, 162-171.	6.0	79
9	971a Inflammation Modulates the Sphingosine-1-Phosphate Pathway in Inflammatory Bowel Disease (IBD). <i>Gastroenterology</i> , 2016, 150, S195.	1.3	0
10	Chemokine receptor CCR7 regulates the intestinal TH1/TH17/Treg balance during Crohn's-like murine ileitis. <i>Journal of Leukocyte Biology</i> , 2015, 97, 1011-1022.	3.3	62
11	Letter: pathogenicity of Th17 cells may differ in ulcerative colitis compared with Crohn's disease " authors' reply. <i>Alimentary Pharmacology and Therapeutics</i> , 2012, 36, 205-205.	3.7	1
12	Late Crohn's disease patients present an increase in peripheral Th17 cells and cytokine production compared with early patients. <i>Alimentary Pharmacology and Therapeutics</i> , 2010, 31, 561-572.	3.7	66
13	Is Lack of "Education" a Mechanism Driving Loss of Tolerance in Crohn's Disease?. <i>Gastroenterology</i> , 2010, 139, 1056-1058.	1.3	0
14	Defective IL-10 production in severe phenotypes of Crohn's disease. <i>Journal of Leukocyte Biology</i> , 2009, 85, 896-903.	3.3	56
15	T1226 Th17 and Th1 Effector Responses in Crohn's Disease. <i>Gastroenterology</i> , 2008, 134, A-511.	1.3	0
16	Btla Signaling in Conventional and Regulatory Lymphocytes Coordinately Tempers Humoral Immunity in the Intestinal Mucosa. <i>SSRN Electronic Journal</i> , 0, , .	0.4	1