Gillian A Whalley

List of Publications by Citations

Source: https://exaly.com/author-pdf/5130747/gillian-a-whalley-publications-by-citations.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

142 papers

5,446 citations

36 h-index

71 g-index

166 ext. papers

6,297 ext. citations

avg, IF

5.61 L-index

#	Paper	IF	Citations
142	Predicting survival in heart failure: a risk score based on 39 372 patients from 30 studies. <i>European Heart Journal</i> , 2013 , 34, 1404-13	9.5	595
141	The survival of patients with heart failure with preserved or reduced left ventricular ejection fraction: an individual patient data meta-analysis. <i>European Heart Journal</i> , 2012 , 33, 1750-7	9.5	508
140	Left ventricular remodeling with carvedilol in patients with congestive heart failure due to ischemic heart disease. Australia-New Zealand Heart Failure Research Collaborative Group. <i>Journal of the American College of Cardiology</i> , 1997 , 29, 1060-6	15.1	329
139	Effects of carvedilol on left ventricular remodeling after acute myocardial infarction: the CAPRICORN Echo Substudy. <i>Circulation</i> , 2004 , 109, 201-6	16.7	244
138	Plasma amino-terminal pro-brain natriuretic peptide and accuracy of heart-failure diagnosis in primary care: a randomized, controlled trial. <i>Journal of the American College of Cardiology</i> , 2003 , 42, 179	93-800	192
137	Randomized, controlled trial of integrated heart failure management: The Auckland Heart Failure Management Study. <i>European Heart Journal</i> , 2002 , 23, 139-46	9.5	180
136	Does a glass of red wine improve endothelial function?. European Heart Journal, 2000, 21, 74-8	9.5	154
135	Ethnic-Specific Normative Reference Values for Echocardiographic LAD LV Size, LV Mass, and Systolic Function: The EchoNoRMAL Study. <i>JACC: Cardiovascular Imaging</i> , 2015 , 8, 656-65	8.4	125
134	The obesity paradox in heart failure patients with preserved versus reduced ejection fraction: a meta-analysis of individual patient data. <i>International Journal of Obesity</i> , 2014 , 38, 1110-4	5.5	123
133	Independent prognostic importance of a restrictive left ventricular filling pattern after myocardial infarction: an individual patient meta-analysis: Meta-Analysis Research Group in Echocardiography acute myocardial infarction. <i>Circulation</i> , 2008 , 117, 2591-8	16.7	123
132	Gender and survival in patients with heart failure: interactions with diabetes and aetiology. Results from the MAGGIC individual patient meta-analysis. <i>European Journal of Heart Failure</i> , 2012 , 14, 473-9	12.3	119
131	Renal dysfunction in patients with heart failure with preserved versus reduced ejection fraction: impact of the new Chronic Kidney Disease-Epidemiology Collaboration Group formula. <i>Circulation: Heart Failure</i> , 2012 , 5, 309-14	7.6	112
130	Independent relationship of left atrial size and mortality in patients with heart failure: an individual patient meta-analysis of longitudinal data (MeRGE Heart Failure). <i>European Journal of Heart Failure</i> , 2009 , 11, 929-36	12.3	109
129	The prognostic significance of heart failure with preserved left ventricular ejection fraction: a literature-based meta-analysis. <i>European Journal of Heart Failure</i> , 2009 , 11, 855-62	12.3	99
128	Three-dimensional assessment of left ventricular systolic strain in patients with type 2 diabetes mellitus, diastolic dysfunction, and normal ejection fraction. <i>American Journal of Cardiology</i> , 2004 , 94, 1391-5	3	99
127	Uptake of self-management strategies in a heart failure management programme. <i>European Journal of Heart Failure</i> , 2003 , 5, 371-80	12.3	98
126	Determinants of left ventricular hypertrophy and systolic dysfunction in chronic renal failure. <i>American Journal of Kidney Diseases</i> , 1994 , 24, 768-76	7.4	91

(2015-2002)

125	Pseudonormal mitral filling pattern predicts hospital re-admission in patients with congestive heart failure. <i>Journal of the American College of Cardiology</i> , 2002 , 39, 1787-95	15.1	85
124	Relationship of serum sodium concentration to mortality in a wide spectrum of heart failure patients with preserved and with reduced ejection fraction: an individual patient data meta-analysis(I): Meta-Analysis Global Group in Chronic heart failure (MAGGIC). European Journal of	12.3	74
123	Independence of restrictive filling pattern and LV ejection fraction with mortality in heart failure: an individual patient meta-analysis. <i>European Journal of Heart Failure</i> , 2008 , 10, 786-92	12.3	61
122	Left ventricular systolic and diastolic function assessed by tissue Doppler imaging and outcome in asymptomatic aortic stenosis. <i>European Heart Journal</i> , 2010 , 31, 2216-22	9.5	60
121	Understanding changing patterns of survival and hospitalization for heart failure over two decades in New Zealand: utility of @ ays alive and out of hospital @ rom epidemiological data. <i>European Journal of Heart Failure</i> , 2010 , 12, 462-8	12.3	52
120	Heart failure in younger patients: the Meta-analysis Global Group in Chronic Heart Failure (MAGGIC). <i>European Heart Journal</i> , 2014 , 35, 2714-21	9.5	50
119	Comparison of ultrasound assessment of flow-mediated dilatation in the radial and brachial artery with upper and forearm cuff positions. <i>Clinical Physiology</i> , 2001 , 21, 9-14		50
118	Association of genetic variation in the natriuretic peptide system with cardiovascular outcomes. Journal of Molecular and Cellular Cardiology, 2011 , 50, 695-701	5.8	49
117	Progression of myocardial remodeling and mechanical dysfunction in the spontaneously hypertensive rat. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2012 , 303, H1353-65	5.2	45
116	Longitudinal left ventricular contractile dysfunction after exercise in aortic stenosis. <i>Heart</i> , 2007 , 93, 732-8	5.1	45
115	Left ventricular diastolic filling and systolic function of young and older trained and untrained men. Journal of Applied Physiology, 2003 , 95, 2570-5	3.7	45
114	Doppler echocardiography and the early diagnosis of carditis in acute rheumatic fever. <i>Australian and New Zealand Journal of Medicine</i> , 1994 , 24, 530-5		44
113	Screening for left ventricular hypertrophy in patients with type 2 diabetes mellitus in the community. <i>Cardiovascular Diabetology</i> , 2011 , 10, 29	8.7	41
112	Effect of early initiation of dialysis on cardiac structure and function: results from the echo substudy of the IDEAL trial. <i>American Journal of Kidney Diseases</i> , 2013 , 61, 262-70	7.4	38
111	A community-based model of care improves blood pressure control and delays progression of proteinuria, left ventricular hypertrophy and diastolic dysfunction in Maori and Pacific patients with type 2 diabetes and chronic kidney disease: a randomized controlled trial. <i>Nephrology Dialysis</i>	4.3	38
110	Transplantation, 2010 , 25, 3260-6 Effects of improved glycaemic control on endothelial function in patients with type 2 diabetes. Internal Medicine Journal, 2001 , 31, 322-8	1.6	38
109	Is heart rate a risk marker in patients with chronic heart failure and concomitant atrial fibrillation? Results from the MAGGIC meta-analysis. <i>European Journal of Heart Failure</i> , 2015 , 17, 1182-91	12.3	37
108	Differing prognostic value of pulse pressure in patients with heart failure with reduced or preserved ejection fraction: results from the MAGGIC individual patient meta-analysis. <i>European Heart Journal</i> , 2015 , 36, 1106-14	9.5	37

107	A common variant at chromosome 9P21.3 is associated with age of onset of coronary disease but not subsequent mortality. <i>Circulation: Cardiovascular Genetics</i> , 2010 , 3, 286-93		37
106	Circulating miR-323-3p and miR-652: candidate markers for the presence and progression of acute coronary syndromes. <i>International Journal of Cardiology</i> , 2014 , 176, 375-85	3.2	36
105	Restrictive filling pattern is a powerful predictor of heart failure events postacute myocardial infarction and in established heart failure: a literature-based meta-analysis. <i>Journal of Cardiac Failure</i> , 2007 , 13, 346-52	3.3	36
104	Association of fat-free mass and training status with left ventricular size and mass in endurance-trained athletes. <i>Journal of the American College of Cardiology</i> , 2004 , 44, 892-6	15.1	36
103	The Validity of Left Ventricular Mass as a Surrogate End Point for All-Cause and Cardiovascular Mortality Outcomes in People With CKD: A Systematic Review and Meta-analysis. <i>American Journal of Kidney Diseases</i> , 2016 , 68, 554-563	7.4	36
102	Atrial fibrillation and the risk of death in patients with heart failure: a literature-based meta-analysis. <i>Internal Medicine Journal</i> , 2010 , 40, 347-56	1.6	32
101	Elevated B-type natriuretic peptide despite normal left ventricular function on rest and exercise stress echocardiography in mitral regurgitation. <i>European Heart Journal</i> , 2008 , 29, 363-70	9.5	32
100	Restrictive diastolic filling predicts death after acute myocardial infarction: systematic review and meta-analysis of prospective studies. <i>Heart</i> , 2006 , 92, 1588-94	5.1	32
99	Bias associated with left ventricular quantification by multimodality imaging: a systematic review and meta-analysis. <i>Open Heart</i> , 2016 , 3, e000388	3	31
98	The prognostic significance of restrictive diastolic filling associated with heart failure: a meta-analysis. <i>International Journal of Cardiology</i> , 2007 , 116, 70-7	3.2	31
97	Massive nitrogen loss in critical surgical illness: effect on cardiac mass and function. <i>Annals of Surgery</i> , 1997 , 226, 191-7	7.8	30
96	A randomized trial of the aldosterone-receptor antagonist eplerenone in asymptomatic moderate-severe aortic stenosis. <i>American Heart Journal</i> , 2008 , 156, 348-55	4.9	29
95	Structural and functional cardiac abnormalities in adolescent girls with poorly controlled type 2 diabetes. <i>Diabetes Care</i> , 2009 , 32, 883-8	14.6	28
94	Prognostic role of echocardiography and brain natriuretic peptide in symptomatic breathless patients in the community. <i>European Heart Journal</i> , 2008 , 29, 509-16	9.5	28
93	Diastolic Dysfunction Assessed Using Contemporary Guidelines and Prognosis Following Myocardial Infarction. <i>Journal of the American Society of Echocardiography</i> , 2018 , 31, 1127-1136	5.8	27
92	Genomic risk variants at 1p13.3, 1q41, and 3q22.3 are associated with subsequent cardiovascular outcomes in healthy controls and in established coronary artery disease. <i>Circulation: Cardiovascular Genetics</i> , 2011 , 4, 636-46		26
91	Pseudonormal mitral filling is associated with similarly poor prognosis as restrictive filling in patients with heart failure and coronary heart disease: a systematic review and meta-analysis of prospective studies. <i>Journal of the American Society of Echocardiography</i> , 2009 , 22, 494-8	5.8	25
90	Left ventricular mass correlates with fat-free mass but not fat mass in adults. <i>Journal of Hypertension</i> , 1999 , 17, 569-74	1.9	25

(2005-1997)

89	Effects of simvastatin and enalapril on serum lipoprotein concentrations and left ventricular mass in patients on dialysis. The Perfect Study Collaborative Group. <i>Journal of Nephrology</i> , 1997 , 10, 33-40	4.8	25
88	Prognostic significance of anaemia in patients with heart failure with preserved and reduced ejection fraction: results from the MAGGIC individual patient data meta-analysis. <i>QJM - Monthly Journal of the Association of Physicians</i> , 2016 , 109, 377-382	2.7	24
87	Known and missing left ventricular ejection fraction and survival in patients with heart failure: a MAGGIC meta-analysis report. <i>European Journal of Heart Failure</i> , 2013 , 15, 1220-7	12.3	22
86	Hyperuricaemia and gout in New Zealand rural and urban MBri and non-MBri communities. <i>Internal Medicine Journal</i> , 2013 , 43, 678-84	1.6	22
85	Geographic variation in left ventricular mass and mass index: a systematic review. <i>Journal of Human Hypertension</i> , 2012 , 26, 420-9	2.6	22
84	The effect of type 2 diabetes on diastolic function. <i>Medicine and Science in Sports and Exercise</i> , 2006 , 38, 1384-8	1.2	21
83	Echo and natriuretic peptide guided therapy improves outcome and reduces worsening renal function in systolic heart failure: An observational study of 1137 outpatients. <i>International Journal of Cardiology</i> , 2016 , 224, 416-423	3.2	20
82	Genetic variation in the renin-angiotensin-aldosterone system is associated with cardiovascular risk factors and early mortality in established coronary heart disease. <i>Journal of Human Hypertension</i> , 2013 , 27, 237-44	2.6	19
81	Angiotensin-converting enzyme 2 A1075G polymorphism is associated with survival in an acute coronary syndromes cohort. <i>American Heart Journal</i> , 2008 , 156, 752-8	4.9	19
80	Redefining normal reference ranges for echocardiography: a major new individual person data meta-analysis. <i>European Heart Journal Cardiovascular Imaging</i> , 2013 , 14, 347-8	4.1	18
79	A Kaupapa MBri approach to a community cohort study of heart disease in New Zealand. <i>Australian and New Zealand Journal of Public Health</i> , 2011 , 35, 249-55	2.3	18
78	A cohort study comparing cardiovascular risk factors in rural Maori, urban Maori and non-Maori communities in New Zealand. <i>BMJ Open</i> , 2012 , 2,	3	18
77	Increased B-type natriuretic peptide is associated with an abnormal blood pressure response to exercise in asymptomatic aortic stenosis. <i>International Journal of Cardiology</i> , 2008 , 127, 313-20	3.2	18
76	Effects of perindopril-indapamide on left ventricular diastolic function and mass in patients with type 2 diabetes: the ADVANCE Echocardiography Substudy. <i>Journal of Hypertension</i> , 2011 , 29, 1439-47	1.9	17
75	Individual patient meta-analyses of restrictive diastolic filling pattern and mortality in patients post acute myocardial infarction and in patients with chronic heart failure. <i>International Journal of Cardiology</i> , 2007 , 122, 207-15	3.2	17
74	The larger exercise stroke volume in endurance-trained men does not result from increased left ventricular early or late inflow or tissue velocities. <i>Acta Physiologica</i> , 2012 , 205, 520-31	5.6	16
73	Pseudonormal diastolic filling unmasked with glyceryl trinitrate in patients with type 2 diabetes with poor metabolic control. <i>Diabetes Care</i> , 2001 , 24, 1307-8	14.6	16
72	Comparison of different methods for detection of diastolic filling abnormalities. <i>Journal of the American Society of Echocardiography</i> , 2005 , 18, 710-7	5.8	15

71	Atherosclerosis and left ventricular hypertrophy: persisting problems in treated hypertensive patients. <i>Journal of Hypertension</i> , 1998 , 16, 1389-95	1.9	15
70	C-Type Natriuretic Peptides in Coronary Disease. Clinical Chemistry, 2017, 63, 316-324	5.5	14
69	Heart failure with preserved ejection fraction. <i>Journal of Geriatric Cardiology</i> , 2013 , 10, 369-76	1.7	14
68	Effects of carvedilol on left ventricular regional wall motion in patients with heart failure caused by ischemic heart disease. Australia-New Zealand Heart Failure Research Collaborative Group. <i>Journal of Cardiac Failure</i> , 2000 , 6, 11-8	3.3	13
67	Left ventricular geometry and all-cause mortality in advanced age. <i>Heart Lung and Circulation</i> , 2015 , 24, 32-9	1.8	12
66	Chronic measurement of left ventricular pressure in freely moving rats. <i>Journal of Applied Physiology</i> , 2013 , 115, 1672-82	3.7	12
65	Assessment and impact of diastolic function by echocardiography in elderly patients. <i>Journal of Geriatric Cardiology</i> , 2016 , 13, 252-60	1.7	12
64	New Diastology Guidelines: Evolution, Validation and Impact on Clinical Practice. <i>Heart Lung and Circulation</i> , 2019 , 28, 1411-1420	1.8	11
63	The prognostic impact of diastolic dysfunction in patients with chronic heart failure and post-acute myocardial infarction: Can age-stratified E/A ratio alone predict survival?. <i>International Journal of Cardiology</i> , 2015 , 181, 362-8	3.2	11
62	KCNE5 polymorphism rs697829 is associated with QT interval and survival in acute coronary syndromes patients. <i>Journal of Cardiovascular Electrophysiology</i> , 2012 , 23, 319-24	2.7	11
61	Effect of tissue harmonic imaging and contrast upon between observer and test-retest reproducibility of left ventricular ejection fraction measurement in patients with heart failure. <i>European Journal of Heart Failure</i> , 2004 , 6, 85-93	12.3	11
60	Definition of physiological hypertrophy in ultramarathon athletes. <i>Journal of the American College of Cardiology</i> , 2004 , 44, 469; author reply 471	15.1	11
59	CYP1A1 MSPI (T6235C) gene polymorphism is associated with mortality in acute coronary syndrome patients. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2010 , 37, 193-8	3	10
58	Serum IGF-I levels are similar in Samoan, MBri and European populations despite differences in body composition. <i>Growth Hormone and IGF Research</i> , 2006 , 16, 57-60	2	10
57	Early detection and significance of structural cardiovascular abnormalities in patients with Type 2 diabetes mellitus. <i>Expert Review of Cardiovascular Therapy</i> , 2008 , 6, 109-25	2.5	9
56	The effect of acute hyperglycaemia on brachial artery flow mediated dilatation in normal volunteers. <i>Australian and New Zealand Journal of Medicine</i> , 2000 , 30, 344-50		9
55	Selection bias in clinical research when subjects are excluded because of failure to estimate left ventricular mass by echocardiography. <i>Journal of the American Society of Echocardiography</i> , 1998 , 11, 1050-5	5.8	9
54	Myocardial tissue characterisation using echocardiographic deformation imaging. <i>Cardiovascular Ultrasound</i> , 2019 , 17, 27	2.4	9

(2005-2015)

53	Genetic markers of repolarization and arrhythmic events after acute coronary syndromes. <i>American Heart Journal</i> , 2015 , 169, 579-86.e3	4.9	8
52	Plasma levels of soluble VEGF receptor isoforms, circulating pterins and VEGF system SNPs as prognostic biomarkers in patients with acute coronary syndromes. <i>BMC Cardiovascular Disorders</i> , 2018 , 18, 169	2.3	8
51	Community screening for cardiovascular risk factors and levels of treatment in a rural MBri cohort. <i>Australian and New Zealand Journal of Public Health</i> , 2011 , 35, 517-23	2.3	8
50	Does rhythm matter? The prognostic importance of atrial fibrillation in heart failure. <i>Heart Lung and Circulation</i> , 2006 , 15, 353-7	1.8	8
49	Hyperendemic rheumatic heart disease in a remote Australian town identified by echocardiographic screening. <i>Medical Journal of Australia</i> , 2020 , 213, 118-123	4	8
48	Relation of Left Atrial Volumes in Patients With Myocardial Infarction to Left Ventricular Filling Pressures and Outcomes. <i>American Journal of Cardiology</i> , 2019 , 124, 325-333	3	7
47	Genetic polymorphism rs6922269 in the MTHFD1L gene is associated with survival and baseline active vitamin B12 levels in post-acute coronary syndromes patients. <i>PLoS ONE</i> , 2014 , 9, e89029	3.7	7
46	Convalescent troponin and cardiovascular death following acute coronary syndrome. <i>Heart</i> , 2019 , 105, 1717-1724	5.1	7
45	Arterial baroreceptor reflex control of renal sympathetic nerve activity following chronic myocardial infarction in male, female, and ovariectomized female rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2015 , 309, R169-78	3.2	6
44	Echocardiographic predictors of all-cause mortality in patients with left ventricular ejection fraction >35%: Value of guideline based assessment of diastolic dysfunction. <i>IJC Heart and Vasculature</i> , 2019 , 24, 100407	2.4	5
43	Understanding differences in results from literature-based and individual patient meta-analyses: an example from meta-analyses of observational data. <i>International Journal of Cardiology</i> , 2011 , 148, 209-1	3 ^{.2}	5
42	Role of echocardiographic left ventricular mass and carotid intima-media thickness in the cardiovascular risk assessment of asymptomatic patients with type 2 diabetes mellitus. <i>Internal Medicine Journal</i> , 2011 , 41, 391-8	1.6	5
41	Plasma N-terminal protype-B natriuretic peptide and restrictive mitral flow to risk-stratify patients with stage B heart failure. <i>Clinical Cardiology</i> , 2009 , 32, 711-7	3.3	5
40	Diastolic Filling Response To Submaximal Exercise In Trained And Untrained Subjects. <i>Medicine and Science in Sports and Exercise</i> , 2006 , 38, S115	1.2	5
39	Correlation between epicardial adipose tissue and body mass index in New Zealand ethnic populations. <i>New Zealand Medical Journal</i> , 2020 , 133, 22-32	0.8	5
38	The relationship between BNP and E/Ea in patients hospitalized with acute heart failure. <i>International Journal of Cardiology</i> , 2008 , 125, 280-2	3.2	4
37	Transmitral flow patterns and the presence of chronic kidney disease provide independent and incremental prognostic information in patients with heart failure and systolic dysfunction. <i>Journal of the American Society of Echocardiography</i> , 2007 , 20, 989-97	5.8	4
36	Role of echocardiography in the contemporary management of chronic heart failure. <i>Expert Review of Cardiovascular Therapy</i> , 2005 , 3, 51-70	2.5	4

35	A comparison of the effects of indexation on standard echocardiographic measurements of the left heart in a healthy multi-racial population. <i>International Journal of Cardiovascular Imaging</i> , 2014 , 30, 749-58	3 5	3
34	Higher prevalence of left ventricular hypertrophy in two MBri cohorts: findings from the Hauora Manawa/Community Heart Study. <i>Australian and New Zealand Journal of Public Health</i> , 2015 , 39, 26-31	.3	3
33	Changes in tissue-Doppler echocardiographic assessment of left ventricular filling during NT-proBNP guided heart failure treatment titration: a pilot study. <i>Heart Lung and Circulation</i> , 2009 , 18, 38-44	.8	3
32	Survey of clinical echocardiography in New Zealand (SCANZ). <i>New Zealand Medical Journal</i> , 2008 , 121, 34-44	.8	3
31	The impact of beta-blockade on right ventricular function in mitral regurgitation. <i>Heart Lung and Circulation</i> , 2014 , 23, 378-80	.8	2
30	Branding, Ingredients and Nutrition Information: Consumer Liking of a Healthier Snack. <i>Journal of Food Research</i> , 2015 , 4, 64	-3	2
29	Association between endothelin type A receptor haplotypes and mortality in coronary heart disease. <i>Personalized Medicine</i> , 2012 , 9, 341-349	.2	2
28	Echocardiographic Left Atrial Volumes are Optimally Indexed to Lean Body Mass to Adjust for Differences in Body Size. <i>Heart Lung and Circulation</i> , 2008 , 17, S45	.8	2
27	Quantitative evaluation of regional endocardial visualisation with second harmonic imaging and contrast left ventricular opacification in heart failure patients. <i>European Journal of Echocardiography</i> , 2005 , 6, 134-43		2
26	Hemodynamic Validation of the E/eQatio as a Measure of Left Ventricular Filling Pressure in Patients With Non-ST Elevation Myocardial Infarction. <i>American Journal of Cardiology</i> , 2020 , 125, 507-512		2
25	The RECARDINA Study protocol: diagnostic utility of ultra-abbreviated echocardiographic protocol for handheld machines used by non-experts to detect rheumatic heart disease. <i>BMJ Open</i> , 2020 , 10, e03 ⁷ 6	609	1
24	Prognostic implications of left ventricular dilation in patients with nonischemic heart failure: interactions with restrictive filling pattern and mitral regurgitation. <i>Congestive Heart Failure</i> , 2012 , 18, 198-204		1
23	Loose tobacco, ethnicity, income and rurality. <i>Australian and New Zealand Journal of Public Health</i> , 2012 , 36, 291-2	.3	1
22	Which cardiovascular risk factors are associated with cardiovascular disease and predict future events in advanced age in New Zealand?. <i>Australasian Journal on Ageing</i> , 2014 , 33, 14-21	.5	1
21	Prediction of ACC/AHA Stage B heart failure by clinical and neurohormonal profiling among patients in the community. <i>Journal of Cardiac Failure</i> , 2010 , 16, 957-63	.3	1
20	Coming of age: affiliate member profile and participation in the annual scientific meeting of the cardiac society of Australia and New Zealand. <i>Heart Lung and Circulation</i> , 2007 , 16, 447-51	.8	1
19	Mechanisms of benefit of sustained weight reduction in morbid obesity: beyond reduction in conventional cardiovascular risk factors. <i>Journal of Hypertension</i> , 2007 , 25, 295-7	.9	1
18	Snack bar compositions and their acute glycaemic and satiety effects. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2017 , 26, 624-629		1

LIST OF PUBLICATIONS

17	Effects of a healthier snack on snacking habits and glycated Hb (HbA1c): a 6-week intervention study. <i>British Journal of Nutrition</i> , 2016 , 116, 2169-2174	3.6	1	
16	Echo and BNP serial assessment in ambulatory heart failure care: Data on loop diuretic use and renal function. <i>Data in Brief</i> , 2016 , 9, 1074-1076	1.2	1	
15	Single-View Echocardiography by Nonexpert Practitioners to Detect Rheumatic Heart Disease: A Prospective Study of Diagnostic Accuracy. <i>Circulation: Cardiovascular Imaging</i> , 2021 , 14, e011790	3.9	1	
14	The 21st Century Echocardiography Laboratory in Australia and New Zealand: Rapid Evolution of Training and Workforce, Practice and Technology. <i>Heart Lung and Circulation</i> , 2019 , 28, 1421-1426	1.8	O	
13	Review: Detection of patient foramen ovale using transcranial Doppler or standard echocardiography. <i>Australasian Journal of Ultrasound in Medicine</i> , 2020 , 23, 210-219	0.6	O	
12	Brain natriuretic peptide in the contemporary management of congestive heart failure. <i>Expert Review of Cardiovascular Therapy</i> , 2005 , 3, 71-84	2.5	O	
11	Sex Disparity in Cardiovascular Disease Outcomes: Do Our Current Echocardiographic Reference Ranges Measure Up?. <i>Heart Lung and Circulation</i> , 2021 , 30, e1-e5	1.8	O	
10	Predictors of quality of life after revascularization for ischemic heart disease: A systematic review. <i>Health Sciences Review</i> , 2022 , 2, 100017		O	
9	The development and feasibility of a composite score of echocardiographic indices that may stratify outcome in patients with diabetes mellitus. <i>International Journal of Cardiology</i> , 2015 , 182, 244-	.9 ^{3.2}		
8	Long-term outcomes in patients with restrictive filling following ST-segment elevation myocardial infarction. <i>Internal Medicine Journal</i> , 2014 , 44, 291-4	1.6		
7	The cardiac sonography workforce in New Zealand. <i>Australasian Journal of Ultrasound in Medicine</i> , 2013 , 16, 77-85	0.6		
6	The Effect of Age and Endurance Training on Exercising Diastolic Function in Healthy Men. <i>Medicine and Science in Sports and Exercise</i> , 2010 , 42, 5	1.2		
5	Detection of pathologic or physiologic left ventricular remodeling in athletes. <i>Journal of the American College of Cardiology</i> , 2005 , 45, 1731; author reply 1731	15.1		
4	Endothelium-dependent dilatation in patients with type 2 diabetes. <i>American Journal of Cardiology</i> , 2002 , 90, 446	3		
3	Integrated management program does not reduce death, but may improve quality of life for people with chronic heart failure. <i>Evidence-based Cardiovascular Medicine</i> , 2002 , 6, 123-124			
2	Hemodynamic and Prognostic Validation of Novel Combined Algorithm to Assess Diastolic Function and Filling Pressures. <i>JACC: Cardiovascular Imaging</i> , 2020 , 13, 2275-2276	8.4		
1	A new approach to assessment of the left ventricle. <i>MethodsX</i> , 2016 , 3, 274-8	1.9		