Gillian A Whalley

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/5130747/gillian-a-whalley-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

142 papers

5,446 citations

36 h-index

g-index

166 ext. papers

6,297 ext. citations

avg, IF

5.61 L-index

#	Paper	IF	Citations
142	Predictors of quality of life after revascularization for ischemic heart disease: A systematic review. Health Sciences Review, 2022 , 2, 100017		O
141	Sex Disparity in Cardiovascular Disease Outcomes: Do Our Current Echocardiographic Reference Ranges Measure Up?. <i>Heart Lung and Circulation</i> , 2021 , 30, e1-e5	1.8	О
140	Single-View Echocardiography by Nonexpert Practitioners to Detect Rheumatic Heart Disease: A Prospective Study of Diagnostic Accuracy. <i>Circulation: Cardiovascular Imaging</i> , 2021 , 14, e011790	3.9	1
139	Review: Detection of patient foramen ovale using transcranial Doppler or standard echocardiography. <i>Australasian Journal of Ultrasound in Medicine</i> , 2020 , 23, 210-219	0.6	O
138	The RECARDINA Study protocol: diagnostic utility of ultra-abbreviated echocardiographic protocol for handheld machines used by non-experts to detect rheumatic heart disease. <i>BMJ Open</i> , 2020 , 10, e03	3 7 609	1
137	Hemodynamic Validation of the E/eQRatio as a Measure of Left Ventricular Filling Pressure in Patients With Non-ST Elevation Myocardial Infarction. <i>American Journal of Cardiology</i> , 2020 , 125, 507-57	12	2
136	Hyperendemic rheumatic heart disease in a remote Australian town identified by echocardiographic screening. <i>Medical Journal of Australia</i> , 2020 , 213, 118-123	4	8
135	Hemodynamic and Prognostic Validation of Novel Combined Algorithm to Assess Diastolic Function and Filling Pressures. <i>JACC: Cardiovascular Imaging</i> , 2020 , 13, 2275-2276	8.4	
134	Correlation between epicardial adipose tissue and body mass index in New Zealand ethnic populations. <i>New Zealand Medical Journal</i> , 2020 , 133, 22-32	0.8	5
133	Relation of Left Atrial Volumes in Patients With Myocardial Infarction to Left Ventricular Filling Pressures and Outcomes. <i>American Journal of Cardiology</i> , 2019 , 124, 325-333	3	7
132	New Diastology Guidelines: Evolution, Validation and Impact on Clinical Practice. <i>Heart Lung and Circulation</i> , 2019 , 28, 1411-1420	1.8	11
131	The 21st Century Echocardiography Laboratory in Australia and New Zealand: Rapid Evolution of Training and Workforce, Practice and Technology. <i>Heart Lung and Circulation</i> , 2019 , 28, 1421-1426	1.8	O
130	Echocardiographic predictors of all-cause mortality in patients with left ventricular ejection fraction >35%: Value of guideline based assessment of diastolic dysfunction. <i>IJC Heart and Vasculature</i> , 2019 , 24, 100407	2.4	5
129	Convalescent troponin and cardiovascular death following acute coronary syndrome. <i>Heart</i> , 2019 , 105, 1717-1724	5.1	7
128	Myocardial tissue characterisation using echocardiographic deformation imaging. <i>Cardiovascular Ultrasound</i> , 2019 , 17, 27	2.4	9
127	Diastolic Dysfunction Assessed Using Contemporary Guidelines and Prognosis Following Myocardial Infarction. <i>Journal of the American Society of Echocardiography</i> , 2018 , 31, 1127-1136	5.8	27
126	Plasma levels of soluble VEGF receptor isoforms, circulating pterins and VEGF system SNPs as prognostic biomarkers in patients with acute coronary syndromes. <i>BMC Cardiovascular Disorders</i> , 2018 , 18, 169	2.3	8

125	C-Type Natriuretic Peptides in Coronary Disease. Clinical Chemistry, 2017, 63, 316-324	5.5	14
124	Snack bar compositions and their acute glycaemic and satiety effects. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2017 , 26, 624-629	1	1
123	Prognostic significance of anaemia in patients with heart failure with preserved and reduced ejection fraction: results from the MAGGIC individual patient data meta-analysis. <i>QJM - Monthly Journal of the Association of Physicians</i> , 2016 , 109, 377-382	2.7	24
122	Bias associated with left ventricular quantification by multimodality imaging: a systematic review and meta-analysis. <i>Open Heart</i> , 2016 , 3, e000388	3	31
121	Assessment and impact of diastolic function by echocardiography in elderly patients. <i>Journal of Geriatric Cardiology</i> , 2016 , 13, 252-60	1.7	12
120	Effects of a healthier snack on snacking habits and glycated Hb (HbA1c): a 6-week intervention study. <i>British Journal of Nutrition</i> , 2016 , 116, 2169-2174	3.6	1
119	Echo and BNP serial assessment in ambulatory heart failure care: Data on loop diuretic use and renal function. <i>Data in Brief</i> , 2016 , 9, 1074-1076	1.2	1
118	A new approach to assessment of the left ventricle. <i>MethodsX</i> , 2016 , 3, 274-8	1.9	
117	The Validity of Left Ventricular Mass as a Surrogate End Point for All-Cause and Cardiovascular Mortality Outcomes in People With CKD: A Systematic Review and Meta-analysis. <i>American Journal of Kidney Diseases</i> , 2016 , 68, 554-563	7.4	36
116	Echo and natriuretic peptide guided therapy improves outcome and reduces worsening renal function in systolic heart failure: An observational study of 1137 outpatients. <i>International Journal of Cardiology</i> , 2016 , 224, 416-423	3.2	20
115	Genetic markers of repolarization and arrhythmic events after acute coronary syndromes. <i>American Heart Journal</i> , 2015 , 169, 579-86.e3	4.9	8
114	Arterial baroreceptor reflex control of renal sympathetic nerve activity following chronic myocardial infarction in male, female, and ovariectomized female rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2015 , 309, R169-78	3.2	6
113	Left ventricular geometry and all-cause mortality in advanced age. <i>Heart Lung and Circulation</i> , 2015 , 24, 32-9	1.8	12
112	Higher prevalence of left ventricular hypertrophy in two MBri cohorts: findings from the Hauora Manawa/Community Heart Study. <i>Australian and New Zealand Journal of Public Health</i> , 2015 , 39, 26-31	2.3	3
111	Is heart rate a risk marker in patients with chronic heart failure and concomitant atrial fibrillation? Results from the MAGGIC meta-analysis. <i>European Journal of Heart Failure</i> , 2015 , 17, 1182-91	12.3	37
110	Branding, Ingredients and Nutrition Information: Consumer Liking of a Healthier Snack. <i>Journal of Food Research</i> , 2015 , 4, 64	1.3	2
109	Differing prognostic value of pulse pressure in patients with heart failure with reduced or preserved ejection fraction: results from the MAGGIC individual patient meta-analysis. <i>European Heart Journal</i> , 2015 , 36, 1106-14	9.5	37
108	Ethnic-Specific Normative Reference Values for Echocardiographic LAland LV Size, LV Mass, and Systolic Function: The EchoNoRMAL Study. <i>JACC: Cardiovascular Imaging</i> , 2015 , 8, 656-65	8.4	125

107	The prognostic impact of diastolic dysfunction in patients with chronic heart failure and post-acute myocardial infarction: Can age-stratified E/A ratio alone predict survival?. <i>International Journal of Cardiology</i> , 2015 , 181, 362-8	3.2	11
106	The development and feasibility of a composite score of echocardiographic indices that may stratify outcome in patients with diabetes mellitus. <i>International Journal of Cardiology</i> , 2015 , 182, 244	-9 ^{3.2}	
105	Circulating miR-323-3p and miR-652: candidate markers for the presence and progression of acute coronary syndromes. <i>International Journal of Cardiology</i> , 2014 , 176, 375-85	3.2	36
104	A comparison of the effects of indexation on standard echocardiographic measurements of the left heart in a healthy multi-racial population. <i>International Journal of Cardiovascular Imaging</i> , 2014 , 30, 74	9- 5 8 ⁵	3
103	The impact of beta-blockade on right ventricular function in mitral regurgitation. <i>Heart Lung and Circulation</i> , 2014 , 23, 378-80	1.8	2
102	Heart failure in younger patients: the Meta-analysis Global Group in Chronic Heart Failure (MAGGIC). <i>European Heart Journal</i> , 2014 , 35, 2714-21	9.5	50
101	The obesity paradox in heart failure patients with preserved versus reduced ejection fraction: a meta-analysis of individual patient data. <i>International Journal of Obesity</i> , 2014 , 38, 1110-4	5.5	123
100	Long-term outcomes in patients with restrictive filling following ST-segment elevation myocardial infarction. <i>Internal Medicine Journal</i> , 2014 , 44, 291-4	1.6	
99	Which cardiovascular risk factors are associated with cardiovascular disease and predict future events in advanced age in New Zealand?. <i>Australasian Journal on Ageing</i> , 2014 , 33, 14-21	1.5	1
98	Genetic polymorphism rs6922269 in the MTHFD1L gene is associated with survival and baseline active vitamin B12 levels in post-acute coronary syndromes patients. <i>PLoS ONE</i> , 2014 , 9, e89029	3.7	7
97	Known and missing left ventricular ejection fraction and survival in patients with heart failure: a MAGGIC meta-analysis report. <i>European Journal of Heart Failure</i> , 2013 , 15, 1220-7	12.3	22
96	Effect of early initiation of dialysis on cardiac structure and function: results from the echo substudy of the IDEAL trial. <i>American Journal of Kidney Diseases</i> , 2013 , 61, 262-70	7.4	38
95	Redefining normal reference ranges for echocardiography: a major new individual person data meta-analysis. <i>European Heart Journal Cardiovascular Imaging</i> , 2013 , 14, 347-8	4.1	18
94	Predicting survival in heart failure: a risk score based on 39 372 patients from 30 studies. <i>European Heart Journal</i> , 2013 , 34, 1404-13	9.5	595
93	Hyperuricaemia and gout in New Zealand rural and urban MBri and non-MBri communities. <i>Internal Medicine Journal</i> , 2013 , 43, 678-84	1.6	22
92	Genetic variation in the renin-angiotensin-aldosterone system is associated with cardiovascular risk factors and early mortality in established coronary heart disease. <i>Journal of Human Hypertension</i> , 2013 , 27, 237-44	2.6	19
91	The cardiac sonography workforce in New Zealand. <i>Australasian Journal of Ultrasound in Medicine</i> , 2013 , 16, 77-85	0.6	
90	Chronic measurement of left ventricular pressure in freely moving rats. <i>Journal of Applied Physiology</i> , 2013 , 115, 1672-82	3.7	12

89	Heart failure with preserved ejection fraction. Journal of Geriatric Cardiology, 2013, 10, 369-76	1.7	14
88	The larger exercise stroke volume in endurance-trained men does not result from increased left ventricular early or late inflow or tissue velocities. <i>Acta Physiologica</i> , 2012 , 205, 520-31	5.6	16
87	Prognostic implications of left ventricular dilation in patients with nonischemic heart failure: interactions with restrictive filling pattern and mitral regurgitation. <i>Congestive Heart Failure</i> , 2012 , 18, 198-204		1
86	Loose tobacco, ethnicity, income and rurality. <i>Australian and New Zealand Journal of Public Health</i> , 2012 , 36, 291-2	2.3	1
85	Relationship of serum sodium concentration to mortality in a wide spectrum of heart failure patients with preserved and with reduced ejection fraction: an individual patient data meta-analysis(I): Meta-Analysis Global Group in Chronic heart failure (MAGGIC). European Journal of	12.3	74
84	Heart Failure, 2012, 14, 1139-46 Gender and survival in patients with heart failure: interactions with diabetes and aetiology. Results from the MAGGIC individual patient meta-analysis. European Journal of Heart Failure, 2012, 14, 473-9	12.3	119
83	KCNE5 polymorphism rs697829 is associated with QT interval and survival in acute coronary syndromes patients. <i>Journal of Cardiovascular Electrophysiology</i> , 2012 , 23, 319-24	2.7	11
82	Progression of myocardial remodeling and mechanical dysfunction in the spontaneously hypertensive rat. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2012 , 303, H1353-65	5 ^{5.2}	45
81	Renal dysfunction in patients with heart failure with preserved versus reduced ejection fraction: impact of the new Chronic Kidney Disease-Epidemiology Collaboration Group formula. <i>Circulation: Heart Failure</i> , 2012 , 5, 309-14	7.6	112
80	Geographic variation in left ventricular mass and mass index: a systematic review. <i>Journal of Human Hypertension</i> , 2012 , 26, 420-9	2.6	22
79	The survival of patients with heart failure with preserved or reduced left ventricular ejection fraction: an individual patient data meta-analysis. <i>European Heart Journal</i> , 2012 , 33, 1750-7	9.5	508
78	Association between endothelin type A receptor haplotypes and mortality in coronary heart disease. <i>Personalized Medicine</i> , 2012 , 9, 341-349	2.2	2
77	A cohort study comparing cardiovascular risk factors in rural Maori, urban Maori and non-Maori communities in New Zealand. <i>BMJ Open</i> , 2012 , 2,	3	18
76	Understanding differences in results from literature-based and individual patient meta-analyses: an example from meta-analyses of observational data. <i>International Journal of Cardiology</i> , 2011 , 148, 209-	13 ^{.2}	5
75	Association of genetic variation in the natriuretic peptide system with cardiovascular outcomes. Journal of Molecular and Cellular Cardiology, 2011 , 50, 695-701	5.8	49
74	Effects of perindopril-indapamide on left ventricular diastolic function and mass in patients with type 2 diabetes: the ADVANCE Echocardiography Substudy. <i>Journal of Hypertension</i> , 2011 , 29, 1439-47	1.9	17
73	Role of echocardiographic left ventricular mass and carotid intima-media thickness in the cardiovascular risk assessment of asymptomatic patients with type 2 diabetes mellitus. <i>Internal Medicine Journal</i> , 2011 , 41, 391-8	1.6	5
7 2	A Kaupapa MBri approach to a community cohort study of heart disease in New Zealand. Australian and New Zealand Journal of Public Health, 2011 , 35, 249-55	2.3	18

71	Community screening for cardiovascular risk factors and levels of treatment in a rural MBri cohort. <i>Australian and New Zealand Journal of Public Health</i> , 2011 , 35, 517-23	2.3	8
70	Screening for left ventricular hypertrophy in patients with type 2 diabetes mellitus in the community. <i>Cardiovascular Diabetology</i> , 2011 , 10, 29	8.7	41
69	Genomic risk variants at 1p13.3, 1q41, and 3q22.3 are associated with subsequent cardiovascular outcomes in healthy controls and in established coronary artery disease. <i>Circulation: Cardiovascular Genetics</i> , 2011 , 4, 636-46		26
68	CYP1A1 MSPI (T6235C) gene polymorphism is associated with mortality in acute coronary syndrome patients. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2010 , 37, 193-8	3	10
67	Left ventricular systolic and diastolic function assessed by tissue Doppler imaging and outcome in asymptomatic aortic stenosis. <i>European Heart Journal</i> , 2010 , 31, 2216-22	9.5	60
66	A community-based model of care improves blood pressure control and delays progression of proteinuria, left ventricular hypertrophy and diastolic dysfunction in Maori and Pacific patients with type 2 diabetes and chronic kidney disease: a randomized controlled trial. <i>Nephrology Dialysis</i>	4.3	38
65	A common variant at chromosome 9P21.3 is associated with age of onset of coronary disease but not subsequent mortality. <i>Circulation: Cardiovascular Genetics</i> , 2010 , 3, 286-93		37
64	Prediction of ACC/AHA Stage B heart failure by clinical and neurohormonal profiling among patients in the community. <i>Journal of Cardiac Failure</i> , 2010 , 16, 957-63	3.3	1
63	Atrial fibrillation and the risk of death in patients with heart failure: a literature-based meta-analysis. <i>Internal Medicine Journal</i> , 2010 , 40, 347-56	1.6	32
62	Understanding changing patterns of survival and hospitalization for heart failure over two decades in New Zealand: utility of Q ays alive and out of hospital Q rom epidemiological data. <i>European Journal of Heart Failure</i> , 2010 , 12, 462-8	12.3	52
61	The Effect of Age and Endurance Training on Exercising Diastolic Function in Healthy Men. <i>Medicine and Science in Sports and Exercise</i> , 2010 , 42, 5	1.2	
60	The prognostic significance of heart failure with preserved left ventricular ejection fraction: a literature-based meta-analysis. <i>European Journal of Heart Failure</i> , 2009 , 11, 855-62	12.3	99
59	Independent relationship of left atrial size and mortality in patients with heart failure: an individual patient meta-analysis of longitudinal data (MeRGE Heart Failure). <i>European Journal of Heart Failure</i> , 2009 , 11, 929-36	12.3	109
58	Structural and functional cardiac abnormalities in adolescent girls with poorly controlled type 2 diabetes. <i>Diabetes Care</i> , 2009 , 32, 883-8	14.6	28
57	Plasma N-terminal protype-B natriuretic peptide and restrictive mitral flow to risk-stratify patients with stage B heart failure. <i>Clinical Cardiology</i> , 2009 , 32, 711-7	3.3	5
56	Changes in tissue-Doppler echocardiographic assessment of left ventricular filling during NT-proBNP guided heart failure treatment titration: a pilot study. <i>Heart Lung and Circulation</i> , 2009 , 18, 38-44	1.8	3
55	Pseudonormal mitral filling is associated with similarly poor prognosis as restrictive filling in patients with heart failure and coronary heart disease: a systematic review and meta-analysis of prospective studies. <i>Journal of the American Society of Echocardiography</i> , 2009 , 22, 494-8	5.8	25
54	Echocardiographic Left Atrial Volumes are Optimally Indexed to Lean Body Mass to Adjust for Differences in Body Size. <i>Heart Lung and Circulation</i> , 2008 , 17, S45	1.8	2

(2006-2008)

53	The relationship between BNP and E/Ea in patients hospitalized with acute heart failure. <i>International Journal of Cardiology</i> , 2008 , 125, 280-2	3.2	4
52	Increased B-type natriuretic peptide is associated with an abnormal blood pressure response to exercise in asymptomatic aortic stenosis. <i>International Journal of Cardiology</i> , 2008 , 127, 313-20	3.2	18
51	A randomized trial of the aldosterone-receptor antagonist eplerenone in asymptomatic moderate-severe aortic stenosis. <i>American Heart Journal</i> , 2008 , 156, 348-55	4.9	29
50	Angiotensin-converting enzyme 2 A1075G polymorphism is associated with survival in an acute coronary syndromes cohort. <i>American Heart Journal</i> , 2008 , 156, 752-8	4.9	19
49	Early detection and significance of structural cardiovascular abnormalities in patients with Type 2 diabetes mellitus. <i>Expert Review of Cardiovascular Therapy</i> , 2008 , 6, 109-25	2.5	9
48	Elevated B-type natriuretic peptide despite normal left ventricular function on rest and exercise stress echocardiography in mitral regurgitation. <i>European Heart Journal</i> , 2008 , 29, 363-70	9.5	32
47	Independence of restrictive filling pattern and LV ejection fraction with mortality in heart failure: an individual patient meta-analysis. <i>European Journal of Heart Failure</i> , 2008 , 10, 786-92	12.3	61
46	Independent prognostic importance of a restrictive left ventricular filling pattern after myocardial infarction: an individual patient meta-analysis: Meta-Analysis Research Group in Echocardiography acute myocardial infarction. <i>Circulation</i> , 2008 , 117, 2591-8	16.7	123
45	Prognostic role of echocardiography and brain natriuretic peptide in symptomatic breathless patients in the community. <i>European Heart Journal</i> , 2008 , 29, 509-16	9.5	28
44	Survey of clinical echocardiography in New Zealand (SCANZ). <i>New Zealand Medical Journal</i> , 2008 , 121, 34-44	0.8	3
43	Transmitral flow patterns and the presence of chronic kidney disease provide independent and incremental prognostic information in patients with heart failure and systolic dysfunction. <i>Journal of the American Society of Echocardiography</i> , 2007 , 20, 989-97	5.8	4
42	Coming of age: affiliate member profile and participation in the annual scientific meeting of the cardiac society of Australia and New Zealand. <i>Heart Lung and Circulation</i> , 2007 , 16, 447-51	1.8	1
41	Restrictive filling pattern is a powerful predictor of heart failure events postacute myocardial infarction and in established heart failure: a literature-based meta-analysis. <i>Journal of Cardiac Failure</i> , 2007 , 13, 346-52	3.3	36
40	Longitudinal left ventricular contractile dysfunction after exercise in aortic stenosis. <i>Heart</i> , 2007 , 93, 732-8	5.1	45
39	Mechanisms of benefit of sustained weight reduction in morbid obesity: beyond reduction in conventional cardiovascular risk factors. <i>Journal of Hypertension</i> , 2007 , 25, 295-7	1.9	1
38	The prognostic significance of restrictive diastolic filling associated with heart failure: a meta-analysis. <i>International Journal of Cardiology</i> , 2007 , 116, 70-7	3.2	31
37	Individual patient meta-analyses of restrictive diastolic filling pattern and mortality in patients post acute myocardial infarction and in patients with chronic heart failure. <i>International Journal of Cardiology</i> , 2007 , 122, 207-15	3.2	17
36	Does rhythm matter? The prognostic importance of atrial fibrillation in heart failure. <i>Heart Lung and Circulation</i> , 2006 , 15, 353-7	1.8	8

35	Restrictive diastolic filling predicts death after acute myocardial infarction: systematic review and meta-analysis of prospective studies. <i>Heart</i> , 2006 , 92, 1588-94	5.1	32
34	Serum IGF-I levels are similar in Samoan, MBri and European populations despite differences in body composition. <i>Growth Hormone and IGF Research</i> , 2006 , 16, 57-60	2	10
33	The effect of type 2 diabetes on diastolic function. <i>Medicine and Science in Sports and Exercise</i> , 2006 , 38, 1384-8	1.2	21
32	Diastolic Filling Response To Submaximal Exercise In Trained And Untrained Subjects. <i>Medicine and Science in Sports and Exercise</i> , 2006 , 38, S115	1.2	5
31	Quantitative evaluation of regional endocardial visualisation with second harmonic imaging and contrast left ventricular opacification in heart failure patients. <i>European Journal of Echocardiography</i> , 2005 , 6, 134-43		2
30	Comparison of different methods for detection of diastolic filling abnormalities. <i>Journal of the American Society of Echocardiography</i> , 2005 , 18, 710-7	5.8	15
29	Detection of pathologic or physiologic left ventricular remodeling in athletes. <i>Journal of the American College of Cardiology</i> , 2005 , 45, 1731; author reply 1731	15.1	
28	Brain natriuretic peptide in the contemporary management of congestive heart failure. <i>Expert Review of Cardiovascular Therapy</i> , 2005 , 3, 71-84	2.5	O
27	Role of echocardiography in the contemporary management of chronic heart failure. <i>Expert Review of Cardiovascular Therapy</i> , 2005 , 3, 51-70	2.5	4
26	Effects of carvedilol on left ventricular remodeling after acute myocardial infarction: the CAPRICORN Echo Substudy. <i>Circulation</i> , 2004 , 109, 201-6	16.7	244
26 25		16.7	244
	CAPRICORN Echo Substudy. <i>Circulation</i> , 2004 , 109, 201-6 Effect of tissue harmonic imaging and contrast upon between observer and test-retest reproducibility of left ventricular ejection fraction measurement in patients with heart failure.	,	
25	CAPRICORN Echo Substudy. <i>Circulation</i> , 2004 , 109, 201-6 Effect of tissue harmonic imaging and contrast upon between observer and test-retest reproducibility of left ventricular ejection fraction measurement in patients with heart failure. <i>European Journal of Heart Failure</i> , 2004 , 6, 85-93 Three-dimensional assessment of left ventricular systolic strain in patients with type 2 diabetes mellitus, diastolic dysfunction, and normal ejection fraction. <i>American Journal of Cardiology</i> , 2004 ,	12.3	11
25 24	CAPRICORN Echo Substudy. <i>Circulation</i> , 2004 , 109, 201-6 Effect of tissue harmonic imaging and contrast upon between observer and test-retest reproducibility of left ventricular ejection fraction measurement in patients with heart failure. <i>European Journal of Heart Failure</i> , 2004 , 6, 85-93 Three-dimensional assessment of left ventricular systolic strain in patients with type 2 diabetes mellitus, diastolic dysfunction, and normal ejection fraction. <i>American Journal of Cardiology</i> , 2004 , 94, 1391-5 Definition of physiological hypertrophy in ultramarathon athletes. <i>Journal of the American College</i>	12.3 3 15.1	11 99
25 24 23	Effect of tissue harmonic imaging and contrast upon between observer and test-retest reproducibility of left ventricular ejection fraction measurement in patients with heart failure. European Journal of Heart Failure, 2004, 6, 85-93 Three-dimensional assessment of left ventricular systolic strain in patients with type 2 diabetes mellitus, diastolic dysfunction, and normal ejection fraction. American Journal of Cardiology, 2004, 94, 1391-5 Definition of physiological hypertrophy in ultramarathon athletes. Journal of the American College of Cardiology, 2004, 44, 469; author reply 471 Association of fat-free mass and training status with left ventricular size and mass in	12.3 3 15.1	11 99 11
25 24 23 22	Effect of tissue harmonic imaging and contrast upon between observer and test-retest reproducibility of left ventricular ejection fraction measurement in patients with heart failure. European Journal of Heart Failure, 2004, 6, 85-93 Three-dimensional assessment of left ventricular systolic strain in patients with type 2 diabetes mellitus, diastolic dysfunction, and normal ejection fraction. American Journal of Cardiology, 2004, 94, 1391-5 Definition of physiological hypertrophy in ultramarathon athletes. Journal of the American College of Cardiology, 2004, 44, 469; author reply 471 Association of fat-free mass and training status with left ventricular size and mass in endurance-trained athletes. Journal of the American College of Cardiology, 2004, 44, 892-6 Uptake of self-management strategies in a heart failure management programme. European	12.3 3 15.1	11 99 11 36
25 24 23 22 21	Effect of tissue harmonic imaging and contrast upon between observer and test-retest reproducibility of left ventricular ejection fraction measurement in patients with heart failure. European Journal of Heart Failure, 2004, 6, 85-93 Three-dimensional assessment of left ventricular systolic strain in patients with type 2 diabetes mellitus, diastolic dysfunction, and normal ejection fraction. American Journal of Cardiology, 2004, 94, 1391-5 Definition of physiological hypertrophy in ultramarathon athletes. Journal of the American College of Cardiology, 2004, 44, 469; author reply 471 Association of fat-free mass and training status with left ventricular size and mass in endurance-trained athletes. Journal of the American College of Cardiology, 2004, 44, 892-6 Uptake of self-management strategies in a heart failure management programme. European Journal of Heart Failure, 2003, 5, 371-80 Left ventricular diastolic filling and systolic function of young and older trained and untrained men.	12.3 3 15.1 15.1 12.3	111 999 111 36 98 45

LIST OF PUBLICATIONS

17	Randomized, controlled trial of integrated heart failure management: The Auckland Heart Failure Management Study. <i>European Heart Journal</i> , 2002 , 23, 139-46	9.5	180
16	Pseudonormal mitral filling pattern predicts hospital re-admission in patients with congestive heart failure. <i>Journal of the American College of Cardiology</i> , 2002 , 39, 1787-95	15.1	85
15	Integrated management program does not reduce death, but may improve quality of life for people with chronic heart failure. <i>Evidence-based Cardiovascular Medicine</i> , 2002 , 6, 123-124		
14	Comparison of ultrasound assessment of flow-mediated dilatation in the radial and brachial artery with upper and forearm cuff positions. <i>Clinical Physiology</i> , 2001 , 21, 9-14		50
13	Effects of improved glycaemic control on endothelial function in patients with type 2 diabetes. <i>Internal Medicine Journal</i> , 2001 , 31, 322-8	1.6	38
12	Pseudonormal diastolic filling unmasked with glyceryl trinitrate in patients with type 2 diabetes with poor metabolic control. <i>Diabetes Care</i> , 2001 , 24, 1307-8	14.6	16
11	The effect of acute hyperglycaemia on brachial artery flow mediated dilatation in normal volunteers. <i>Australian and New Zealand Journal of Medicine</i> , 2000 , 30, 344-50		9
10	Does a glass of red wine improve endothelial function?. <i>European Heart Journal</i> , 2000 , 21, 74-8	9.5	154
9	Effects of carvedilol on left ventricular regional wall motion in patients with heart failure caused by ischemic heart disease. Australia-New Zealand Heart Failure Research Collaborative Group. <i>Journal of Cardiac Failure</i> , 2000 , 6, 11-8	3.3	13
8	Left ventricular mass correlates with fat-free mass but not fat mass in adults. <i>Journal of Hypertension</i> , 1999 , 17, 569-74	1.9	25
7	Selection bias in clinical research when subjects are excluded because of failure to estimate left ventricular mass by echocardiography. <i>Journal of the American Society of Echocardiography</i> , 1998 , 11, 1050-5	5.8	9
6	Atherosclerosis and left ventricular hypertrophy: persisting problems in treated hypertensive patients. <i>Journal of Hypertension</i> , 1998 , 16, 1389-95	1.9	15
5	Left ventricular remodeling with carvedilol in patients with congestive heart failure due to ischemic heart disease. Australia-New Zealand Heart Failure Research Collaborative Group. <i>Journal of the American College of Cardiology</i> , 1997 , 29, 1060-6	15.1	329
4	Massive nitrogen loss in critical surgical illness: effect on cardiac mass and function. <i>Annals of Surgery</i> , 1997 , 226, 191-7	7.8	30
3	Effects of simvastatin and enalapril on serum lipoprotein concentrations and left ventricular mass in patients on dialysis. The Perfect Study Collaborative Group. <i>Journal of Nephrology</i> , 1997 , 10, 33-40	4.8	25
2	Doppler echocardiography and the early diagnosis of carditis in acute rheumatic fever. <i>Australian and New Zealand Journal of Medicine</i> , 1994 , 24, 530-5		44
1	Determinants of left ventricular hypertrophy and systolic dysfunction in chronic renal failure. <i>American Journal of Kidney Diseases</i> , 1994 , 24, 768-76	7.4	91