Hideki Yamamoto

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5131180/publications.pdf

Version: 2024-02-01

1040056 940533 16 319 9 16 citations h-index g-index papers 16 16 16 422 citing authors docs citations times ranked all docs

#	Article	IF	CITATIONS
1	Contribution of Invariant Natural Killer T Cells to the Clearance of Pseudomonas aeruginosa from Skin Wounds. International Journal of Molecular Sciences, 2021, 22, 3931.	4.1	5
2	Deficiency of lung-specific claudin-18 leads to aggravated infection with Cryptococcus deneoformans through dysregulation of the microenvironment in lungs. Scientific Reports, 2021, 11, 21110.	3.3	8
3	Effect of CARD9 Deficiency on Neutrophil-Mediated Host Defense against Pulmonary Infection with Streptococcus pneumoniae. Infection and Immunity, 2020, 89, .	2.2	7
4	Limited Role of Mincle in the Host Defense against Infection with Cryptococcus deneoformans. Infection and Immunity, 2020, 88, .	2.2	8
5	Production of IL-17A at Innate Immune Phase Leads to Decreased Th1 Immune Response and Attenuated Host Defense against Infection with <i>Cryptococcus deneoformans</i> . Journal of Immunology, 2020, 205, 686-698.	0.8	13
6	Novel Toll-Like Receptor 9 Agonist Derived from <i>Cryptococcus neoformans</i> Attenuates Allergic Inflammation Leading to Asthma Onset in Mice. International Archives of Allergy and Immunology, 2020, 181, 651-664.	2.1	2
7	Dectinâ€2â€mediated signaling triggered by the cell wall polysaccharides of <i>Cryptococcus neoformans</i> . Microbiology and Immunology, 2019, 63, 500-512.	1.4	10
8	Contribution of <scp>CARD</scp> 9â€mediated signalling to wound healing in skin. Experimental Dermatology, 2017, 26, 1097-1104.	2.9	10
9	Invariant NKT cells promote skin wound healing by preventing a prolonged neutrophilic inflammatory response. Wound Repair and Regeneration, 2017, 25, 805-815.	3.0	39
10	Cryptococcus neoformans Infection in Mice Lacking Type I Interferon Signaling Leads to Increased Fungal Clearance and IL-4-Dependent Mucin Production in the Lungs. PLoS ONE, 2015, 10, e0138291.	2.5	25
11	Dectin-2 Deficiency Promotes Th2 Response and Mucin Production in the Lungs after Pulmonary Infection with Cryptococcus neoformans. Infection and Immunity, 2015, 83, 671-681.	2.2	64
12	Involvement of Gr-1dull+ Cells in the Production of TNF- \hat{l}_{\pm} and IL-17 and Exacerbated Systemic Inflammatory Response Caused by Lipopolysaccharide. Inflammation, 2014, 37, 186-195.	3.8	4
13	Defect of CARD9 Leads to Impaired Accumulation of Gamma Interferon-Producing Memory Phenotype T Cells in Lungs and Increased Susceptibility to Pulmonary Infection with Cryptococcus neoformans. Infection and Immunity, 2014, 82, 1606-1615.	2.2	60
14	Toll-Like Receptor 9-Dependent Activation of Bone Marrow-Derived Dendritic Cells by <i>URA5 </i> PONA from Cryptococcus neoformans. Infection and Immunity, 2012, 80, 778-786.	2.2	23
15	Continuous hydrothermal synthesis of 3,4-dihydroxyhydrocinnamic acid-modified magnetite nanoparticles with stealth-functionality against immunological response. Journal of Materials Chemistry, 2012, 22, 9041.	6.7	33
16	Cryptococcus neoformanssuppresses the activation of bone marrow–derived dendritic cells stimulated with its own DNA, but not with DNA from other fungi. FEMS Immunology and Medical Microbiology, 2011, 63, 363-372.	2.7	8