

# Yu-Ping Ning

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5137209/publications.pdf>

Version: 2024-02-01

75  
papers

1,352  
citations

361045

20  
h-index

414034

32  
g-index

78  
all docs

78  
docs citations

78  
times ranked

1358  
citing authors

#	ARTICLE	IF	CITATIONS
1	Rapid and longer-term antidepressant effects of repeated-dose intravenous ketamine for patients with unipolar and bipolar depression. <i>Journal of Psychiatric Research</i> , 2018, 106, 61-68.	1.5	119
2	Adjunctive intranasal esketamine for major depressive disorder: A systematic review of randomized double-blind controlled-placebo studies. <i>Journal of Affective Disorders</i> , 2020, 265, 63-70.	2.0	68
3	Antidepressant effect of repeated ketamine administration on kynurenine pathway metabolites in patients with unipolar and bipolar depression. <i>Brain, Behavior, and Immunity</i> , 2018, 74, 205-212.	2.0	63
4	Major depressive disorder and suicide risk among adult outpatients at several general hospitals in a Chinese Han population. <i>PLoS ONE</i> , 2017, 12, e0186143.	1.1	59
5	Neurocognitive effects of six ketamine infusions and the association with antidepressant response in patients with unipolar and bipolar depression. <i>Journal of Psychopharmacology</i> , 2018, 32, 1118-1126.	2.0	53
6	Neurocognitive performance and repeated-dose intravenous ketamine in major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 246, 241-247.	2.0	50
7	A preliminary study of anti-suicidal efficacy of repeated ketamine infusions in depression with suicidal ideation. <i>Journal of Affective Disorders</i> , 2019, 251, 205-212.	2.0	46
8	Alterations of multiple peripheral inflammatory cytokine levels after repeated ketamine infusions in major depressive disorder. <i>Translational Psychiatry</i> , 2020, 10, 246.	2.4	42
9	Hypomethylation of LINE-1 elements in schizophrenia and bipolar disorder. <i>Journal of Psychiatric Research</i> , 2018, 107, 68-72.	1.5	41
10	Cross-sectional relationship between kynurenine pathway metabolites and cognitive function in major depressive disorder. <i>Psychoneuroendocrinology</i> , 2019, 101, 72-79.	1.3	40
11	Cognitive Impairment and Structural Abnormalities in Late Life Depression with Olfactory Identification Impairment: an Alzheimer's Disease-Like Pattern. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 640-648.	1.0	37
12	Investigation of medical effect of multiple ketamine infusions on patients with major depressive disorder. <i>Journal of Psychopharmacology</i> , 2019, 33, 494-501.	2.0	35
13	A preliminary study of adjunctive ketamine for treatment-resistant bipolar depression. <i>Journal of Affective Disorders</i> , 2020, 275, 38-43.	2.0	31
14	Serum brain-derived neurotrophic factor and nerve growth factor decreased in chronic ketamine abusers. <i>Drug and Alcohol Dependence</i> , 2014, 142, 290-294.	1.6	30
15	A Selective Review of the Excitatory-Inhibitory Imbalance in Schizophrenia: Underlying Biology, Genetics, Microcircuits, and Symptoms. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 664535.	1.8	30
16	Volumetric changes in subcortical structures following repeated ketamine treatment in patients with major depressive disorder: a longitudinal analysis. <i>Translational Psychiatry</i> , 2020, 10, 264.	2.4	28
17	Repeated intravenous infusions of ketamine: Neurocognition in patients with anxious and nonanxious treatment-resistant depression. <i>Journal of Affective Disorders</i> , 2019, 259, 1-6.	2.0	27
18	Adjunctive minocycline for major mental disorders: A systematic review. <i>Journal of Psychopharmacology</i> , 2019, 33, 1215-1226.	2.0	27

#	ARTICLE	IF	CITATIONS
19	Impairment in the goal-directed corticostriatal learning system as a biomarker for obsessive-compulsive disorder. <i>Psychological Medicine</i> , 2020, 50, 1490-1500.	2.7	26
20	Expert consensus on the prevention and treatment of substance use and addictive behaviour-related disorders during the COVID-19 pandemic. <i>Annals of General Psychiatry</i> , 2020, 33, e100252.	1.1	26
21	Plasma inflammatory cytokines and treatment-resistant depression with comorbid pain: improvement by ketamine. <i>Journal of Neuroinflammation</i> , 2021, 18, 200.	3.1	26
22	Adjunctive raloxifene for postmenopausal women with schizophrenia: A meta-analysis of randomized, double-blind, placebo-controlled trials. <i>Schizophrenia Research</i> , 2018, 197, 288-293.	1.1	25
23	Interactive Effect of Depression and Cognitive Impairment on Olfactory Identification in Elderly People. <i>Journal of Alzheimer's Disease</i> , 2018, 66, 1645-1655.	1.2	19
24	Relationship between hippocampal volume and inflammatory markers following six infusions of ketamine in major depressive disorder. <i>Journal of Affective Disorders</i> , 2020, 276, 608-615.	2.0	19
25	&lt;p&gt;Increased Plasma Kynurenic Acid Levels are Associated with Impaired Attention/Vigilance and Social Cognition in Patients with Schizophrenia&lt;/p&gt;. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 263-271.	1.0	18
26	Working memory associated with anti-suicidal ideation effect of repeated-dose intravenous ketamine in depressed patients. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 431-438.	1.8	17
27	Association between depression subtypes and response to repeated-dose intravenous ketamine. <i>Acta Psychiatrica Scandinavica</i> , 2019, 140, 446-457.	2.2	16
28	Association between cognition and suicidal ideation in patients with major depressive disorder: A longitudinal study. <i>Journal of Affective Disorders</i> , 2020, 272, 146-151.	2.0	16
29	Gender differences in the prevalence and clinical correlates of sleep disturbance in general hospital outpatients. <i>Psychiatry Research</i> , 2018, 269, 134-139.	1.7	15
30	Predictors of response to repeated ketamine infusions in depression with suicidal ideation: An ROC curve analysis. <i>Journal of Affective Disorders</i> , 2020, 264, 263-271.	2.0	15
31	Sleep improvement is associated with the antidepressant efficacy of repeated-dose ketamine and serum BDNF levels: a post-hoc analysis. <i>Pharmacological Reports</i> , 2021, 73, 594-603.	1.5	15
32	Comparative effectiveness of repeated ketamine infusions in treating anhedonia in bipolar and unipolar depression. <i>Journal of Affective Disorders</i> , 2022, 300, 109-113.	2.0	15
33	Interactive effects of elevated homocysteine and late-life depression on cognitive impairment. <i>Journal of Affective Disorders</i> , 2020, 277, 212-217.	2.0	14
34	Prevalence, clinical correlates and IQ of suicidal ideation in drug-naïve Chinese Han patients with major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 248, 59-64.	2.0	13
35	Difference of olfactory deficit in patients with acute episode of schizophrenia and major depressive episode. <i>Schizophrenia Research</i> , 2019, 212, 99-106.	1.1	12
36	The additive effect of late-life depression and olfactory dysfunction on the risk of dementia was mediated by hypersynchronization of the hippocampus/fusiform gyrus. <i>Translational Psychiatry</i> , 2021, 11, 172.	2.4	12

#	ARTICLE	IF	CITATIONS
37	The interaction between estradiol change and the serotonin transporter gene (5-HTTLPR) polymorphism is associated with postpartum depressive symptoms. <i>Psychiatric Genetics</i> , 2019, 29, 97-102.	0.6	10
38	<p>BDNF and the Antidepressant Effects of Ketamine and Propofol in Electroconvulsive Therapy: A Preliminary Study</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 901-908.	1.0	9
39	Cardiovascular effects of repeated subanaesthetic ketamine infusion in depression. <i>Journal of Psychopharmacology</i> , 2021, 35, 159-167.	2.0	9
40	The potential pro-cognitive effects with intravenous subanesthetic ketamine in adults with treatment-resistant major depressive or bipolar disorders and suicidality. <i>Journal of Psychiatric Research</i> , 2021, 144, 312-319.	1.5	9
41	Consistent brain structural abnormalities and multisite individualised classification of schizophrenia using deep neural networks. <i>British Journal of Psychiatry</i> , 2022, 221, 732-739.	1.7	9
42	Functional connectivity between the habenula and default mode network and its association with the antidepressant effect of ketamine. <i>Depression and Anxiety</i> , 2022, 39, 352-362.	2.0	9
43	The relationship between plasma cytokine levels and antidepressant response in patients with first-episode major depressive disorder. <i>Journal of Affective Disorders</i> , 2021, 287, 327-333.	2.0	8
44	Retrospective assessment of factors associated with readmission in a large psychiatric hospital in Guangzhou, China. <i>Shanghai Archives of Psychiatry</i> , 2014, 26, 138-48.	0.7	8
45	Association between plasma levels of BDNF and the antisuicidal effects of repeated ketamine infusions in depression with suicidal ideation. <i>Therapeutic Advances in Psychopharmacology</i> , 2020, 10, 204512532097379.	1.2	7
46	Prevalence and risk factors for mental health symptoms in community epidemic prevention workers during the postpandemic era of COVID-19 in China. <i>Psychiatry Research</i> , 2021, 304, 114132.	1.7	7
47	Baseline insomnia as a predictor of antidepressant efficacy to repeated intravenous ketamine for unipolar and bipolar depression: A preliminary study. <i>Journal of Affective Disorders</i> , 2020, 271, 1-8.	2.0	6
48	Diabetes in late-life schizophrenia: Prevalence, factors, and association with clinical symptoms. <i>Journal of Psychiatric Research</i> , 2021, 132, 44-49.	1.5	6
49	Association of VEGF With Antianhedonic Effects of Repeated-Dose Intravenous Ketamine in Treatment-Refractory Depression. <i>Frontiers in Psychiatry</i> , 2021, 12, 780975.	1.3	6
50	Long-term outcomes of repeated ketamine infusions in patients with unipolar and bipolar depression: A naturalistic follow-up study. <i>Journal of Affective Disorders</i> , 2022, 300, 172-178.	2.0	6
51	The effectiveness of repeated intravenous ketamine on subjective and objective psychosocial function in patients with treatment-resistant depression and suicidal ideation. <i>Journal of Affective Disorders</i> , 2022, 304, 78-84.	2.0	6
52	Odor-related brain hyper-reactivity in euthymic bipolar disorder: An fMRI and ERP study. <i>Psychiatry Research</i> , 2019, 278, 218-227.	1.7	5
53	A Preliminary Study of Adjunctive Nonconvulsive Electrotherapy for Treatment-Refractory Depression. <i>Psychiatric Quarterly</i> , 2021, 92, 311-320.	1.1	5
54	Association of plasma VEGF levels and the antidepressant effects of ketamine in patients with depression. <i>Therapeutic Advances in Psychopharmacology</i> , 2021, 11, 204512532110143.	1.2	5

#	ARTICLE	IF	CITATIONS
55	Development of a checklist of short-term and long-term psychological symptoms associated with ketamine use. Shanghai Archives of Psychiatry, 2015, 27, 186-94.	0.7	5
56	Neurocognitive effects of six ketamine infusions and the association with antidepressant effects in treatment-resistant bipolar depression: a preliminary study. PeerJ, 2020, 8, e10208.	0.9	5
57	Neuropsychiatric Symptoms Mediated the Relationship Between Odor Identification and Cognition in Alzheimer's Disease Spectrum: A Structural Equation Model Analysis. Frontiers in Aging Neuroscience, 2021, 13, 732840.	1.7	5
58	Predictors of 4-week antidepressant outcome in patients with first-episode major depressive disorder: An ROC curve analysis. Journal of Affective Disorders, 2022, 304, 59-65.	2.0	5
59	Mediating effect of neurocognition between severity of symptoms and social-occupational function in anxious depression. Journal of Affective Disorders, 2019, 246, 667-673.	2.0	4
60	Determining the effects of LLD and MCI on brain decline according to machine learning and a structural covariance network analysis. Journal of Psychiatric Research, 2020, 126, 43-54.	1.5	4
61	&lt;p&gt;Serum BDNF Levels are Not Associated with the Antidepressant Effects of Nonconvulsive Electrotherapy&lt;/p&gt;. Neuropsychiatric Disease and Treatment, 2020, Volume 16, 1555-1560.	1.0	4
62	Reduction of phonetic mismatch negativity may depict illness course and predict functional outcomes in schizophrenia. Journal of Psychiatric Research, 2021, 137, 290-297.	1.5	4
63	Differences in Odor Identification in Early-Onset and Late-Onset Depression. Brain Sciences, 2022, 12, 276.	1.1	4
64	Plasma VEGF Concentrations and Ketamine's Effects on Suicidal Ideation in Depression With Suicidal Ideation. Frontiers in Psychiatry, 2022, 13, 855995.	1.3	4
65	Cognitive Function Mediates the Anti-suicide Effect of Repeated Intravenous Ketamine in Adult Patients With Suicidal Ideation. Frontiers in Psychiatry, 2022, 13, 779326.	1.3	4
66	Resting-state functional connectivity of the amygdala in major depressive disorder with suicidal ideation. Journal of Psychiatric Research, 2022, 153, 189-196.	1.5	4
67	Prevalence and risk factors for depression in outpatient departments of three general hospitals in China: a cross-sectional study. International Journal of Psychiatry in Clinical Practice, 2020, 24, 88-95.	1.2	3
68	&lt;p&gt;Physical Comorbidities are Independently Associated with Higher Rates of Psychiatric Readmission in a Chinese Han Population&lt;/p&gt;. Neuropsychiatric Disease and Treatment, 2020, Volume 16, 2073-2082.	1.0	3
69	Adjunctive intranasal esketamine for major depressive disorder: a systematic review of randomized double-blind controlled-placebo studies - Authors's reply. Journal of Affective Disorders, 2020, 274, 955.	2.0	3
70	&lt;p&gt;Clozapine Use in Patients with Early-Stage Schizophrenia in a Chinese Psychiatric Hospital&lt;/p&gt;. Neuropsychiatric Disease and Treatment, 2020, Volume 16, 2827-2836.	1.0	3
71	The challenge and response of mental health institutions in COVID-19 pandemic: from chaos to new normal. Translational Psychiatry, 2020, 10, 386.	2.4	3
72	Interactive Effects of Agitation and Cognitive Impairment on Odor Identification in Patients With Late-Life Depression. Frontiers in Psychiatry, 2022, 13, 839012.	1.3	3

#	ARTICLE	IF	CITATIONS
73	Disrupted olfactory functional connectivity in patients with late-life depression. <i>Journal of Affective Disorders</i> , 2022, 306, 174-181.	2.0	3
74	Pre-treatment Pain Symptoms Influence Antidepressant Response to Ketamine in Depressive Patients. <i>Frontiers in Psychiatry</i> , 2022, 13, 793677.	1.3	2
75	Aberrant Dynamic Functional Connectivity of Posterior Cingulate Cortex Subregions in Major Depressive Disorder With Suicidal Ideation. <i>Frontiers in Neuroscience</i> , 0, 16, .	1.4	2