

Yu-Ping Ning

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5137209/publications.pdf>

Version: 2024-02-01

75
papers

1,352
citations

361045

20
h-index

414034

32
g-index

78
all docs

78
docs citations

78
times ranked

1358
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Long-term outcomes of repeated ketamine infusions in patients with unipolar and bipolar depression: A naturalistic follow-up study. <i>Journal of Affective Disorders</i> , 2022, 300, 172-178. | 2.0 | 6 |
| 2 | Comparative effectiveness of repeated ketamine infusions in treating anhedonia in bipolar and unipolar depression. <i>Journal of Affective Disorders</i> , 2022, 300, 109-113. | 2.0 | 15 |
| 3 | Predictors of 4-week antidepressant outcome in patients with first-episode major depressive disorder: An ROC curve analysis. <i>Journal of Affective Disorders</i> , 2022, 304, 59-65. | 2.0 | 5 |
| 4 | Consistent brain structural abnormalities and multisite individualised classification of schizophrenia using deep neural networks. <i>British Journal of Psychiatry</i> , 2022, 221, 732-739. | 1.7 | 9 |
| 5 | Differences in Odor Identification in Early-Onset and Late-Onset Depression. <i>Brain Sciences</i> , 2022, 12, 276. | 1.1 | 4 |
| 6 | Interactive Effects of Agitation and Cognitive Impairment on Odor Identification in Patients With Late-Life Depression. <i>Frontiers in Psychiatry</i> , 2022, 13, 839012. | 1.3 | 3 |
| 7 | Pre-treatment Pain Symptoms Influence Antidepressant Response to Ketamine in Depressive Patients. <i>Frontiers in Psychiatry</i> , 2022, 13, 793677. | 1.3 | 2 |
| 8 | The effectiveness of repeated intravenous ketamine on subjective and objective psychosocial function in patients with treatment-resistant depression and suicidal ideation. <i>Journal of Affective Disorders</i> , 2022, 304, 78-84. | 2.0 | 6 |
| 9 | Disrupted olfactory functional connectivity in patients with late-life depression. <i>Journal of Affective Disorders</i> , 2022, 306, 174-181. | 2.0 | 3 |
| 10 | Functional connectivity between the habenula and default mode network and its association with the antidepressant effect of ketamine. <i>Depression and Anxiety</i> , 2022, 39, 352-362. | 2.0 | 9 |
| 11 | Plasma VEGF Concentrations and Ketamine's Effects on Suicidal Ideation in Depression With Suicidal Ideation. <i>Frontiers in Psychiatry</i> , 2022, 13, 855995. | 1.3 | 4 |
| 12 | Cognitive Function Mediates the Anti-suicide Effect of Repeated Intravenous Ketamine in Adult Patients With Suicidal Ideation. <i>Frontiers in Psychiatry</i> , 2022, 13, 779326. | 1.3 | 4 |
| 13 | Resting-state functional connectivity of the amygdala in major depressive disorder with suicidal ideation. <i>Journal of Psychiatric Research</i> , 2022, 153, 189-196. | 1.5 | 4 |
| 14 | Cardiovascular effects of repeated subanaesthetic ketamine infusion in depression. <i>Journal of Psychopharmacology</i> , 2021, 35, 159-167. | 2.0 | 9 |
| 15 | A Preliminary Study of Adjunctive Nonconvulsive Electrotherapy for Treatment-Refractory Depression. <i>Psychiatric Quarterly</i> , 2021, 92, 311-320. | 1.1 | 5 |
| 16 | Diabetes in late-life schizophrenia: Prevalence, factors, and association with clinical symptoms. <i>Journal of Psychiatric Research</i> , 2021, 132, 44-49. | 1.5 | 6 |
| 17 | Association of plasma VEGF levels and the antidepressant effects of ketamine in patients with depression. <i>Therapeutic Advances in Psychopharmacology</i> , 2021, 11, 204512532110143. | 1.2 | 5 |
| 18 | The additive effect of late-life depression and olfactory dysfunction on the risk of dementia was mediated by hypersynchronization of the hippocampus/fusiform gyrus. <i>Translational Psychiatry</i> , 2021, 11, 172. | 2.4 | 12 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | The relationship between plasma cytokine levels and antidepressant response in patients with first-episode major depressive disorder. <i>Journal of Affective Disorders</i> , 2021, 287, 327-333. | 2.0 | 8 |
| 20 | Reduction of phonetic mismatch negativity may depict illness course and predict functional outcomes in schizophrenia. <i>Journal of Psychiatric Research</i> , 2021, 137, 290-297. | 1.5 | 4 |
| 21 | Plasma inflammatory cytokines and treatment-resistant depression with comorbid pain: improvement by ketamine. <i>Journal of Neuroinflammation</i> , 2021, 18, 200. | 3.1 | 26 |
| 22 | Prevalence and risk factors for mental health symptoms in community epidemic prevention workers during the postpandemic era of COVID-19 in China. <i>Psychiatry Research</i> , 2021, 304, 114132. | 1.7 | 7 |
| 23 | Working memory associated with anti-suicidal ideation effect of repeated-dose intravenous ketamine in depressed patients. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 431-438. | 1.8 | 17 |
| 24 | Sleep improvement is associated with the antidepressant efficacy of repeated-dose ketamine and serum BDNF levels: a post-hoc analysis. <i>Pharmacological Reports</i> , 2021, 73, 594-603. | 1.5 | 15 |
| 25 | A Selective Review of the Excitatory-Inhibitory Imbalance in Schizophrenia: Underlying Biology, Genetics, Microcircuits, and Symptoms. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 664535. | 1.8 | 30 |
| 26 | The potential pro-cognitive effects with intravenous subanesthetic ketamine in adults with treatment-resistant major depressive or bipolar disorders and suicidality. <i>Journal of Psychiatric Research</i> , 2021, 144, 312-319. | 1.5 | 9 |
| 27 | Association of VEGF With Antianhedonic Effects of Repeated-Dose Intravenous Ketamine in Treatment-Refractory Depression. <i>Frontiers in Psychiatry</i> , 2021, 12, 780975. | 1.3 | 6 |
| 28 | Neuropsychiatric Symptoms Mediated the Relationship Between Odor Identification and Cognition in Alzheimer's Disease Spectrum: A Structural Equation Model Analysis. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, 732840. | 1.7 | 5 |
| 29 | Impairment in the goal-directed corticostriatal learning system as a biomarker for obsessive-compulsive disorder. <i>Psychological Medicine</i> , 2020, 50, 1490-1500. | 2.7 | 26 |
| 30 | Prevalence and risk factors for depression in outpatient departments of three general hospitals in China: a cross-sectional study. <i>International Journal of Psychiatry in Clinical Practice</i> , 2020, 24, 88-95. | 1.2 | 3 |
| 31 | Adjunctive intranasal esketamine for major depressive disorder: A systematic review of randomized double-blind controlled-placebo studies. <i>Journal of Affective Disorders</i> , 2020, 265, 63-70. | 2.0 | 68 |
| 32 | <p>Physical Comorbidities are Independently Associated with Higher Rates of Psychiatric Readmission in a Chinese Han Population</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 2073-2082. | 1.0 | 3 |
| 33 | Adjunctive intranasal esketamine for major depressive disorder: a systematic review of randomized double-blind controlled-placebo studies - Authors' reply. <i>Journal of Affective Disorders</i> , 2020, 274, 955. | 2.0 | 3 |
| 34 | Association between plasma levels of BDNF and the antisuicidal effects of repeated ketamine infusions in depression with suicidal ideation. <i>Therapeutic Advances in Psychopharmacology</i> , 2020, 10, 204512532097379. | 1.2 | 7 |
| 35 | <p>Clozapine Use in Patients with Early-Stage Schizophrenia in a Chinese Psychiatric Hospital</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 2827-2836. | 1.0 | 3 |
| 36 | Alterations of multiple peripheral inflammatory cytokine levels after repeated ketamine infusions in major depressive disorder. <i>Translational Psychiatry</i> , 2020, 10, 246. | 2.4 | 42 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Volumetric changes in subcortical structures following repeated ketamine treatment in patients with major depressive disorder: a longitudinal analysis. <i>Translational Psychiatry</i> , 2020, 10, 264. | 2.4 | 28 |
| 38 | Relationship between hippocampal volume and inflammatory markers following six infusions of ketamine in major depressive disorder. <i>Journal of Affective Disorders</i> , 2020, 276, 608-615. | 2.0 | 19 |
| 39 | Interactive effects of elevated homocysteine and late-life depression on cognitive impairment. <i>Journal of Affective Disorders</i> , 2020, 277, 212-217. | 2.0 | 14 |
| 40 | Expert consensus on the prevention and treatment of substance use and addictive behaviour-related disorders during the COVID-19 pandemic. <i>Annals of General Psychiatry</i> , 2020, 33, e100252. | 1.1 | 26 |
| 41 | The challenge and response of mental health institutions in COVID-19 pandemic: from chaos to new normal. <i>Translational Psychiatry</i> , 2020, 10, 386. | 2.4 | 3 |
| 42 | Association between cognition and suicidal ideation in patients with major depressive disorder: A longitudinal study. <i>Journal of Affective Disorders</i> , 2020, 272, 146-151. | 2.0 | 16 |
| 43 | Determining the effects of LLD and MCI on brain decline according to machine learning and a structural covariance network analysis. <i>Journal of Psychiatric Research</i> , 2020, 126, 43-54. | 1.5 | 4 |
| 44 | <p>Increased Plasma Kynurenic Acid Levels are Associated with Impaired Attention/Vigilance and Social Cognition in Patients with Schizophrenia</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 263-271. | 1.0 | 18 |
| 45 | <p>Serum BDNF Levels are Not Associated with the Antidepressant Effects of Nonconvulsive Electrotherapy</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 1555-1560. | 1.0 | 4 |
| 46 | A preliminary study of adjunctive ketamine for treatment-resistant bipolar depression. <i>Journal of Affective Disorders</i> , 2020, 275, 38-43. | 2.0 | 31 |
| 47 | Predictors of response to repeated ketamine infusions in depression with suicidal ideation: An ROC curve analysis. <i>Journal of Affective Disorders</i> , 2020, 264, 263-271. | 2.0 | 15 |
| 48 | <p>BDNF and the Antidepressant Effects of Ketamine and Propofol in Electroconvulsive Therapy: A Preliminary Study</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 901-908. | 1.0 | 9 |
| 49 | Baseline insomnia as a predictor of antidepressant efficacy to repeated intravenous ketamine for unipolar and bipolar depression: A preliminary study. <i>Journal of Affective Disorders</i> , 2020, 271, 1-8. | 2.0 | 6 |
| 50 | Neurocognitive effects of six ketamine infusions and the association with antidepressant effects in treatment-resistant bipolar depression: a preliminary study. <i>PeerJ</i> , 2020, 8, e10208. | 0.9 | 5 |
| 51 | Difference of olfactory deficit in patients with acute episode of schizophrenia and major depressive episode. <i>Schizophrenia Research</i> , 2019, 212, 99-106. | 1.1 | 12 |
| 52 | Repeated intravenous infusions of ketamine: Neurocognition in patients with anxious and nonanxious treatment-resistant depression. <i>Journal of Affective Disorders</i> , 2019, 259, 1-6. | 2.0 | 27 |
| 53 | Adjunctive minocycline for major mental disorders: A systematic review. <i>Journal of Psychopharmacology</i> , 2019, 33, 1215-1226. | 2.0 | 27 |
| 54 | Association between depression subtypes and response to repeatedâledose intravenous ketamine. <i>Acta Psychiatrica Scandinavica</i> , 2019, 140, 446-457. | 2.2 | 16 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Odor-related brain hyper-reactivity in euthymic bipolar disorder: An fMRI and ERP study. <i>Psychiatry Research</i> , 2019, 278, 218-227. | 1.7 | 5 |
| 56 | A preliminary study of anti-suicidal efficacy of repeated ketamine infusions in depression with suicidal ideation. <i>Journal of Affective Disorders</i> , 2019, 251, 205-212. | 2.0 | 46 |
| 57 | Investigation of medical effect of multiple ketamine infusions on patients with major depressive disorder. <i>Journal of Psychopharmacology</i> , 2019, 33, 494-501. | 2.0 | 35 |
| 58 | The interaction between estradiol change and the serotonin transporter gene (5-HTTLPR) polymorphism is associated with postpartum depressive symptoms. <i>Psychiatric Genetics</i> , 2019, 29, 97-102. | 0.6 | 10 |
| 59 | Neurocognitive performance and repeated-dose intravenous ketamine in major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 246, 241-247. | 2.0 | 50 |
| 60 | Mediating effect of neurocognition between severity of symptoms and social-occupational function in anxious depression. <i>Journal of Affective Disorders</i> , 2019, 246, 667-673. | 2.0 | 4 |
| 61 | Prevalence, clinical correlates and IQ of suicidal ideation in drug naïve Chinese Han patients with major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 248, 59-64. | 2.0 | 13 |
| 62 | Cross-sectional relationship between kynurenine pathway metabolites and cognitive function in major depressive disorder. <i>Psychoneuroendocrinology</i> , 2019, 101, 72-79. | 1.3 | 40 |
| 63 | Adjunctive raloxifene for postmenopausal women with schizophrenia: A meta-analysis of randomized, double-blind, placebo-controlled trials. <i>Schizophrenia Research</i> , 2018, 197, 288-293. | 1.1 | 25 |
| 64 | Cognitive Impairment and Structural Abnormalities in Late Life Depression with Olfactory Identification Impairment: an Alzheimer's Disease-Like Pattern. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 640-648. | 1.0 | 37 |
| 65 | Interactive Effect of Depression and Cognitive Impairment on Olfactory Identification in Elderly People. <i>Journal of Alzheimer's Disease</i> , 2018, 66, 1645-1655. | 1.2 | 19 |
| 66 | Neurocognitive effects of six ketamine infusions and the association with antidepressant response in patients with unipolar and bipolar depression. <i>Journal of Psychopharmacology</i> , 2018, 32, 1118-1126. | 2.0 | 53 |
| 67 | Rapid and longer-term antidepressant effects of repeated-dose intravenous ketamine for patients with unipolar and bipolar depression. <i>Journal of Psychiatric Research</i> , 2018, 106, 61-68. | 1.5 | 119 |
| 68 | Hypomethylation of LINE-1 elements in schizophrenia and bipolar disorder. <i>Journal of Psychiatric Research</i> , 2018, 107, 68-72. | 1.5 | 41 |
| 69 | Antidepressant effect of repeated ketamine administration on kynurenine pathway metabolites in patients with unipolar and bipolar depression. <i>Brain, Behavior, and Immunity</i> , 2018, 74, 205-212. | 2.0 | 63 |
| 70 | Gender differences in the prevalence and clinical correlates of sleep disturbance in general hospital outpatients. <i>Psychiatry Research</i> , 2018, 269, 134-139. | 1.7 | 15 |
| 71 | Major depressive disorder and suicide risk among adult outpatients at several general hospitals in a Chinese Han population. <i>PLoS ONE</i> , 2017, 12, e0186143. | 1.1 | 59 |
| 72 | Development of a checklist of short-term and long-term psychological symptoms associated with ketamine use. <i>Shanghai Archives of Psychiatry</i> , 2015, 27, 186-94. | 0.7 | 5 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Serum brain-derived neurotrophic factor and nerve growth factor decreased in chronic ketamine abusers. <i>Drug and Alcohol Dependence</i> , 2014, 142, 290-294. | 1.6 | 30 |
| 74 | Retrospective assessment of factors associated with readmission in a large psychiatric hospital in Guangzhou, China. <i>Shanghai Archives of Psychiatry</i> , 2014, 26, 138-48. | 0.7 | 8 |
| 75 | Aberrant Dynamic Functional Connectivity of Posterior Cingulate Cortex Subregions in Major Depressive Disorder With Suicidal Ideation. <i>Frontiers in Neuroscience</i> , 0, 16, . | 1.4 | 2 |