

# Michał Zakliczyński

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5138559/publications.pdf>

Version: 2024-02-01

27  
papers

55  
citations

1684188

5  
h-index

1872680

6  
g-index

27  
all docs

27  
docs citations

27  
times ranked

133  
citing authors

#	ARTICLE	IF	CITATIONS
1	Patient-Related Preoperative Clinical Factors Influencing 1-Year Survival After Orthotopic Heart Transplantation – A Single Center Polish Experience. <i>Annals of Transplantation</i> , 2022, 27, e934185.	0.9	1
2	COVID-19 mortality in patients after orthotopic heart transplantation: A single-center one-year observational study. <i>Kardiologia Polska</i> , 2022, 80, 215-217.	0.6	0
3	Kidney Re-Transplantation after Simultaneous Heart and Kidney Transplant: Case Study and Literature Review. <i>Transplantology</i> , 2022, 3, 124-129.	0.6	0
4	Occurrence of Professional Burnout and Severity of Depressive Symptoms among Cardiac Nurses: A Cross-Sectional Study. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 12038.	2.6	3
5	P1689COMPARISON OF NON-STEROIDAL ANTI-INFLAMMATORY DRUGS(NSAIDS) AND/OR PAINKILLERS USE BY KIDNEY (KTRS) AND HEART TRANSPLANT RECIPIENTS (HTRS). <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
6	Prevalence and Risk Factors of New-Onset Diabetes After Transplantation (NODAT). <i>Annals of Transplantation</i> , 2020, 25, e926556.	0.9	9
7	Short and long-term outcomes of continuous-flow left ventricular assist device therapy in 79 patients with end-stage heart failure. <i>Polish Archives of Internal Medicine</i> , 2020, 130, 589-597.	0.4	0
8	Mechanical circulatory support is effective to treat pulmonary hypertension in heart transplant candidates disqualified due to unacceptable pulmonary vascular resistance. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2018, 15, 23-26.	0.1	1
9	Prognostic utility of the N terminal prohormone of brain natriuretic peptide and the modified Model for End Stage Liver Disease in patients with end stage heart failure. <i>Polish Archives of Internal Medicine</i> , 2018, 128, 235-243.	0.4	5
10	The first successful bridging with an Impella CP® to minimally invasive HeartMate 3 LVAD implantation in Poland. <i>Kardiologia Polska</i> , 2018, 76, 209-209.	0.6	0
11	Cardiac retransplantation as a promising treatment option for late graft failure – Zabrze experience. <i>Kardiologia Polska</i> , 2018, 76, 1015-1017.	0.6	0
12	Emergency HeartWare Ventricular Assist Device (HVAD) exchange due to pump thrombosis using minimally invasive technique. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2017, 1, 76-78.	0.1	0
13	One year survival of ambulatory patients with the end-stage heart failure: the analysis of prognostic factors. <i>Polish Archives of Internal Medicine</i> , 2017, 127, 254-260.	0.4	5
14	Monitoring hemostasis parameters in left ventricular assist device recipients – a preliminary report. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2016, 3, 224-228.	0.1	2
15	Preoperative Echocardiography Examination of Right Ventricle Function in Patients Scheduled for LVAD Implantation Correlates with Postoperative Hemodynamic Examinations. <i>Annals of Transplantation</i> , 2016, 21, 500-507.	0.9	5
16	Recurrent asymptomatic acute cellular rejection after heart transplantation: monitoring with speckle-tracking echocardiography. <i>Polish Archives of Internal Medicine</i> , 2016, 126, 700-703.	0.4	0
17	Activated Hemostatic Biomarkers in Patients with Implanted Left Ventricle Assist Devices: Are Heparin and/or Clopidogrel Justified?. <i>Cardiology</i> , 2015, 131, 172-176.	1.4	2
18	Professional and social activity of patients after heart transplant. <i>International Journal of Occupational Medicine and Environmental Health</i> , 2015, 28, 741-749.	1.3	3

#	ARTICLE	IF	CITATIONS
19	Left ventricular assist as a key factor for heart self-restoration in a 20-year-old patient with severe left ventricular dysfunction: five-year echocardiographic observation. <i>Kardiologia Polska</i> , 2015, 73, 783-783.	0.6	0
20	The Model for End-Stage Liver Disease (MELD) can predict outcomes in ambulatory patients with advanced heart failure who have been referred for cardiac transplantation evaluation. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2014, 2, 178-181.	0.1	7
21	Coincidence of cellular and antibody mediated rejection in heart transplant recipients – preliminary report. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2014, 1, 52-55.	0.1	2
22	“second delivery after heart transplantation” – a case study. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2014, 3, 339-342.	0.1	1
23	Bridge to recovery in two cases of dilated cardiomyopathy after long-term mechanical circulatory support. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2014, 2, 169-172.	0.1	3
24	Influence of proliferation signal inhibitors on vascular endothelial growth factor production in heart transplant recipients – preliminary report. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2014, 2, 173-177.	0.1	0
25	Five-year survival of patients with chronic systolic heart failure of ischemic and non-ischemic etiology: analysis of prognostic factors. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2014, 1, 56-62.	0.1	2
26	Depression is the strongest predictor of long-term outcome in patients with chronic nonischemic heart failure. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2014, 1, 102-108.	0.1	1
27	HEART AND LUNG FAILURE, TRANSPLANTOLOGY Low health-related quality of life is a predictor of major adverse cardiovascular events in patients with chronic nonischemic heart failure. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2014, 3, 283-288.	0.1	3