

Shinsuke Isobe

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5138774/publications.pdf>

Version: 2024-02-01

37
papers

406
citations

840776
11
h-index

794594
19
g-index

38
all docs

38
docs citations

38
times ranked

508
citing authors

#	ARTICLE	IF	CITATIONS
1	Circadian rhythm of blood pressure and the renin-angiotensin system in the kidney. Hypertension Research, 2017, 40, 413-422.	2.7	52
2	Disturbed circadian rhythm of the intrarenal renin-angiotensin system: relevant to nocturnal hypertension and renal damage. Clinical and Experimental Nephrology, 2015, 19, 231-239.	1.6	38
3	Impaired endogenous nighttime melatonin secretion relates to intrarenal renin-angiotensin system activation and renal damage in patients with chronic kidney disease. Clinical and Experimental Nephrology, 2016, 20, 878-884.	1.6	38
4	The pivotal role of melatonin in ameliorating chronic kidney disease by suppression of the renin-angiotensin system in the kidney. Hypertension Research, 2019, 42, 761-768.	2.7	36
5	Melatonin ameliorates intrarenal renin-angiotensin system in a 5/6 nephrectomy rat model. Clinical and Experimental Nephrology, 2018, 22, 539-549.	1.6	34
6	Augmented circadian rhythm of the intrarenal renin-angiotensin systems in anti-thymocyte serum nephritis rats. Hypertension Research, 2016, 39, 312-320.	2.7	22
7	Plasma Soluble (Pro)renin Receptor Reflects Renal Damage. PLoS ONE, 2016, 11, e0156165.	2.5	17
8	Alogliptin improves steroid-induced hyperglycemia in treatment-naïve Japanese patients with chronic kidney disease by decrease of plasma glucagon levels. Medical Science Monitor, 2014, 20, 587-593.	1.1	16
9	Intrarenal renin-angiotensin system activity is augmented after initiation of dialysis. Hypertension Research, 2017, 40, 364-370.	2.7	15
10	Prevalence of neural epidermal growth factor-like 1- and exostosin 1/exostosin 2-associated membranous nephropathy: a single-center retrospective study in Japan. Scientific Reports, 2022, 12, 2967.	3.3	15
11	Increased nocturnal blood pressure variability is associated with renal arteriolar hyalinosis in normotensive patients with IgA nephropathy. Hypertension Research, 2017, 40, 921-926.	2.7	12
12	Increased heart rate is associated with intrarenal renin-angiotensin system activation in chronic kidney disease patients. Clinical and Experimental Nephrology, 2019, 23, 1109-1118.	1.6	10
13	Primary Membranous Nephropathy With Enhanced Staining of Exostosin 1/Exostosin 2 in the Glomeruli: A Report of 2 Cases. Kidney Medicine, 2021, 3, 669-673.	2.0	10
14	A Rare Case of Lupus Nephritis Presenting as Thrombotic Microangiopathy with Diffuse Pseudotubulization Possibly Caused by Atypical Hemolytic Uremic Syndrome. Internal Medicine, 2018, 57, 1617-1623.	0.7	8
15	The level of urinary ± 1 microglobulin excretion is a useful marker of peritubular capillaritis in antineutrophil cytoplasmic antibody associated vasculitis. Clinical and Experimental Nephrology, 2015, 19, 851-858.	1.6	7
16	Night-time activation of the intrarenal renin-angiotensin system due to nocturnal hypertension is associated with renal arteriosclerosis in normotensive IgA nephropathy patients. Hypertension Research, 2018, 41, 334-341.	2.7	7
17	Hyperuricaemia is associated with renal damage independently of hypertension and intrarenal renin-angiotensin system activation, as well as their circadian rhythms. Nephrology, 2015, 20, 814-819.	1.6	6
18	Salt Loading Aggravates the Relationship between Melatonin and Proteinuria in Patients with Chronic Kidney Disease. Internal Medicine, 2019, 58, 1557-1564.	0.7	6

#	ARTICLE	IF	CITATIONS
19	The Relationship between the Intrarenal Dopamine System and Intrarenal Renin-angiotensin System Depending on the Renal Function. Internal Medicine, 2018, 57, 3241-3247.	0.7	5
20	Regular exercise and branched-chain amino acids prevent ischemic acute kidney injury-related muscle wasting in mice. Physiological Reports, 2020, 8, e14557.	1.7	5
21	Sodium-Glucose Cotransporter-2 Inhibitor Immediately Decreases Serum Uric Acid Levels in Type 2 Diabetic Patients. Medical Science Monitor, 2020, 26, e926086.	1.1	5
22	The Sequential Development of Antiglomerular Basement Membrane Nephritis and Myeloperoxidase-antineutrophil Cytoplasmic Antibody-associated Vasculitis. Internal Medicine, 2017, 56, 2617-2621.	0.7	4
23	The Intrarenal Renin-angiotensin System Is Activated Immediately after Kidney Donation in Kidney Transplant Donors. Internal Medicine, 2019, 58, 643-648.	0.7	4
24	The Urinary Angiotensinogen to Urinary Albumin Ratio Reflects Whether the Renin-angiotensin System in the Kidney Is Activated due to Filtration of Plasma Angiotensinogen through the Damaged Glomeruli or the Production of Angiotensinogen in the Proximal Tubules. Internal Medicine, 2020, 59, 357-364.	0.7	4
25	Salt intake causes B-type natriuretic peptide elevation independently of blood pressure elevation in the general population without hypertension and heart disease. Medicine (United States), 2021, 100, e25931.	1.0	4
26	Nicotinic acetylcholine receptor agonist reduces acute lung injury after renal ischemia-reperfusion injury by acting on splenic macrophages in mice. American Journal of Physiology - Renal Physiology, 2022, 322, F540-F552.	2.7	4
27	Focal segmental glomerulosclerosis associated with cutaneous and systemic plasmacytosis. CEN Case Reports, 2017, 6, 206-209.	0.9	3
28	A Case of Recurrent Atypical Anti-Glomerular Basement Membrane Nephritis Suspicion after Renal Transplantation. Nephron, 2020, 144, 49-53.	1.8	3
29	Circadian rhythm of the intrarenal renin-angiotensin system is caused by glomerular filtration of liver-derived angiotensinogen depending on glomerular capillary pressure in adriamycin nephropathy rats. Hypertension Research, 2021, 44, 618-627.	2.7	3
30	Baseline Urinary Angiotensinogen Excretion Predicts Deterioration of the Kidney Function in Patients with Chronic Kidney Disease. Internal Medicine, 2021, 60, 2201-2206.	0.7	3
31	Positive relationships between annual changes in salt intake and plasma B-type natriuretic peptide levels in the general population without hypertension and heart diseases. Hypertension Research, 2022, 45, 944-953.	2.7	3
32	A case of hypertensive disorders of pregnancy that developed at 9 weeks of gestation. CEN Case Reports, 2021, 10, 476-482.	0.9	2
33	Chronotherapy with a Renin-angiotensin System Inhibitor Ameliorates Renal Damage by Suppressing Intrarenal Renin-angiotensin System Activation. Internal Medicine, 2020, 59, 2237-2244.	0.7	2
34	A case report of thin basement membrane nephropathy accompanied by sporadic glomerulocystic kidney disease. BMC Nephrology, 2019, 20, 248.	1.8	1
35	Association between adrenal function and dialysis vintage in hemodialysis patients. Clinical and Experimental Nephrology, 0, , .	1.6	1
36	SP098CLINICAL USEFULNESS OF ADJUSTED D-DIMER CUTOFF VALUE TO EXCLUDE THROMBOEMBOLIC COMPLICATIONS IN THE PATIENTS WITH NEPHROTIC SYNDROME. Nephrology Dialysis Transplantation, 2018, 33, i377-i377.	0.7	0

#	ARTICLE	IF	CITATIONS
37	Case report: increased single-nephron estimated glomerular filtration rate in an adult patient with low birth weight. BMC Nephrology, 2020, 21, 75.	1.8	0