

# Shinsuke Isobe

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5138774/publications.pdf>

Version: 2024-02-01

37  
papers

406  
citations

840119

11  
h-index

794141

19  
g-index

38  
all docs

38  
docs citations

38  
times ranked

508  
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence of neural epidermal growth factor-like 1- and exostosin 1/exostosin 2-associated membranous nephropathy: a single-center retrospective study in Japan. <i>Scientific Reports</i> , 2022, 12, 2967.	1.6	15
2	Nicotinic acetylcholine receptor agonist reduces acute lung injury after renal ischemia-reperfusion injury by acting on splenic macrophages in mice. <i>American Journal of Physiology - Renal Physiology</i> , 2022, 322, F540-F552.	1.3	4
3	Positive relationships between annual changes in salt intake and plasma B-type natriuretic peptide levels in the general population without hypertension and heart diseases. <i>Hypertension Research</i> , 2022, 45, 944-953.	1.5	3
4	Circadian rhythm of the intrarenal renin-angiotensin system is caused by glomerular filtration of liver-derived angiotensinogen depending on glomerular capillary pressure in adriamycin nephropathy rats. <i>Hypertension Research</i> , 2021, 44, 618-627.	1.5	3
5	A case of hypertensive disorders of pregnancy that developed at 9 weeks of gestation. <i>CEN Case Reports</i> , 2021, 10, 476-482.	0.5	2
6	Salt intake causes B-type natriuretic peptide elevation independently of blood pressure elevation in the general population without hypertension and heart disease. <i>Medicine (United States)</i> , 2021, 100, e25931.	0.4	4
7	Baseline Urinary Angiotensinogen Excretion Predicts Deterioration of the Kidney Function in Patients with Chronic Kidney Disease. <i>Internal Medicine</i> , 2021, 60, 2201-2206.	0.3	3
8	Primary Membranous Nephropathy With Enhanced Staining of Exostosin 1/Exostosin 2 in the Glomeruli: A Report of 2 Cases. <i>Kidney Medicine</i> , 2021, 3, 669-673.	1.0	10
9	The Urinary Angiotensinogen to Urinary Albumin Ratio Reflects Whether the Renin-angiotensin System in the Kidney Is Activated due to Filtration of Plasma Angiotensinogen through the Damaged Glomeruli or the Production of Angiotensinogen in the Proximal Tubules. <i>Internal Medicine</i> , 2020, 59, 357-364.	0.3	4
10	Regular exercise and branched-chain amino acids prevent ischemic acute kidney injury-related muscle wasting in mice. <i>Physiological Reports</i> , 2020, 8, e14557.	0.7	5
11	A Case of Recurrent Atypical Anti-Glomerular Basement Membrane Nephritis Suspicion after Renal Transplantation. <i>Nephron</i> , 2020, 144, 49-53.	0.9	3
12	Case report: increased single-nephron estimated glomerular filtration rate in an adult patient with low birth weight. <i>BMC Nephrology</i> , 2020, 21, 75.	0.8	0
13	Sodium-Glucose Cotransporter-2 Inhibitor Immediately Decreases Serum Uric Acid Levels in Type 2 Diabetic Patients. <i>Medical Science Monitor</i> , 2020, 26, e926086.	0.5	5
14	Chronotherapy with a Renin-angiotensin System Inhibitor Ameliorates Renal Damage by Suppressing Intrarenal Renin-angiotensin System Activation. <i>Internal Medicine</i> , 2020, 59, 2237-2244.	0.3	2
15	A case report of thin basement membrane nephropathy accompanied by sporadic glomerulocystic kidney disease. <i>BMC Nephrology</i> , 2019, 20, 248.	0.8	1
16	Increased heart rate is associated with intrarenal renin-angiotensin system activation in chronic kidney disease patients. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 1109-1118.	0.7	10
17	Salt Loading Aggravates the Relationship between Melatonin and Proteinuria in Patients with Chronic Kidney Disease. <i>Internal Medicine</i> , 2019, 58, 1557-1564.	0.3	6
18	The Intrarenal Renin-angiotensin System Is Activated Immediately after Kidney Donation in Kidney Transplant Donors. <i>Internal Medicine</i> , 2019, 58, 643-648.	0.3	4

#	ARTICLE	IF	CITATIONS
19	The pivotal role of melatonin in ameliorating chronic kidney disease by suppression of the renin-angiotensin system in the kidney. <i>Hypertension Research</i> , 2019, 42, 761-768.	1.5	36
20	Melatonin ameliorates intrarenal renin-angiotensin system in a 5/6 nephrectomy rat model. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 539-549.	0.7	34
21	Night-time activation of the intrarenal renin-angiotensin system due to nocturnal hypertension is associated with renal arteriosclerosis in normotensive IgA nephropathy patients. <i>Hypertension Research</i> , 2018, 41, 334-341.	1.5	7
22	A Rare Case of Lupus Nephritis Presenting as Thrombotic Microangiopathy with Diffuse Pseudotubulization Possibly Caused by Atypical Hemolytic Uremic Syndrome. <i>Internal Medicine</i> , 2018, 57, 1617-1623.	0.3	8
23	SP098CLINICAL USEFULNESS OF ADJUSTED D-DIMER CUTOFF VALUE TO EXCLUDE THROMBOEMBOLIC COMPLICATIONS IN THE PATIENTS WITH NEPHROTIC SYNDROME. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i377-i377.	0.4	0
24	The Relationship between the Intrarenal Dopamine System and Intrarenal Renin-angiotensin System Depending on the Renal Function. <i>Internal Medicine</i> , 2018, 57, 3241-3247.	0.3	5
25	Circadian rhythm of blood pressure and the renin-angiotensin system in the kidney. <i>Hypertension Research</i> , 2017, 40, 413-422.	1.5	52
26	Focal segmental glomerulosclerosis associated with cutaneous and systemic plasmacytosis. <i>CEN Case Reports</i> , 2017, 6, 206-209.	0.5	3
27	Increased nocturnal blood pressure variability is associated with renal arteriolar hyalinosis in normotensive patients with IgA nephropathy. <i>Hypertension Research</i> , 2017, 40, 921-926.	1.5	12
28	Intrarenal renin-angiotensin system activity is augmented after initiation of dialysis. <i>Hypertension Research</i> , 2017, 40, 364-370.	1.5	15
29	The Sequential Development of Antiglomerular Basement Membrane Nephritis and Myeloperoxidase-antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Internal Medicine</i> , 2017, 56, 2617-2621.	0.3	4
30	Plasma Soluble (Pro)renin Receptor Reflects Renal Damage. <i>PLoS ONE</i> , 2016, 11, e0156165.	1.1	17
31	Augmented circadian rhythm of the intrarenal renin-angiotensin systems in anti-thymocyte serum nephritis rats. <i>Hypertension Research</i> , 2016, 39, 312-320.	1.5	22
32	Impaired endogenous nighttime melatonin secretion relates to intrarenal renin-angiotensin system activation and renal damage in patients with chronic kidney disease. <i>Clinical and Experimental Nephrology</i> , 2016, 20, 878-884.	0.7	38
33	Hyperuricaemia is associated with renal damage independently of hypertension and intrarenal renin-angiotensin system activation, as well as their circadian rhythms. <i>Nephrology</i> , 2015, 20, 814-819.	0.7	6
34	The level of urinary $\hat{\pm}$ 1 microglobulin excretion is a useful marker of peritubular capillaritis in antineutrophil cytoplasmic antibody associated vasculitis. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 851-858.	0.7	7
35	Disturbed circadian rhythm of the intrarenal renin-angiotensin system: relevant to nocturnal hypertension and renal damage. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 231-239.	0.7	38
36	Alogliptin improves steroid-induced hyperglycemia in treatment-naïve Japanese patients with chronic kidney disease by decrease of plasma glucagon levels. <i>Medical Science Monitor</i> , 2014, 20, 587-593.	0.5	16

#	ARTICLE	IF	CITATIONS
37	Association between adrenal function and dialysis vintage in hemodialysis patients. Clinical and Experimental Nephrology, 0, , .	0.7	1