David A Sassoon

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/513888/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Paternally expressed gene 3 (Pw1/Peg3) promotes sexual dimorphism in metabolism and behavior. PLoS Genetics, 2022, 18, e1010003. | 1.5 | 3 |
| 2 | Plateletâ€Derived Growth Factor Receptor Type α Activation Drives Pulmonary Vascular Remodeling Via Progenitor Cell Proliferation and Induces Pulmonary Hypertension. Journal of the American Heart Association, 2022, 11, e023021. | 1.6 | 5 |
| 3 | Hypoxia promotes a perinatal-like progenitor state in the adult murine epicardium. Scientific Reports, 2022, 12, . | 1.6 | 3 |
| 4 | Stem Cell Biology: Structure and Function – The Adult Stem Cell Niche: Multiple Cellular Players in Tissue Homeostasis and Regeneration. , 2022, , . | | 0 |
| 5 | Anti-integrin αv therapy improves cardiac fibrosis after myocardial infarction by blunting cardiac PW1+ stromal cells. Scientific Reports, 2020, 10, 11404. | 1.6 | 28 |
| 6 | The imprinted gene Pw1/Peg3 regulates skeletal muscle growth, satellite cell metabolic state, and self-renewal. Scientific Reports, 2018, 8, 14649. | 1.6 | 17 |
| 7 | Inhibition of the Activin Receptor Type-2B Pathway Restores Regenerative Capacity in Satellite Cell-Depleted Skeletal Muscle. Frontiers in Physiology, 2018, 9, 515. | 1.3 | 11 |
| 8 | FAPs are sensors for skeletal myofibre atrophy. Nature Cell Biology, 2018, 20, 864-865. | 4.6 | 4 |
| 9 | Odd skipped-related 1 (Osr1) identifies muscle-interstitial fibro-adipogenic progenitors (FAPs) activated by acute injury. Stem Cell Research, 2018, 32, 8-16. | 0.3 | 64 |
| 10 | Does cardiac development provide heart research with novel therapeutic approaches?. F1000Research, 2018, 7, 1756. | 0.8 | 7 |
| 11 | Peg3/PW1 Is a Marker of a Subset of Vessel Associated Endothelial Progenitors. Stem Cells, 2017, 35, 1328-1340. | 1.4 | 22 |
| 12 | Odd skipped-related 1 identifies a population of embryonic fibro-adipogenic progenitors regulating myogenesis during limb development. Nature Communications, 2017, 8, 1218. | 5.8 | 95 |
| 13 | Fibrogenic Potential of PW1/Peg3 Expressing Cardiac Stem Cells. Journal of the American College of Cardiology, 2017, 70, 728-741. | 1.2 | 27 |
| 14 | Expression Analysis of the Stem Cell Marker Pw1/Peg3 Reveals a CD34 Negative Progenitor Population in the Hair Follicle. Stem Cells, 2017, 35, 1015-1027. | 1.4 | 13 |
| 15 | Transplantation of Allogeneic PW1pos/Pax7neg Interstitial Cells EnhanceÂEndogenous Repair of InjuredÂPorcine Skeletal Muscle. JACC Basic To Translational Science, 2017, 2, 717-736. | 1.9 | 4 |
| 16 | The zinc finger transcription factor PW1/PEG3 restrains murine beta cell cycling. Diabetologia, 2016, 59, 1474-1479. | 2.9 | 5 |
| 17 | Fatty acid metabolism—the first trigger for cachexia?. Nature Medicine, 2016, 22, 584-585. | 15.2 | 4 |
| 18 | Phosphotyrosine phosphatase inhibitor bisperoxovanadium endows myogenic cells with enhanced muscle stem cell functions <i>via</i> epigenetic modulation of Scaâ€1 and Pw1 promoters. FASEB Journal, 2016, 30, 1404-1415. | 0.2 | 6 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Resident PW1 ⁺ Progenitor Cells Participate in Vascular Remodeling During Pulmonary Arterial Hypertension. Circulation Research, 2016, 118, 822-833. | 2.0 | 34 |
| 20 | A Novel Mutant Allele of Pw1/Peg3 Does Not Affect Maternal Behavior or Nursing Behavior. PLoS Genetics, 2016, 12, e1006053. | 1.5 | 22 |
| 21 | PW1/Peg3 expression regulates key properties that determine mesoangioblast stem cell competence. Nature Communications, 2015, 6, 6364. | 5.8 | 120 |
| 22 | The extraocular muscle stem cell niche is resistant to ageing and disease. Frontiers in Aging Neuroscience, 2014, 6, 328. | 1.7 | 28 |
| 23 | Porcine Skeletal Muscle-Derived Multipotent PW1pos/Pax7negInterstitial Cells: Isolation, Characterization, and Long-Term Culture. Stem Cells Translational Medicine, 2014, 3, 702-712. | 1.6 | 17 |
| 24 | Nâ€ <scp>WASP</scp> is required for Amphiphysinâ€2/ <scp>BIN</scp> 1â€dependent nuclear positioning and triad organization in skeletal muscle and is involved in the pathophysiology of centronuclear myopathy. EMBO Molecular Medicine, 2014, 6, 1455-1475. | 3.3 | 87 |
| 25 | Defining skeletal muscle resident progenitors and their cell fate potentials. Development (Cambridge), 2013, 140, 2879-2891. | 1.2 | 139 |
| 26 | Fibroadipogenic progenitors mediate the ability of HDAC inhibitors to promote regeneration in dystrophic muscles of young, but not old Mdx mice. EMBO Molecular Medicine, 2013, 5, 626-639. | 3.3 | 201 |
| 27 | An Unbiased Assessment of the Role of Imprinted Genes in an Intergenerational Model of Developmental Programming. PLoS Genetics, 2012, 8, e1002605. | 1.5 | 105 |
| 28 | Loss of a single allele for Ku80 leads to progenitor dysfunction and accelerated aging in skeletal muscle. EMBO Molecular Medicine, 2012, 4, 910-923. | 3.3 | 35 |
| 29 | Stem cells in the hood: the skeletal muscle niche. Trends in Molecular Medicine, 2012, 18, 599-606. | 3.5 | 106 |
| 30 | <i>PW1</i> gene/paternally expressed gene 3 (PW1/Peg3) identifies multiple adult stem and progenitor cell populations. Proceedings of the National Academy of Sciences of the United States of America, 2011, 108, 11470-11475. | 3.3 | 84 |
| 31 | Identification and characterization of a non-satellite cell muscle resident progenitor during postnatal development. Nature Cell Biology, 2010, 12, 257-266. | 4.6 | 390 |
| 32 | Skeletal Muscle Phenotypically Converts and Selectively Inhibits Metastatic Cells in Mice. PLoS ONE, 2010, 5, e9299. | 1.1 | 26 |
| 33 | A parable about environment and our daughters' health. Trends in Endocrinology and Metabolism, 2010, 21, 335-336. | 3.1 | 1 |
| 34 | Non-canonical Wnt signaling regulates cell polarity in female reproductive tract development via van gogh-like 2. Development (Cambridge), 2009, 136, 1559-1570. | 1.2 | 63 |
| 35 | Effects of p21 deletion in mouse models of premature aging. Cell Cycle, 2009, 8, 2002-2004. | 1.3 | 11 |
| 36 | Modulation of Caspase Activity Regulates Skeletal Muscle Regeneration and Function in Response to Vasopressin and Tumor Necrosis Factor. PLoS ONE, 2009, 4, e5570. | 1.1 | 39 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Tumor Necrosis Factor-Î \pm Inhibition of Skeletal Muscle Regeneration Is Mediated by a Caspase-Dependent Stem Cell Response. Stem Cells, 2008, 26, 997-1008. | 1.4 | 65 |
| 38 | PCBs Exert an Estrogenic Effect through Repression of the Wnt7a Signaling Pathway in the Female Reproductive Tract. Environmental Health Perspectives, 2006, 114, 898-904. | 2.8 | 59 |
| 39 | Muscle cachexia is regulated by a p53-PW1/Peg3-dependent pathway. Genes and Development, 2006, 20, 3440-3452. | 2.7 | 104 |
| 40 | Tumor necrosis factor-α gene transfer induces cachexia and inhibits muscle regeneration. Genesis, 2005, 43, 120-128. | 0.8 | 113 |
| 41 | Diethylstilbestrol exposure in utero: A paradigm for mechanisms leading to adult disease. Birth Defects Research Part A: Clinical and Molecular Teratology, 2005, 73, 133-135. | 1.6 | 21 |
| 42 | Embryonic deregulation of muscle stress signaling pathways leads to altered postnatal stem cell behavior and a failure in postnatal muscle growth. Developmental Biology, 2005, 281, 171-183. | 0.9 | 36 |
| 43 | Wnt5a is required for proper epithelial-mesenchymal interactions in the uterus. Development (Cambridge), 2004, 131, 2061-2072. | 1.2 | 216 |
| 44 | Wnt7a Is a Suppressor of Cell Death in the Female Reproductive Tract and Is Required for Postnatal and Estrogen-Mediated Growth1. Biology of Reproduction, 2004, 71, 444-454. | 1.2 | 54 |
| 45 | A role for Wnt/β-catenin signaling in lens epithelial differentiation. Developmental Biology, 2003, 259, 48-61. | 0.9 | 125 |
| 46 | Wnt signaling mediates reorientation of outer hair cell stereociliary bundles in the mammalian cochlea. Development (Cambridge), 2003, 130, 2375-2384. | 1.2 | 183 |
| 47 | The Receptor Tyrosine Kinase Regulator Sprouty1 Is a Target of the Tumor Suppressor WT1 and Important for Kidney Development. Journal of Biological Chemistry, 2003, 278, 41420-41430. | 1.6 | 72 |
| 48 | FAP-1 Association with Fas (Apo-1) Inhibits Fas Expression on the Cell Surface. Molecular and Cellular Biology, 2003, 23, 3623-3635. | 1.1 | 100 |
| 49 | Induction of Homologue of Slimb Ubiquitin Ligase Receptor by Mitogen Signaling. Journal of Biological Chemistry, 2002, 277, 36624-36630. | 1.6 | 48 |
| 50 | TNFalpha inhibits skeletal myogenesis through a PW1-dependent pathway by recruitment of caspase pathways. EMBO Journal, 2002, 21, 631-642. | 3.5 | 93 |
| 51 | Cellular and cis-regulation of En-2 expression in the mandibular arch. Mechanisms of Development, 2002, 111, 125-136. | 1.7 | 16 |
| 52 | Expression of theboc gene during murine embryogenesis. Developmental Dynamics, 2002, 223, 379-388. | 0.8 | 43 |
| 53 | Siah ubiquitin ligase is structurally related to TRAF and modulates TNF-α signaling. Nature Structural Biology, 2002, 9, 68-75. | 9.7 | 129 |
| 54 | Stress-induced decrease in TRAF2 stability is mediated by Siah2. EMBO Journal, 2002, 21, 5756-5765. | 3.5 | 109 |

4

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 55 | A Role for Engrailed-2 in Determination of Skeletal Muscle Physiologic Properties. Developmental Biology, 2001, 231, 175-189. | 0.9 | 28 |
| 56 | Identification of a Novel Stretch-Responsive Skeletal Muscle Gene (Smpx). Genomics, 2001, 72, 260-271. | 1.3 | 44 |
| 57 | Developmental expression pattern of thecdo gene. Developmental Dynamics, 2000, 219, 40-49. | 0.8 | 46 |
| 58 | The emergence of molecular gynecology: homeobox and Wnt genes in the female reproductive tract. BioEssays, 2000, 22, 902-910. | 1.2 | 57 |
| 59 | Pw1/Peg3 is a potential cell death mediator and cooperates with Siah1a in p53-mediated apoptosis. Proceedings of the National Academy of Sciences of the United States of America, 2000, 97, 2105-2110. | 3.3 | 151 |
| 60 | ldentification of Ankrd2, a Novel Skeletal Muscle Gene Coding for a Stretch-Responsive Ankyrin-Repeat Protein. Genomics, 2000, 66, 229-241. | 1.3 | 115 |
| 61 | Wnt genes and endocrine disruption of the female reproductive tract: a genetic approach. Molecular and Cellular Endocrinology, 1999, 158, 1-5. | 1.6 | 49 |
| 62 | Fetal exposure to DES results in de-regulation of Wnt7a during uterine morphogenesis. Nature Genetics, 1998, 20, 228-230. | 9.4 | 146 |
| 63 | Peg3/Pw1 is an imprinted gene involved in the TNF-NFκB signal transduction pathway. Nature Genetics, 1998, 18, 287-291. | 9.4 | 148 |
| 64 | Msh homeobox genes regulate cadherin-mediated cell adhesion and cell–cell sorting. , 1998, 70, 22-28. | | 28 |
| 65 | Differential expression patterns of Wnt genes in the murine female reproductive tract during development and the estrous cycle. Mechanisms of Development, 1998, 76, 91-99. | 1.7 | 150 |
| 66 | CDO, A Robo-related Cell Surface Protein that Mediates Myogenic Differentiation. Journal of Cell Biology, 1998, 143, 403-413. | 2.3 | 86 |
| 67 | Msx2 Is a Transcriptional Regulator in the BMP4-Mediated Programmed Cell Death Pathway. Developmental Biology, 1997, 186, 127-138. | 0.9 | 143 |
| 68 | Pw1, a Novel Zinc Finger Gene Implicated in the Myogenic and Neuronal Lineages. Developmental Biology, 1996, 177, 383-396. | 0.9 | 135 |
| 69 | MSX1 inhibits MyoD expression in fibroblast × 10T½ cell hybrids. Cell, 1995, 82, 611-620. | 13.5 | 141 |
| 70 | Ectoderm-Mesenchyme and Mesenchyme-Mesenchyme Interactions Regulate Msx-1 Expression and Cellular Differentiation in the Murine Limb Bud. Developmental Biology, 1995, 168, 374-382. | 0.9 | 100 |
| 71 | Restricted expression of type-II TGFβ receptor in murine embryonic development suggests a central role in tissue modeling and CNS patterning. Mechanisms of Development, 1995, 52, 275-289. | 1.7 | 42 |
| 72 | Myogenic Regulatory Factors: Dissecting Their Role and Regulation during Vertebrate Embryogenesis. Developmental Biology, 1993, 156, 11-23. | 0.9 | 146 |

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 73 | [24] Detection of messenger RNA by in Situ hybridization. Methods in Enzymology, 1993, 225, 384-404. | 0.4 | 118 |
| 74 | Hox Genes: A Role for Tissue Development. American Journal of Respiratory Cell and Molecular Biology, 1992, 7, 1-2. | 1.4 | 2 |
| 75 | Loss of N-myc function results in embryonic lethality and failure of the epithelial component of the embryo to develop Genes and Development, 1992, 6, 2235-2247. | 2.7 | 328 |
| 76 | In-Situ Hybridization of Tropoelastin mRNA during the Development of the Multilayered Neonatal Rat Aortic Smooth Muscle Cell Culture. Matrix Biology, 1992, 12, 321-332. | 1.8 | 19 |
| 77 | A transgene target for positional regulators marks early rostrocaudal specification of myogenic lineages. Cell, 1992, 69, 79-93. | 13.5 | 79 |
| 78 | Molecular aspects of regeneration in developing vertebrate limbs. Developmental Biology, 1992, 152, 37-49. | 0.9 | 94 |
| 79 | Expression of Hox-7.1 in myoblasts inhibits terminal differentiation and induces cell transformation. Nature, 1992, 360, 477-481. | 13.7 | 215 |
| 80 | Multiple sites of HOX-7 expression during mouse embryogenesis: Comparison with retinoic acid receptor mRNA localization. Molecular Reproduction and Development, 1992, 32, 303-314. | 1.0 | 55 |
| 81 | ld expression during mouse development: A role in morphogenesis. Developmental Dynamics, 1992, 194, 222-230. | 0.8 | 56 |
| 82 | Myogenesis in the Mouse. Novartis Foundation Symposium, 1992, 165, 111-131. | 1.2 | 12 |
| 83 | Expression of the muscle regulatory factor MRF4 during somite and skeletal myofiber development. Developmental Biology, 1991, 147, 144-156. | 0.9 | 274 |
| 84 | The expression of myosin genes in developing skeletal muscle in the mouse embryo Journal of Cell Biology, 1990, 111, 1465-1476. | 2.3 | 259 |
| 85 | Developmental regulation of myosin gene expression in mouse cardiac muscle Journal of Cell Biology, 1990, 111, 2427-2436. | 2.3 | 381 |
| 86 | Expression of two myogenic regulatory factors myogenin and MyoDl during mouse embryogenesis. Nature, 1989, 341, 303-307. | 13.7 | 647 |
| 87 | Myogenin, a factor regulating myogenesis, has a domain homologous to MyoD. Cell, 1989, 56, 607-617. | 13.5 | 1,248 |
| 88 | The protein encoded by a murine male germ cell-specific transcript is a putative ATP-dependent RNA helicase. Cell, 1989, 57, 549-559. | 13.5 | 167 |
| 89 | A developmental study of the abnormal expression of α-cardiac and α-skeletal actins in the striated muscle of a mutant mouse. Developmental Biology, 1989, 134, 236-245. | 0.9 | 36 |
| 90 | Development and hormone regulation of androgen receptor levels in the sexually dimorphic larynx of Xenopus laevis. Developmental Biology, 1989, 131, 111-118. | 0.9 | 51 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 91 | Androgen-induced myogenesis and chondrogenesis in the larynx of Xenopus laevis. Developmental Biology, 1986, 113, 135-140. | 0.9 | 71 |
| 92 | The sexually dimorphic larynx ofXenopus laevis: Development and androgen regulation. American Journal of Anatomy, 1986, 177, 457-472. | 0.9 | 108 |